Seeking and reporting of adverse effects in orthodontic research

Pauline A.J. Steegmans

Seeking and reporting of adverse effects in orthodontic research

Pauline Antoinette Josephine Steegmans

Financial support for printing and distribution of this dissertation was kindly provided by:







PS

ISBN: 978-94-6483-668-4 Lay-out: Ridderprint & Pauline Steegmans Cover design: Illustration by Pauline Steegmans. 'Class I molar-occlusion' Printing: Ridderprint | www.ridderprint.nl

Copyright 2024 © P.A.J. Steegmans, Amsterdam, The Netherlands.

All rights reserved. No parts of this dissertation may be reported or transmitted in any form or by any means without permission of the author.

Seeking and reporting of adverse effects in orthodontic research

ACADEMISCH PROEFSCHRIFT

ter verkrijging van de graad van doctor aan de Universiteit van Amsterdam op gezag van de Rector Magnificus prof. dr. ir. P.P.C.C. Verbeek ten overstaan van een door het College voor Promoties ingestelde commissie, in het openbaar te verdedigen in de Aula der Universiteit op woensdag 14 februari 2024, te 11.00 uur

> door Pauline Antoinette Josephine Steegmans geboren te Leiderdorp

Promotiecommissie

Promotor:	prof. dr. J. de Lange	Universiteit van Amsterdam
Copromotores:	dr. R.A. Meursinge Reynders dr. R.E.G. Jonkman	Universiteit van Amsterdam Vrije Universiteit Amsterdam
Overige leden:	prof. dr. A.G. Becking prof. dr. F. Abbas prof. dr. L. Dubois prof. dr. F.R. Rozema dr. F.N. van der Weijden dr. M.A. Jongsma	Universiteit van Amsterdam Rijksuniversiteit Groningen Universiteit van Amsterdam Universiteit van Amsterdam Universiteit van Amsterdam Jongsma & Jongsma

Faculteit der Tandheelkunde

Paranymphs

Michiel Steegmans

Floor Tielbeke

Dedication

This dissertation is affectionately dedicated to my beloved father (in memoriam), mother and brother.

Highlights of this dissertation

What is new?

Key findings

This dissertation summarized the results of 7 publications on adverse effects of orthodontic interventions. Key findings are:

- A new framework was developed to categorize and define adverse effects of orthodontic interventions.
- Relapse and stability issues (19% (36/195)) and undesired treatment results (22% (43/195)) were the predominant adverse effects sought and reported in systematic reviews of orthodontic interventions, but many adverse effects were underassessed and underreported.
- 36% (35/98) of systematic reviews of orthodontic interventions defined seeking of adverse effects of interventions as a research objective of the systematic review.
- 85% (83/98) of systematic reviews of orthodontic interventions reported findings related to adverse effects of interventions sought in the studies included in the review.
- 91% (89/98) of systematic reviews of orthodontic interventions considered, discussed (weighed) potential adverse effects of interventions somewhere in the manuscript.
- 77% (75/98) of systematic reviews of orthodontic interventions reported or considered (i.e., discussed, weighed) potential adverse effects of interventions in the abstracts.
- 41% (40/98) of systematic reviews of orthodontic interventions had spin on adverse effects in the abstracts of these reviews.
- Misleading reporting was the predominant type (90% (36/40)) of spin on adverse effects in the abstracts of systematic reviews of orthodontic interventions.

What this adds to what was known?

- This dissertation presented a new framework for categorizing and defining adverse effects of orthodontic interventions.
- This dissertation showed that most systematic reviews of orthodontic interventions assessed and reported adverse effects but these actions were not systematic, incomplete, and selective.
- This dissertation showed that reporting on adverse effects in abstracts of systematic reviews of orthodontic interventions was suboptimal with a high prevalence of spin.

What is the implication and what should change now

- This dissertation showed that what is assessed and reported on adverse effects in systematic reviews of orthodontic interventions was often incomplete and misleading, which could lead to inadequate clinical decision making.
- Focus on developing, assessing, and reporting of core adverse effects of orthodontic interventions (using our new framework) in primary studies and in systematic reviews.
- Besides conducting traditional systematic reviews of interventions, consider undertaking systematic reviews that focus exclusively on adverse effects.

Table of contents

Title page3
Members of the PhD committee4
Highlights of this dissertation7
Chapter 1
Chapter 2
Chapter 3
Chapter 4
Chapter 5
Chapter 6
Chapter 7
Chapter 8
Chapter 9
Chapter 10155 Summary and conclusions
Chapter 11

Chapter 12
List of publications168
List of contributing authors169
Author's contributions170
Funding172
Conflict of interest172
Data sharing statement172
Reprints and Permission173
Acknowledgment174
Curriculum vitae175
Chapter 13
Additional files chapter 2178
Additional files chapter 3187
Additional files chapter 4229
Additional files chapter 5237
Additional files chapter 8250



CHAPTER 1

General introduction and objectives

The central story

Systematic reviews identify the current knowledge status on a particular health issue by synthesizing and appraising best evidence. When well conducted, these reviews save end-users considerable time, energy, and resources. However, when done poorly, these reviews can be misleading and even cause harm. This can be worrisome considering the high-quality status of systematic reviews in the evidence hierarchy [1] and the high and increasing number of such reviews published in the literature (Figure 1).




Figure 1. Line chart showing the exponential growth of systematic reviews of interventions in healthcare since 2000 in PubMed.

Search strategy in PubMed: (systematic review* AND (healthcare* OR health* OR intervention* OR therapy* OR treatment*))

The consequences of low-quality reviews could be even more damaging when the adverse effects of interventions are poorly assessed and reported. Therefore prior to implementing the findings of systematic reviews, clinicians need to critically appraise these studies and evaluate their credibility regarding the assessment and reporting of adverse effects of interventions. Cochrane states that "it is critical that outcomes used to assess adverse effects as well as outcomes used to assess beneficial effects are among those addressed by a review" [2]. It is also critical that when presenting findings on adverse effects that they are free of spin, i.e., a distorted presentation of study results [3]. This is particularly important in abstracts, because titles and abstracts are the most and often the only read sections of biomedical papers [4]. This dissertation assessed a broad spectrum of items on defining, seeking and reporting of adverse effects in systematic reviews of orthodontic interventions.

Description of the condition

Classifying and defining adverse effects of interventions is challenging [5,6]. For instance, harms, adverse events, side effects, complications, safety, and toxicity are closely related terms, but their definitions can vary between studies [7]. The definition of adverse effects depends on the context and type of intervention [6]. For example, in orthodontics, anchorage loss of first molars that are reinforced with implants can be considered an adverse effect. Instead, anchorage loss can also be a beneficial effect, e.g., in extraction cases where anchorage loss is desired to avoid over-retraction of maxillary incisors. In this dissertation, we adopted Cochrane's definition of an adverse effect: "An adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility" [5].

Preoteasa et al. [8] categorized a variety of adverse effects of orthodontic interventions and presented them in a framework of subgroups (Table 1). Root resorption is an example of the dental subgroup of adverse effects linked to orthodontics and is illustrated in Figure 2. The framework by Preoteasa et al. [8] was used for defining and assessing adverse effects in the research studies of this dissertation. These assessments subsequently resulted in a new framework for defining adverse effects of orthodontic interventions, which is presented in additional file 2 of chapter 3, additional file 2 of chapter 9.




Figure 2. Root resorption is visible on a large number of maxillary and mandibular teeth [9]

Subgroup	Description
Local adverse effects	
Dental	Crown: decalcifications, decays, tooth wear, enamel cracks and fractures; discolorations, deterioration of prosthetic crown (as fracturing a ceramic one during debonding) Root: root resorption, early closure of root apex, ankylosis Pulp: ischemia, pulpitis, necrosis
Periodontal	 Gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscencer fenestrations, interdental fold, dark triangles
Temporomandibular joint	 Condylar resorption, temporomandibular dysfunction
Soft tissues of the oral and maxillofacial region	 Trauma (e.g., long archwires, headgear related), mucosal ulcerations or hyperplasia, chemical burns (e.g., etching related), thermal injuries (e.g., overheated burs), stomatitis, clumsy handling of dental instruments
Unsatisfactory treatment outcome	 Inadequate morpho-functional, esthetic or functional final result, relapse, failure to complete treatment due to treatment dropout
Systemic adverse effects	
Psychological	 Teasing, behavioral changes of patients and parents; disconfort associated with pain presence and esthetic look discontents during orthodontic appliance usage
Gastro-intestinal	 Accidental swallowing of small parts of the orthodontic device (tubes, brackets)
Allergies	To nickel or latex
Cardiac	Infective endocarditis
Chronic fatigue syndrome	
Cross infections	 From doctor to patient, patient to doctor, patient to patient

Table 1. Adverse effects hypothetically linked to orthodontic interventions according to Preoteasa et al. [8]*

*Permission to reproduce this table was obtained on 16 August 2018 from InTech's Publishing Ethics and Legal Affairs Department

Description of the problems and what is published on the problems

Three major methodological and reporting problems regarding adverse effects in systematic reviews can jeopardize the information published on these effects, which can subsequently affect clinical decision making.

1. Poor seeking, reporting, and synthesizing of adverse effects of interventions in systematic reviews

Epidemiological research has shown that seeking, reporting, and synthesizing of adverse effects of interventions was suboptimal in systematic reviews of interventions [10, 11, 12, 13, 14]. Poor methodological rigor in assessing adverse effects of interventions was also identified in Cochrane reviews [15]. The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) harms checklist was published in 2016 to improve the reporting of adverse effects in systematic reviews [7] but little evidence is available on the consequences of this reporting guideline on the quality of reporting of adverse effects in systematic reviews [16].

2. Poor seeking, reporting, and synthesizing of adverse effects of interventions in primary studies

The lack of rigor in seeking and reporting of adverse effects in the primary studies that feed systematic reviews is another major problem that affects the validity on what is reported on adverse effects in these reviews. Multiple epidemiological studies have identified these limitations in primary studies [17, 18, 19, 20, 21, 22]. For example, clinical trials on the evaluation of the same drugs showed that different adverse effects were assessed and reported [23, 24]. Further, a higher number and range of adverse effects were reported in unpublished versions of clinical trials compared to the final published versions of these trials [18]. To improve the reporting of adverse effects in randomized controlled trials (RCTs), an extension of the Consolidated Standards of Reporting Trials (CONSORT) was developed in 2004 [25]. Several studies have investigated the effect of this guideline on reporting adverse effects, but only slight improvements were found [17, 19, 20, 26].

3. Misleading reporting, interpretation, and extrapolation (spin) of adverse effects in systematic reviews.

Misleading presentation of study results regarding adverse effects is a major problem that can affect clinical decision making. Distorted presentation of study results is called 'spin' [3]. A wide variety of terms and definitions have been used for spin in the medical literature. For example; misrepresentation [27, 28], distorted presentation [29], inappropriate extrapolation [28], overinterpretation, and misreporting [30]. Spin has been divided into three subcategories: 'misleading reporting', 'misleading interpretation', and 'misleading extrapolation' of study results [31]. This dissertation will adopt the definitions described by Lazarus et al. [32] for these three categories of spin. Table 2 gives an overview of these definitions and key terminology used in this dissertation.

Spin in the abstracts of research studies can be particularly harmful because titles and abstracts are the most read and often also the only read sections of biomedical papers [4]. The presence of spin in abstracts can influence readers' interpretation [27], which could result in inadequate decisions on healthcare interventions. The Methodological Expectations of Cochrane Intervention Reviews (MECIR) state: "The abstract of the review should aim to reflect a balanced summary of the benefits and harms of the intervention" [40]. Epidemiological studies have examined the presence of spin in abstracts of randomized controlled trials and systematic reviews in different research fields [27, 29, 41, 42, 43, 44]. Such studies identified spin in more than 50% of abstracts of medical RCTs [29, 43, 45], and in more than 30% of abstracts of systematic reviews of proximal humerus fractures treatments [46]. Similar findings were reported in orthodontic RCTs [42] and systematic reviews [41]. Spin on adverse effects in abstracts of systematic reviews of interventions could be even more damaging for clinical decision making, but its magnitude and consequences have not been assessed in the literature.

Table 2. Glossary of terms

Term	Definition
Systematic review	Cochrane [33] defines a systematic review as follows: 'A systematic review attempts to identify, appraise and synthesize all the empirical evidence that meets pre-specified eligibility criteria to answer a specific research question. Researchers conducting systematic reviews use explicit, systematic methods that are selected with a view aimed at minimizing bias, to produce more reliable findings to inform decision making.'
Intervention review	Cochrane [33] defines an intervention review as follows: 'Intervention reviews assess the benefits and harms of interventions used in healthcare and health policy.'
Orthodontic interventions	Steegmans et al. [34] defined orthodontic interventions as follows: 'Orthodontic interventions refer to the use of any type of orthodontic appliance that are used to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodontic treatment, for example retainers.'
Adverse effect	Cochrane [35, 36] defines an adverse effect as 'an adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility'.
Spin [3]	'Distorted presentation of study results'.
Spin [3]	'A misrepresentation of study results, regardless of motive (intentionally or unintentionally) that overemphasizes the beneficial effects of the intervention and overstates safety compared with that shown by the results'.
Spin [37]	'A specific intentional or unintentional reporting that fails to faithfully reflect the nature and range of findings and that could affect the impression the results produce in readers'
Spin [38]	'A specific reporting that fails to faithfully reflect the nature and range of findings and that could affect the impression that the results produce in readers, a way to distort science reporting without actually lying'
Misleading reporting related spin [32]	'Incomplete reporting of the study results that could be misleading for the reader'.
Misleading interpretation related spin [32]	Inadequate interpretation of the study results overestimating the beneficial effect of the intervention.
Misleading (inappropriate) extrapolation related spin [32]	'Inappropriate generalization of the study results by inadequate 1) extrapolation from the population, interventions or outcome actually assessed in the study to a larger population, different interventions or outcomes, or 2) inadequate implications for clinical practice.
Spin (in the abstract) on adverse effects of interventions [39]	Incomplete or inadequate reporting, interpretation, or extrapolation (or a combination of these variables) of findings on adverse effects of interventions in the abstract that could be misleading for the reader.

Why this study is necessary and for who, and what are the objectives of this paper

Above we reported on the magnitude and the problems associated with incomplete, misleading, and non-assessing and reporting of adverse effects in research studies. In this dissertation we assessed a series of these issues in systematic reviews of orthodontic interventions. Cross-sectional studies were developed, which addressed 12 research questions (Figures 3 and 4).




Figure 3. Flow diagram "seeking adverse effects of interventions in systematic reviews of orthodontic interventions"




Figure 4.a. Reporting or considering adverse effects of orthodontic interventions in the abstract **b**. Spin on adverse effects of orthodontics in the abstract

Our scoping searches showed that these research questions were not assessed previously in the orthodontic literature. Besides our cross-sectional studies, we also explored adverse effects of orthodontic interventions in two critical appraisals and a case report. A detailed list of all research objectives is reported under here. Patients, clinicians, researchers, editors, peer reviewers, guideline developers, policymakers, and research funders, can all benefit from the findings of this dissertation.

The objectives of this dissertation are:

- 1. To assess whether seeking adverse effects was defined as a research objective in systematic reviews of orthodontic interventions (chapters 2 and 3).
- To assess whether systematic reviews of orthodontic interventions sought any findings related to adverse effects of interventions in the included studies (chapters 2 and 3).
- 3. To assess whether systematic reviews of orthodontic interventions reported any findings related to adverse effects of interventions sought in the included studies (chapters 2 and 3).
- 4. To assess whether potential adverse effects of orthodontic interventions were considered, discussed (weighed) anywhere in systematic reviews of orthodontic interventions. (chapters 2 and 3).
- 5. To assess each type of adverse effect sought in systematic reviews of orthodontic interventions (chapters 2 and 3).
- 6. To assess whether potential adverse effects were reported or considered (i.e., discussed, weighed, etc.) in abstracts of the systematic reviews of orthodontic interventions (chapters 4 and 5).
- 7. To assess whether spin was identified on adverse effects of orthodontic interventions in abstracts of systematic reviews of orthodontic interventions (chapters 4 and 5).
- 8. To assess what type of spin was identified on adverse effects of orthodontic interventions in the abstract of systematic reviews of orthodontic interventions (chapters 4 and 5).
- 9. To critically appraise a systematic review that assessed the effect of fixed orthodontic retainers on periodontal health [47] (chapter 6).
- To critically appraise a randomized controlled trial that assessed the post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers [48] (chapter 7).
- 11. To assess a case with an adverse effect of an orthodontic intervention (chapter 8).

References

- Ioannidis JP. The Mass Production of Redundant, Misleading, and Conflicted Systematic Reviews and Meta-analyses. Milbank Q. 2016 Sep;94(3):485-514. doi: 10.1111/1468-0009.12210. PMID: 27620683; PMCID: PMC5020151.
- O'Connor D, Green S, Higgins JPT (editors). Chapter 5: Defining the review question and developing criteria for including studies. In: Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. [online] Available from: <u>www.cochrane-handbook.org</u>. (accessed August 20th 2018).
- 3. Haneef R, Yavchitz A, Ravaud P, Baron G, Oranksy I, Schwitzer G, Boutron I. Interpretation of health news items reported with or without spin: protocol for a prospective meta-analysis of 16 randomised controlled trials. BMJ Open. 2017 Nov 17;7(11):e017425. doi: 10.1136/bmjopen-2017-017425.
- Dogan RI, Murray GC, Névéol A, Lu Z. Understanding PubMed user search behavior through log analysis. Database (Oxford). 2009;2009:bap018. Epub 2009 Nov 27.
- Peryer G, Golder S, Junqueira D, Vohra S, Loke YK. Chapter 19: Adverse effects. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.3 (updated February 2022). Cochrane, 2022. Available from www.training.cochrane.org/handbook. (Accessed 8 August 2023)
- Qureshi R, Mayo-Wilson E, Li T, Summaries of harms in systematic reviews are unreliable Paper 1: An introduction to research on harms, *Journal of Clinical Epidemiology* (2021), doi: https://doi.org/10.1016/j.jclinepi.2021.10.023
- Zorzela L, Loke YK, Ioannidis JP, Golder S, Santaguida P, Altman DG, Moher D, Vohra S; PRISMA Harms Group. PRISMA harms checklist: improving harms reporting in systematic reviews. BMJ. 2016 Feb 1;352:i157. doi: 10.1136/bmj.i157.
- Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F. (editor). Orthodontics-Basic aspects and clinical considerations. March 9, 2012 under CC BY 3.0 license. www.intechopen.com. [online] Available from: https://cdn.intechopen.com/pdfs/31388/InTech-
 Didentification of the tribulation of tribulation of the tribulation of the tribulation of tribulation of tribulation of the tribulation of trib
 - <u>Risks_and_complicationsassociated_with_orthodontic_treatment.pdf</u> (Accessed 4 December 2021).
- 9. Steegmans PAJ, Jonkman REG, de Lange J. Fixed Flexible Spiral Wire Retainers and Unwanted Tooth movements: A Case Report. *Appl. Sci.* 2023, *13*(2), 922.
- Li L, Xu C, Deng K, Zhou X, Liu Z, Busse JW, Ren Y, Zou K, Sun X. The reporting of safety among drug systematic reviews was poor before the implementation of the PRISMA harms checklist. J Clin Epidemiol. 2019 Jan;105:125-135. doi: 10.1016/j.jclinepi.2018.09.014. Epub 2018 Sep 29. PMID: 30278212.
- 11. Mahady SE, Schlub T, Bero L, Moher D, Tovey D, George J, Craig JC. Side effects are incompletely reported among systematic reviews in gastroenterology. J Clin Epidemiol. 2015 Feb;68(2):144-53.
- 12. Qureshi R, Mayo-Wilson E, Rittiphairoj T, McAdams-DeMarco M, Guallar E, Li T. Summaries of harms in systematic reviews are unreliable Paper 2: Methods used to assess harms are neglected in systematic reviews of gabapentin, Journal of Clinical Epidemiology (2021), doi: https://doi.org/10.1016/j.jclinepi.2021.10.024
- Qureshi R, Mayo-Wilson E, Rittiphairoj T, McAdams-DeMarco M, Guallar E, Li T. Summaries of harms in systematic reviews are unreliable Paper 3: Given the same data sources, systematic reviews of gabapentin have different results for harms, Journal of Clinical Epidemiology (2021), doi:<u>https://doi.org/10.1016/i.jclinepi.2021.10.025</u>
- Zorzela L, Golder S, Liu Y, Pilkington K, Hartling L, Joffe A, Loke Y, Vohra S. Quality of reporting in systematic reviews of adverse events: systematic review. BMJ. 2014 Jan 8;348:f7668. doi: 10.1136/bmj.f7668. PMID: 24401468; PMCID: PMC3898583.
- Johansson M, Borys F, Peterson H, Bilamour G, Bruschettini M, Jørgensen KJ. Addressing harms of screening-A review of outcomes in Cochrane reviews and suggestions for next steps. J Clin Epidemiol. 2021 Jan;129:68-73. doi: 10.1016/j.jclinepi.2020.09.030. Epub 2020 Oct 1.PMID: 33010402
- Chang T, Cui Y, Zhang Y, Ma J, Tan J, Wang J. Reporting Quality of oral TCM Systematic Reviews Based on the PRISMA Harms checklist from 2013 to 2020. Evid Based Complement Alternat Med. 2023 Jan 24;2023:4612036. doi: 10.1155/2023/4612036. eCollection 2023.PMID: 36733845

- Bagul NB, Kirkham JJ. The reporting of harms in randomized controlled trials of hypertension using the CONSORT criteria for harm reporting. Clin Exp Hypertens. 2012;34(8):548-54. doi: 10.3109/10641963.2012.681724. Epub 2012 May 9.
- Golder S, Loke YK, Wright K, Norman G. Reporting of Adverse Events in Published and Unpublished Studies of Health Care Interventions: A Systematic Review. PLoS Med. 2016 Sep 20;13(9):e1002127. doi: 10.1371/journal.pmed.1002127. eCollection 2016 Sep.
- Hodkinson A, Kirkham JJ, Tudur-Smith C, Gamble C. Reporting of harms data in RCTs: a systematic review of empirical assessments against the CONSORT harms extension. BMJ Open. 2013 Sep 27;3(9):e003436. doi: 10.1136/bmjopen-2013-003436.
- Péron J, Maillet D, Gan HK, Chen EX, You B. Adherence to CONSORT adverse event reporting guidelines in randomized clinical trials evaluating systemic cancer therapy: a systematic review. J Clin Oncol. 2013 Nov 1;31(31):3957-63. doi: 10.1200/JCO.2013.49.3981. Epub 2013 Sep 23.
- Smith SM, Chang RD, Pereira A, Shah N, Gilron I, Katz NP, Lin AH, McDermott MP, Rappaport BA, Rowbotham MC, Sampaio C, Turk DC, Dworkin RH. Adherence to CONSORT harmsreporting recommendations in publications of recent analgesic clinical trials: an ACTTION systematic review. Pain. 2012 Dec;153(12):2415-21. doi: 10.1016/j.pain.2012.08.009. Epub 2012 Sep 15.
- Tang E, Ravaud P, Riveros C, Perrodeau E, Dechartres A. Comparison of serious adverse events posted at ClinicalTrials.gov and published in corresponding journal articles. BMC Med. 2015 Aug 14;13:189. doi: 10.1186/s12916-015-0430-4. PMID: 26269118; PMCID: PMC4535304.
- Mayo-Wilson E, Fusco N, Li T, Hong H, Canner JK, Dickersin K; MUDS investigators. Harms are assessed inconsistently and reported inadequately part 1: systematic adverse events. J Clin Epidemiol. 2019 Sep;113:20-27. doi: 10.1016/j.jclinepi.2019.04.022. Epub 2019 May 2. PMID: 31055175.
- Mayo-Wilson E, Fusco N, Li T, Hong H, Canner JK, Dickersin K; MUDS investigators. Harms are assessed inconsistently and reported inadequately Part 2: nonsystematic adverse events. J Clin Epidemiol. 2019 Sep;113:11-19. doi: 10.1016/j.jclinepi.2019.04.020. Epub 2019 May 2. PMID: 31055176.
- Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, Schulz K, Moher D; CONSORT Group. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med. 2004 Nov 16;141(10):781-8.
- Junqueira DR, Phillips R, Zorzela L, Golder S, Loke Y, Moher D, Ioannidis JPA, Vohra S. Time to improve the reporting of harms in randomized controlled trials. J Clin Epidemiol. 2021 Aug;136:216-220. doi: 10.1016/j.jclinepi.2021.04.020. Epub 2021 May 10. PMID: 33984494.
- Boutron I, Altman DG, Hopewell S, Vera-Badillo F, Tannock I, Ravaud P. Impact of spin in the abstracts of articles reporting results of randomized controlled trials in the field of cancer: the SPIIN randomized controlled trial. J Clin Oncol. 2014 Dec 20;32(36):4120-6. doi: 10.1200/JCO.2014.56.7503. Epub 2014 Nov 17. PMID: 25403215.
- Yavchitz A, Boutron I, Bafeta A, Marroun I, Charles P, Mantz J, Ravaud P. Misrepresentation of randomized controlled trials in press releases and news coverage: a cohort study. PLoS Med. 2012;9(9):e1001308. doi: 10.1371/journal.pmed.1001308. Epub 2012 Sep 11. PMID: 22984354; PMCID: PMC3439420.
- Boutron I, Dutton S, Ravaud P, Altman DG. Reporting and interpretation of randomized controlled trials with statistically nonsignificant results for primary outcomes. JAMA. 2010 May 26;303(20):2058-64. doi: 10.1001/jama.2010.651.
- Ochodo EA, de Haan MC, Reitsma JB, Hooft L, Bossuyt PM, Leeflang MM. Overinterpretation and misreporting of diagnostic accuracy studies: evidence of "spin". Radiology. 2013 May;267(2):581-8. doi: 10.1148/radiol.12120527. Epub 2013 Jan 29.
- Haneef R, Lazarus C, Ravaud P, Yavchitz A, Boutron I. Interpretation of Results of Studies Evaluating an Intervention Highlighted in Google Health News: A Cross-Sectional Study of News. PLoS One. 2015 Oct 16;10(10):e0140889. doi: 10.1371/journal.pone.0140889. eCollection 2015.
- Lazarus C, Haneef R, Ravaud P, Boutron I. Classification and prevalence of spin in abstracts of nonrandomized studies evaluating an intervention. <u>BMC Med Res Methodol.</u> 2015 Oct 13;15:85. doi: 10.1186/s12874-015-0079-x.
- Cochrane library. Available from: https://www.cochranelibrary.com/ about/about-cochrane-reviews. Accessed 27 June 2023.
- Steegmans PAJ, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study. Syst Rev. 2019 Apr 5;8(1):89. doi: 10.1186/s13643-019-1000-1. PMID: 30953538; PMCID: PMC6449933.

- 35. Glossary of terms in the Cochrane Collaboration. Version 4.2.5. Updated May 2005. Available from: http://aaz.hr/resources/pages/57/7.%20Coc hrane%20glossary.pdf. Accessed 4 Dec 2021.
- Peryer G, Golder S, Junqueira D, Vohra S, Loke YK. Chapter 19: Adverse effects. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021). Cochrane, 2021. Available from <u>www.training.cochrane.org/handbook</u>.
- 37. Boutron I, Ravaud P. Misrepresentation and distortion of research in biomedical literature. <u>Proc Natl</u> <u>Acad Sci U S A.</u> 2018 Mar 13;115(11):2613-2619. doi: 10.1073/pnas.1710755115.
- Boutron I. Spin in Scientific Publications: A Frequent Detrimental Research Practice. Ann Emerg Med. 2020 Mar;75(3):432-434. doi: 10.1016/j.annemergmed.2019.11.002. Epub 2019 Dec 23. PMID: 31874770.
- 39. Steegmans, P.A.J., Di Girolamo, N, Meursinge Reynders, R.A. Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews. *Res Integr Peer Rev* 4, 27 (2019). https://doi.org/10.1186/s41073-019-0084-4
- 40. Higgins JPT, Lasserson T, Chandler J, Tovey D, Thomas, J, Flemyng E, Churchill R. Methodological Expectations of Cochrane Intervention Reviews. Cochrane: London, Version February 2022
- Makou O, Eliades T, Koletsi D. Reporting, interpretation, and extrapolation issues (SPIN) in abstracts of orthodontic meta-analyses published from 2000 to 2020. Eur J Orthod. 2021 Mar 19:567-575. doi: 10.1093/ejo/cjab009. Epub ahead of print. PMID: 33740054.
- Guo F, Fang X, Li C, Qin D, Hua F, He H. The presence and characteristics of 'spin' among randomized controlled trial abstracts in orthodontics. Eur J Orthod. 2021 Oct 4;43(5):576-582. doi: 10.1093/ejo/cjab044. PMID: 34397084.
- Khan MS, Lateef N, Siddiqi TJ, Rehman KA, Alnaimat S, Khan SU, Riaz H, Murad MH, Mandrola J, Doukky R, Krasuski RA. Level and Prevalence of Spin in Published Cardiovascular Randomized Clinical Trial Reports With Statistically Nonsignificant Primary Outcomes: A Systematic Review. JAMA Netw Open. 2019 May 3;2(5):e192622. doi: 10.1001/jamanetworkopen.2019.2622. PMID: 31050775; PMCID: PMC6503494.
- Mathieu S, Giraudeau B, Soubrier M, Ravaud P. Misleading abstract conclusions in randomized controlled trials in rheumatology: comparison of the abstract conclusions and the results section. Joint Bone Spine. 2012 May;79(3):262-7. doi: 10.1016/j.jbspin.2011.05.008. Epub 2011 Jul 5. PMID: 21733728.
- 45. Wang D, Chen L, Wang L, Hua F, Li J, Li Y, Zhang Y, Fan H, Li W, Clarke M. Abstracts for reports of randomised trials of COVID-19 interventions had low quality and high spin. J Clin Epidemiol. 2021 Jul 2;139:107-120. doi: 10.1016/j.jclinepi.2021.06.027. Online ahead of print.PMID: 34224834
- 46. Jones C, Rulon Z, Arthur W, Ottwell R, Checketts J, Detweiler B, Calder M, Adil A, Hartwell M, Wright DN, Vassar M. Evaluation of spin in the abstracts of systematic reviews and meta-analyses related to the treatment of proximal humerus fractures. J Shoulder Elbow Surg. 2021 Jan 19:S1058-2746(21)00016-1. doi: 10.1016/j.jse.2020.11.026. Epub ahead of print. PMID: 33482369.
- Arn ML, Dritsas K, Pandis N, Kloukos D. The effects of fixed orthodontic retainers on periodontal health: A systematic review. Am J Orthod Dentofacial Orthop. 2020;157(2):156–164.e17. doi:10.1016/j.ajodo.2019.10.010
- Krämer A, Sjöström M, Apelthun C, Hallman M, Feldmann I. Post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers-a randomized controlled trial. Eur J Orthod 2022; DOI: 10.1093/ejo/cjac043.



CHAPTER 2

Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study

This is a non-edited version of the manuscript:

Steegmans PAJ, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study. Syst Rev. 2019 Apr 5;8(1):89.

Abstract

Background: Before implementing healthcare interventions, clinicians need to weigh the beneficial and adverse effects of interventions. However, a large body of evidence has demonstrated that seeking and reporting of adverse effects is suboptimal in clinical trials and in systematic reviews of interventions. This cross-sectional study will investigate the status of this problem in orthodontics. This study will assess whether adverse effects were sought and whether findings related to adverse effects were reported in systematic reviews of orthodontic interventions in the five leading orthodontic journals and in the Cochrane Database of Systematic Reviews.

Methods: Systematic reviews of clinical orthodontic interventions published between 01 August 2009 and 31 July 2019 in the five leading orthodontic journals and in the Cochrane Database will be included. Empty reviews will be excluded. The reporting of outcomes on adverse effects will not determine eligibility, i.e., reviews will not be excluded, because they did not report usable data. Study selection and data extraction will be conducted independently by two authors. Our primary outcome will be the prevalence of systematic reviews of orthodontic interventions that sought any findings related to adverse effects in the included studies. Additional prevalence statistics will be calculated on a series of items related to seeking of adverse effects in the eligible reviews. All statistics will be calculated for (1) all journals together, (2) the group of five orthodontic journals and the Cochrane Database of Systematic Reviews separately, and (3) each individual journal separately. Chisquare tests of independence will be used to compare these groups.

Discussion: This study will assess whether adverse effects were sought in systematic reviews of orthodontic interventions. This knowledge is important, because reviews that present an incomplete picture on adverse effects can have unfavorable consequences for the endusers. Also not reporting that no adverse effects were assessed in eligible studies included in a systematic review can mislead pertinent stakeholders. Our findings could have policy implications for making judgments on accepting or rejecting an intervention systematic review for publication, for example, by directing editors and peer-reviewers to adopt the various items on adverse effects defined in the MECIR standards and in the PRISMA harm checklist.

Keywords: Orthodontics, Reporting, Systematic review, Interventions, Adverse effect, Adverse event, Harm, Safety, Side effect, Patient-important outcomes

Background

Making balanced decisions on healthcare interventions requires reliable evidence on both their beneficial and adverse effects. In the Cochrane systematic reviews of interventions, it is therefore mandatory to seek both types of outcomes and include at least one undesirable outcome as a primary outcome measure [1, 2]. In both Cochrane and non-Cochrane reviews of orthodontic interventions, we will assess whether adverse effects were sought and whether findings related to adverse effects were reported.

Since its foundation in 1993, Cochrane has set the standard for medical research-synthesis publications [3]. Systematic reviews with or without meta-analyses are the core of such syntheses and are the foundations for evidence-based practice guidelines and policy. Cochrane reviews of interventions aim at including outcomes that are likely to be important for patients, clinicians, the general public, guideline developers, administrators, and policy makers [2]. Cochrane states: "It is critical that outcomes used to assess adverse effects as well as outcomes used to assess beneficial effects are among those addressed by a review" (chapter 5.4.1) [2]. This issue is important, because a balanced perspective of an intervention can only be obtained when both types of outcomes are assessed and reported with the same rigor. Cochrane has formulated the following definition of an adverse effect: "An adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility" [4, 5]. We adopted Cochrane's definitions of adverse effects, systematic reviews, and interventions reviews in this manuscript (Table 1) [4–6].

Numerous epidemiological studies have shown that adverse effects of interventions are often under-assessed or under-reported in primary research studies [7–11]. In addition, much information on adverse events remains unpublished and the number and range of these events are higher in unpublished compared to published versions of the same study [12]. To improve the reporting of harms in randomized trials, an extension of Consolidated Standards of Reporting Trials (CONSORT) Statement was developed [13]. The reporting of adverse events has improved over time since the publication of this extension, but was still suboptimal for a wide variety of clinical trials [9, 11, 14]. Systematic reviewers have an important role in bringing these issues to the foreground. However, epidemiological studies have shown that seeking and reporting adverse effects of interventions is also suboptimal in systematic reviews [15–18]. In 2016, the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) harm checklist [19] to improve harm reporting in systematic reviews was published, but the consequences of this checklist are still unknown.

In this study, we will assess whether adverse effects were sought and reported in systematic reviews of orthodontic interventions. We will scrutinize such reviews in the five leading orthodontic journals and those registered in the Cochrane Database of Systematic Reviews. Reporting on pain as a result of tooth movement and the various categories of known orthodontic adverse effects as defined by Preoteasa et al. [20] will be assessed in these reviews (Table 2). Scoping searches in the orthodontic literature confirmed the knowledge gaps on our research questions. Our pilot studies on intervention reviews of the Cochrane Database of Systematic Reviews and those published in the five leading orthodontic journals quantified these gaps and further showed the need to undertake this research study. Addressing our research objectives is crucial for patients, clinicians, researchers, policy

makers, and research sponsors. These questions are particularly important, because systematic reviews are increasingly consulted by patients [21].

Table 1 Glossary of terms

Term	Definition
Systematic review	The Cochrane glossary [5] defines a systematic review as "A review of a clearly formulated question that uses systematic and explicit methods to identify, select, and critically appraise relevant research, and to collect and analyse data from the studies that are included in the review. Statistical methods (meta-analysis) may or may not be used to analyse and summarise the results of the included studies."
Intervention review	Cochrane (5) defines an intervention review as follows: "Intervention reviews assess the benefits and harms of interventions used in healthcare and health policy."
Orthodontic interventions	Orthodontic interventions refer to the use of any type of orthodontic appliances that are used to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodontic treatment, for example retainers.
Adverse effect	Cochrane [4, 5] defines an adverse effect as "an adverse event for which the causal relation between the intervention and th event is at least a reasonable possibility."

Subgroup	Description
Local adverse effects	
Dental	 Crown: decalcifications, decays, tooth wear, enamel cracks and fractures; discolorations, deterioration of prosthetic crown (as fracturing a ceramic one during debonding) Root root resorption, early closure of root apex, ankylosis Pulp: ischemia, pulpitis, necrosis
Periodontal	 Gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscencer fenestrations, interdental fold, dark triangles
Temporomandibular joint	Condylar resorption, temporomandibular dysfunction
Soft tissues of the oral and maxilofacial region	 Trauma (e.g., long archwites, headgear related), mucosal ulcerations or hyperplasia, chemical burns (e.g., etching related), thermal injuries (e.g., overheated burs), stomatitis, clumsy handling of dental instruments
Unsatisfactory treatment outcome	 Inadequate morpho-functional, esthetic or functional final result, relapse, failure to complete treatment due to treatment dropout
Systemic adverse effects	
Psychological	 Teasing, behavioral changes of patients and parents; disconfort associated with pain presence and esthetic look discontents during orthodontic appliance usage
Gastro-intestinal	 Accidental swallowing of small parts of the orthodontic device (tubes, brackets)
Allergies	To nickel or latex
Cardiac	Infective endocarditis
Chronic fatigue syndrome	
Cross infections	· From doctor to patient, patient to doctor, patient to patient

Table 2 Adverse effects hypothetically linked to orthodontic interventions [20]

Permission to reproduce this table was obtained on 16 August 2018 from InTech's Publishing Ethics and Legal Affairs Department


Objectives

The main research question of this cross-sectional study is the following: "Do reviewers seek adverse effects in systematic reviews of orthodontic interventions?" To address this question, we have defined the following objectives:

- To calculate the prevalence of eligible systematic reviews of orthodontic interventions that defined seeking of adverse effects as a research objective of the review
- To calculate the prevalence of eligible systematic reviews of orthodontic interventions that sought any findings related to adverse effects in the included studies
- To calculate the prevalence of eligible systematic reviews of orthodontic interventions that considered and discussed (weighed) potential adverse effects of the intervention anywhere in the review
- To calculate the prevalence of each type of adverse effect sought in the review

Methods

We used the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P) 2015 statement as the guideline for reporting this protocol [22, 23]. The PRISMA-P checklist is included as Additional file 1. Figure 1 represents the flow diagram of our research methods. Our first step was to conduct scoping searches to identify knowledge gaps and prioritize research questions on seeking and reporting of adverse effects in systematic reviews of orthodontic interventions. Two reviewers (PS and RMR) subsequently conducted pilot tests to assess the validity of these questions and the research methods and to fine-tune them. The sample size for the pilot test was calculated a priori [24], and random numbers were generated to select pilot systematic reviews [25]. The procedures for our pilot tests are reported in Additional file 2. In the following sections, we presented our planned methods based on these pilot tests.



Eligibility criteria

Study designs

- We will include systematic reviews of orthodontic interventions. The definition of a systematic review, an intervention review, and orthodontic interventions listed in the Glossary of terms will be used to assess whether a review is eligible (Table 1).
- We will exclude (1) non-interventional reviews such as "Methodology," "Diagnostic," "Qualitative," and "Prognostic"; (2) rapid and scoping reviews; (3) systematic reviews that focus exclusively on adverse effects of interventions; and (4) systematic reviews of interventions that did not find any eligible studies (empty reviews).

Participants

- We will include systematic reviews on any type of patients undergoing orthodontic interventions, i.e., patients of any health status, sex, age, demographics, and socio-economic status.
- We will exclude (1) intervention reviews that focus exclusively on patients with congenital anomalies, for example, with cleft lip and palate, and (2) systematic reviews of animal or laboratory studies.

Interventions

- We will include the following: (1) Systematic reviews that assess the effects of clinical orthodontic interventions. Clinical orthodontic interventions refer to the use of any type of orthodontic appliances that are used to move teeth or change the jaw size or position for orthodontic purposes. (2) Systematic reviews of interventions with appliances to maintain or stabilize the results of orthodontic treatment, for example, retainers. (3) Systematic reviews of orthodontic interventions that compare the effects of orthodontic treatment with or without additional interventions such as pharmacological or small surgical procedures, e.g., periodontal or implant surgery.
- We will exclude (1) systematic reviews in which patients receive orthodontic treatment, but in which the effects of other interventions, e.g., periodontal surgery, are compared and not the effects of orthodontic interventions; (2) systematic reviews of interventions in which orthodontic appliances are specifically used for other purposes, e.g., changing jaw positions to treat respiration or temporomandibular disorders; and (3) systematic reviews of orthodontic interventions that included orthognathic surgery.
- No exclusion criteria will be applied to the characteristics of the operator who conducted the interventions.

Outcomes

- Any adverse effect of an orthodontic intervention scored at any endpoint or timing will be eligible.
- The effects of orthodontic interventions do not refer just to outcomes related to tooth and jaw size and positions, but also to broader outcomes such as periodontal

health, esthetic changes, the health of the temporomandibular joint, patient health experiences, and economic issues associated with the intervention.

• The reporting of outcomes on adverse effects will not determine the eligibility of reviews for this cross- sectional study, i.e., reviews will not be excluded because they did not provide "usable" data [2].

Setting

• No exclusion criteria will be applied to the type of setting, e.g., university or private practice, etc., in which the interventions were conducted.

Information sources

We will manually search eligible systematic reviews between 01 August 2009 and 31 July 2019 in the Cochrane Database of Systematic Reviews [26] and in the websites of the five leading orthodontic journals. We consulted the journal citation reports by Clarivate Analytics [27] to identify the five leading orthodontic journals based on their impact factor. Based on these reports, the following orthodontic journals were included: European Journal of Orthodontics [EJO], American Journal of Orthodontics, and Dentofacial Orthopedics [AJODO], Angle Orthodontist, The Korean Journal of Orthodontics, and Orthodontics and Craniofacial Research. Recently launched orthodontic journals, i.e., covering less than 10 years of journal publication, will not be eligible. The first of August 2009 was chosen as the incept data for our searches, because it coincides with the launch of the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement and guidance on 21 July 2009 [28, 29].

Study records

Data management

- All study selection and data extraction procedures will be conducted by two authors (PS and RMR) independently.
- Our pilot tests were also used to train both reviewers in applying our methods consistently and to calibrate them [23].
- Disagreement on the eligibility of a paper or the extraction of data will be resolved through (1) discussions between reviewers, (2) rereading the pertinent paper, or (3) contacting its authors by email [28]. Persistent disagreements will be resolved through the consultation of a methodologist (SB).
- All eligible systematic reviews will be downloaded as PDFs, and all data will be extracted to an Excel spreadsheet [30].

Selection process

- All titles and abstracts will be screened for eligibility in the websites of the five orthodontic journals. We will search the section "Dentistry and Oral health" for eligible reviews in the Cochrane Database of Systematic Reviews [26].
- When updates of reviews are identified, we will only consider the latest version.
- Authors suspect of multiple publications of the same systematic review will be contacted by email. We plan to consider the first publication, but this decision will be weighed on a case-by-case basis. Our rationale for these decisions will be reported in the completed study.
- A PRISMA flow diagram will illustrate our selection procedures [28, 29].
- All eligible and excluded systematic reviews will be presented in tables. The rationale for exclusion will be listed for each excluded review.

Data collection process

- Eligible studies and their pertinent supplemental files will be merged into binder PDFs, and multiple search terms will be applied to facilitate data extraction [31, 32].
- We consulted various articles on adverse effects [4, 13, 18, 19, 33, 34] and thesauri to develop these search terms. A table with all search terms is listed in Additional file 3.
- All pertinent data items will be extracted using our pilot tested data collection forms. These forms are presented in Additional file 4 and incorporate all our research questions. We consulted the PRISMA [28, 29] and the PRISMA-P [22, 23] checklists to develop these data collection forms.
- Criteria for scoring the pertinent data items are defined in these forms.
- We will search the entire eligible review, i.e., the text, tables, figures, and supplemental files. The plain language summary in eligible Cochrane systematic reviews will not be scrutinized for data items.
- Modifications made in the collection forms during data extraction will be reported in the section "Differences between the protocol and review" together with the rationale for these changes.

Scoring adverse effects of orthodontic interventions

- We will adopt a priori the various categories of known orthodontic adverse effects as defined by Preoteasa et al. [20], which were divided into two main types: local and systemic, with their pertinent subtypes (Table 2).
- We will also consider pain as a result of tooth movement and additional adverse effects of orthodontic interventions that are identified post hoc, i.e., during data extraction, and are not listed in Table 2. We will explain the rationale for including specific additional effects as adverse and will produce a framework for categorizing them.
- Ambiguous outcomes that could be interpreted as either beneficial or adverse will not be scored as "adverse." We will also present the rationale for this score.

Ambiguous outcomes will only be scored as adverse when the authors of the pertinent review define these outcomes as such.

Outcomes and statistical analyses

- All research questions are presented in flow diagrams (Fig. 2).
- All planned outcomes are presented in a summary of findings table (Table 3).
- All prevalence data will be calculated and reported with their 95% confidence levels.
- Prevalence statistics will be calculated for (1) all journals together, (2) the group of five orthodontic journals and the Cochrane Database of Systematic Reviews separately, and (3) each individual journal separately. Comparisons between these statistics will be calculated. These statistics will be compared with chi-square tests of independence. We will report the value of chi-square, the degrees of freedom (df), and the p value. A p value of < 0.05 will be considered to be statistically significant. We will use Stata software (Stata Corporation, College Station, TX, USA) version 15 for all the statistical analyses [35].
- We will report all outcomes that will be introduced or eliminated post hoc together with the rationale for inclusion or exclusion.




Table 3 Summary of findings		
Description of outcomes from the main text	Statistic	
The number of retrieved systematic reviews	Number	
The prevalence of eligible systematic reviews of orthodontic interventions	Prevalence	
The prevalence of eligible systematic reviews of orthodontic interventions that defined seeking of adverse effects of interventions as a research objective of the review	Prevalence	
The prevalence of eligible systematic reviews of orthodontic interventions that sought any findings related to adverse effects of interventions in the included studies	Prevalence	
The prevalence of eligible systematic reviews of orthodontic interventions that reported findings related to the adverse effects of interventions sought in the included studies	Prevalence	
The prevalence of eligible systematic reviews of orthodontic interventions that considered, discussed (weighed) potential adverse effects of the intervention anywhere in the review	Prevalence	
The prevalence of each type of adverse effect of interventions defined in the objectives of the review	Prevalence	
The prevalence of each type of adverse effect of interventions sought in the review	Prevalence	

Reporting of the research study and data management

- We will adopt The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement as the guideline for reporting the completed cross-sectional study [36].
- We prepared a data management plan for the long- term storage of our research data [37]. This plan guarantees that (1) all our project data will be made freely available and (2) our submitted article will be accompanied by additional files with all raw data of the completed study or with a link to a repository where these files will be deposited. In the latter case, we will register our repository in the Registry of Research Data Repositories [38]. (3) Our project data will be presented in a format that permits other scientists to understand, cite, and reuse the data. (4) Sensitive data will be protected. (5) Our data management plan will be frequently reassessed and updated if necessary [37, 38].

Differences between the protocol and the completed study

All differences between the protocol and the final research study will be reported together with the rationale for these changes. We will also present the consequences of these modifications on the magnitude, direction, and validity of the outcomes [39].

Discussion

Strengths

We point at four key strengths of this research study. First, extensive scoping searches and pilot studies were conducted to fine-tune our research questions and procedures. Our pilot studies also confirmed the importance of our research questions. Second, the research team consisted of two topic experts (PS and RMR) and two methodologists (RMR and SB). Third, all study selection and data extraction procedures were conducted by two operators (PS and RMR) independently. Fourth, this study will permit reproducibility, because we will publish the protocol a priori and all raw data of the completed study will be reported in additional files or will be deposited in an open access repository [37, 40].

Limitations

The limitations of this research study include the following: (1) It does not cover all journals that have published orthodontic intervention systematic reviews, but only a subgroup, i.e., those published in the five leading orthodontic journals and in the Cochrane Database of Systematic Reviews. However, we expect that the choice of this subgroup of the leading orthodontic literature will produce outcomes that will underestimate the true severity of the problem. (2) Including only systematic reviews of orthodontic interventions published in the last 10 years could introduce the risk of publication bias. However, we chose this period, because it will represent the actual knowledge status on assessing adverse effects in orthodontic intervention systematic reviews. Further, this period coincides with the launch in 2009 of the PRISMA reporting checklist, which is an important update on how to report items in systematic reviews [28, 29].

Importance and beneficiaries

In this research study, we will assess whether adverse effects were sought and reported in both Cochrane and non-Cochrane systematic reviews of orthodontic interventions. This is important, because of the following: (1) The validity of the findings of systematic reviews of interventions depends on a balanced presentation of both the benefits and adverse effects of the intervention [19]. (2) There is a large body of evidence that has demonstrated that seeking and reporting of adverse effects is suboptimal in a wide variety of clinical trials [7–11]. Systematic reviewers can have a crucial role as whistle blowers by bringing these knowledge gaps to the foreground. However, their position can also be damaging, because reviews that present an incomplete picture on these gaps can have unfavorable consequences for the end-users. For example, not reporting that no adverse effects were assessed in eligible studies included in a systematic review can mislead readers.

Our findings could have policy implications for making judgments on accepting or rejecting a systematic reviews of orthodontic interventions for publication, for example, by directing editors and peer-reviewers to adopt the various items on adverse effects defined in the Methodological Expectations of Cochrane Intervention Reviews (MECIR) standards [1] and the PRISMA harm checklist [19]. Patients, clinicians, researchers, editors, peer-reviewers,

guideline developers, policy makers, and research funders will all benefit from the findings of this research study.

Additional files

Additional file 1: Checklist for the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P) 2015 statement. (DOCX 33 kb) Additional file 2: Pilot tests. (DOCX 21 kb) Additional file 3: Search terms and their derivatives. (DOCX 15 kb) Additional file 4: Data collection forms. (DOCX 16 kb)

Abbreviations

EQUATOR Network: Enhancing the Quality and Transparency Of health Research Network; MECIR: Methodological Expectations of Cochrane Intervention Reviews; PRISMA: Preferred Reporting Items for Systematic reviews and Meta-Analyses; PRISMA-P: Preferred Reporting Items for Systematic review and Meta-Analysis-Protocols

Acknowledgements

Not applicable

Funding

All expenses for preparing this protocol and for conducting the subsequent research study will be paid evenly by each author.

Availability of data and materials

Not applicable

Authors' contributions

PS and RMR conceived and designed the study protocol for this cross- sectional study. RMR is the guarantor. PS and RMR conducted the pilot testing of the study selection procedures and data extraction forms and fine-tuned the research protocol after the pilot testing. SB provided support on methodological and statistical issues and assisted in the overall fine-tuning of this protocol. All authors read and approved the final protocol.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

Reint Meursinge Reynders and Shandra Bipat are both Associate Editors for Systematic Reviews. The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Orthodontics, Academisch Centrum Tandheelkunde Amsterdam (ACTA), University of Amsterdam, Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands.

²Department of Radiology, Academic Medical Center, University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.

³Studio di ortodonzia, Via Matteo Bandello 15, 20123 Milan, Italy.

⁴Department of Oral and Maxillofacial Surgery, Academic Medical Center, University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.

Received: 22 August 2018 Accepted: 25 March 2019

Published online: 05 april 2019

References

- Higgins JPT, Lasserson T, Chandler J, Tovey D, Churchill R. Methodological Expectations of Cochrane Intervention Reviews (MECIR). Standards for the conduct and reporting of new Cochrane Intervention Reviews, reporting of protocols and the planning, conduct and reporting of updates. London: Cochrane; 2016. [online] Available from: <u>https://community</u>.cochrane.org/ mecir-manual (Accessed 7 Dec 2018)
- O'Connor D, Green S, Higgins JPT. Chapter 5: defining the review question and developing criteria for including studies. In: JPT H, Green S, editors. Cochrane handbook for systematic reviews of interventions version 5.1.0 (updated March 2011): The Cochrane Collaboration; 2011. [online] Available from: <u>http://handbook</u>-5-1.cochrane.org/. (Accessed 20 Aug 2018).
- 3. Gurevitch J, Koricheva J, Nakagawa S, Stewart G. Meta-analysis and the science of research synthesis. Nature. 2018;555(7695):175–82.
- Loke YK, Price D, Herxheimer A. Chapter 14: adverse 38ochran. In: JPT H, Green S, editors. Cochrane handbook for systematic reviews of interventions version 5.1.0 (updated March 2011): The Cochrane Collaboration; 2011. [online] Available from: <u>http://handbook</u>-5-1. Cochrane.org/. (Accessed 20 Aug 2018).
- Glossary of terms in the Cochrane Collaboration. Version 4.2.5. Updated May 2005. [online] Available from: <u>http://aaz</u>.hr/resources/pages/57/7. %20Cochrane%20glossary.pdf (Accessed 20 Aug 2018).
- Cochrane library. [online] Available from: <u>https://www</u>.cochranelibrary.com/ about/about-cochranereviews (Accessed 20 Aug 2018).
- Bagul NB, Kirkham JJ. The reporting of harms in randomized controlled trials of hypertension using the CONSORT criteria for harm reporting. Clin Exp Hypertens. 2012;34(8):548–54 doi: 10.3109/10641963.2012.681724. Epub 2012 May 9.
- Hodkinson A, Kirkham JJ, Tudur-Smith C, Gamble C. Reporting of harms data in RCTs: a systematic review of empirical assessments against the CONSORT harms extension. BMJ Open. 2013;3(9):e003436. <u>https://doi.org/10.1136/bmjopen-2013-003436</u>.
- Péron J, Maillet D, Gan HK, Chen EX, You B. Adherence to CONSORT adverse event reporting guidelines in randomized clinical trials evaluating systemic cancer therapy: a systematic review. J Clin Oncol. 2013;31(31): 3957–63. https://doi.org/10.1200/JCO.2013.49.3981 Epub 2013 Sep 23.
- 10. Pitrou I, Boutron I, Ahmad N, Ravaud P. Reporting of safety results in published reports of randomized controlled trials. Arch Intern Med. 2009; 169(19):1756–61.
- Smith SM, Chang RD, Pereira A, Shah N, Gilron I, Katz NP, Lin AH, MP MD, Rappaport BA, Rowbotham MC, Sampaio C, Turk DC, Dworkin RH. Adherence to CONSORT harms-reporting recommendations in publications of recent analgesic clinical trials: an ACTTION systematic review. Pain. 2012;153(12):2415–21. <u>https://doi</u>.org/10.1016/j.pain.2012.08.009 Epub 2012 Sep 15.
- Golder S, Loke YK, Wright K, Norman G. Reporting of adverse events in published and unpublished studies of health care interventions: a systematic review. PloS Med. 2016;13(9):e1002127. <u>https://doi</u>.org/10.1371/ journal.pmed.1002127 eCollection 2016 Sep.
- Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, Schulz K, Moher D, CONSORT Group. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med. 2004;141(10):781–8.
- Haidich AB, Birtsou C, Dardavessis T, Tirodimos I, Arvanitidou M. The quality of safety reporting in trials is still suboptimal: survey of major general medical journals. J Clin Epidemiol. 2011;64(2):124–35. <u>https://doi.org/10.1016/j.jclinepi.2010.03.005 Epub 2010 Jun 17.</u>
- Hopewell S, Wolfenden L, Clarke M. Reporting of adverse events in systematic reviews can be improved: survey results. J Clin Epidemiol. 2008;61(6):597–602. https://doi.org/10.1016/j.jclinepi.2007.10.005 Epub 2008 Apr 14.
- Hammad TA, Neyarapally GA, Pinheiro SP, Iyasu S, Rochester G, Dal Pan G. Reporting of meta-analyses of randomized controlled trials with a focus on drug safety: an empirical assessment. Clin Trials. 2013;10(3):389–97. https:// doi.org/10.1177/1740774513479467. Epub 2013 Mar 18.
- Mahady SE, Schlub T, Bero L, Moher D, Tovey D, George J, Craig JC. Side effects are incompletely reported among systematic reviews in gastroenterology. J Clin Epidemiol. 2015;68(2):144–53.

- Zorzela L, Golder S, Liu Y, Pilkington K, Hartling L, Joffe A, Loke Y, Vohra S. Quality of reporting in systematic reviews of adverse events: systematic review. BMJ. 2014;348:f7668. <u>https://doi.org/10.1136/bmj.f7668 Review.</u>
- Zorzela L, Loke YK, Ioannidis JP, Golder S, Santaguida P, Altman DG, Moher D, Vohra S, PRISMA Harms Group. PRISMA harms checklist: improving harms reporting in systematic reviews. BMJ. 2016;352:i157. <u>https://doi.org/10.1136/ bmj.i157</u>.
- Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: risks and complications associated with orthodontic treatment. In: Bourzgui F, editor. Orthodontics- basic aspects and clinical considerations; 2012. Under CC BY 3.0 license. www. Intechopen.com. [online] Available from:
 <u>https://www</u>.intechopen.com/books/ orthodontics-basic-aspects-and-clinical-considerations/risks-and-complications- associated-with-orthodontic-treatment (Accessed 20 Aug 2018).
- 21. Reeves BC. Reporting of harms in systematic reviews and their primary studies. BMJ. 2014;349:g6819. <u>https://doi</u>.org/10.1136/bmj.g6819.
- 22. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. Preferred reporting items for systematic review and meta- analysis protocols (PRISMA-P) 2015 statement. Syst Rev. 2015;4(1):1.
- Shamseer L, Moher D, Clarke M, Ghersi D, Deceased LA, Petticrew M, Shekelle P, Stewart LA, the PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;349:g7647.
- Viechtbauer W, Smits L, Kotz D, Budé L, Spigt M, Serroyen J, Crutzen R. A simple formula for the calculation of sample size in pilot studies. J Clin Epidemiol. 2015;68(11):1375–9. <u>https://doi.org/10.1016/j.jclinepi.2015.04.014</u> Epub 2015 Jun 6.
- 25. Random.org. Randomness and Integrity Services Ltd. [online] Available from: <u>https://www</u>.random.org/ (Accessed 20 Aug 2018).
- 26. Cochrane library. Cochrane database of systematic reviews. [online] Available from: <u>https://www</u>.cochranelibrary.com (Accessed 20 Aug 2018).
- 27. Clarivate Analytics. [online] Available from: https://clarivate.com/(Accessed 20 Aug 2018).
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. PloS Med. 2009;6:e1000100 Epub 2009 Jul 21.
- 29. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PloS Med. 2009;6:e1000097.
- Microsoft Excel. [online] Available from: <u>https://products</u>.office.com/it-IT/ excel (Accessed 20 Aug 2018).
- Adobe Merge PDFs, combine files into one PDF. [online] Available from: <u>http://www</u>.wikihow.com/Merge-PDF-Files (Accessed 20 Aug 2018).
- Acrobat for legal professionals. Searching and marking multiple words. [online] Available from: <u>http://blogs</u>.adobe.com/acrolaw/2010/04/searching- and-marking-multiple-words-in-a-pdf/ (Accessed 20 Aug 2018).
- Golder S, Loke YK, Zorzela L. Some improvements are apparent in identifying adverse effects in systematic reviews from 1994 to 2011. J Clin Epidemiol. 2013;66(3):253–60. <u>https://doi.org/10.1016/j.jclinepi.2012.09.013</u>.
- Saini P, Loke YK, Gamble C, Altman DG, Williamson PR, Kirkham JJ. Selective reporting bias of harm outcomes within studies: findings from a cohort of systematic reviews. BMJ. 2014;349:g6501. <u>https://doi.org/10.1136/bmj.g6501.</u>
- 35. StataCorp. Stata statistical software: release 15. College Station: StataCorp LLC; 2017.
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, Initiative STROBE. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Ann Intern Med. 2007;147(8):573–7.
- Schiermeier Q. For the record. Making project data freely available for open science. Nature. 2018;555:403–5.
- Registry of Research Data Repositories. [online] Available from: <u>https://www</u>. Re3data.org/ (Accessed 20 Aug 2018).
- Higgins JPT, Deeks JJ, Altman DG (editors). Chapter 16: special topics in statistics. In: Higgins JPT, Green S (editors), Cochrane handbook for systematic reviews of interventions version 5.1.0 (updated March

2011). The Cochrane Collaboration, 2011. [online] Available from: <u>http://handbook</u>-5-1. 40ochrane.org/ (Accessed 20 Aug 2018).

 Goodman SN, Fanelli D, Ioannidis JP. What does research reproducibility mean? Sci Transl Med. 2016;8(341):341 ps12. <u>https://doi</u>.org/10.1126/ scitranslmed.aaf5027.



CHAPTER 3

Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1)

This is a non-edited version of the manuscript:

Steegmans PAJ, Di Girolamo N, Bipat S, Reynders RAM. Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1). Syst Rev. 2023 Jul 3;12(1):112.

Abstract

Background Systematic reviews that assess the benefits of interventions often do not completely capture all dimensions of the adverse effects. This cross-sectional study (part 1 of 2 studies) assessed whether adverse effects were sought, whether the findings on these effects were reported, and what types of adverse effects were identified in systematic reviews of orthodontic interventions.

Methods Systematic reviews of orthodontic interventions on human patients of any health status, sex, age, and demographics, and socio-economic status, in any type of setting assessing any type of adverse effect scored at any endpoint or timing were eligible. The Cochrane Database of Systematic Reviews and 5 leading orthodontic journals were manually searched for eligible reviews between August 1 2009 and July 31 2021. Study selection and data extraction was conducted by two researchers independently. Prevalence proportions were calculated for four outcomes on seeking and reporting of adverse effects of orthodontic interventions. Univariable logistic regression models were used to determine the association between each one of these outcomes and the journal in which the systematic review was published using the eligible Cochrane reviews as reference.

Results Ninety-eight eligible systematic reviews were identified. 35.7% (35/98) of reviews defined seeking of adverse effects as a research objective, 85.7% (84/98) sought adverse effects, 84.7% (83/98) reported findings related to adverse effects, and 90.8% (89/98) considered or discussed potential adverse effects in the review. Reviews in the journal Orthodontics and Craniofacial Research compared with Cochrane reviews had approximately 7 times the odds (OR 7.20, 95% CI 1.08 to 47.96) to define seeking of adverse effects in the research objectives. Five of the 12 categories of adverse effects accounted for 83.1% (162/195) of all adverse effects sought and reported.

Conclusions Although the majority of included reviews sought and reported adverse effects of orthodontic interventions, end-users of these reviews should beware that these findings do not give the complete spectrum on these effects and that they could be jeopardized by the risk of non-systematically assessing and reporting of adverse effects in these reviews and in the primary studies that feed them. Much research is ahead such as developing core outcome sets on adverse effects of interventions for both primary studies and systematic reviews.

Keywords Orthodontics, Reporting, Systematic review, Interventions, Adverse effect, Adverse event, Harm, Safety, Side effect, Patient important outcomes

Background

To get a balanced perspective of an intervention, systematic reviewers need to report both its beneficial and adverse effects [1]. In this cross-sectional study we assessed whether adverse effects were sought, whether the findings on these effects were reported, and what types of adverse effects were identified in systematic reviews published in the Cochrane Database of Systematic reviews [2] and in 5 leading orthodontic journals.

'Cochrane defines an adverse effect as 'an adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility' [3, 4]. This definition and other key terms used in this manuscript are listed in Table 1 [5, 6]. A wide body of epidemiological studies has shown that adverse effects of interventions in primary research studies are often under-assessed, and/or under-reported, and/or distorted [7–13]. These issues can misinform anyone trying to make valid decisions on a healthcare intervention. An extension of the Consolidated Standards of Reporting Trials (CONSORT) Statement was developed to tackle poor reporting of harms in randomized trials [14]. Since the publication of this statement, the reporting of adverse events in clinical trials has improved, but is still suboptimal [10, 12, 15, 16].

Systematic reviews could provide even more information on adverse effects, because they assess large amounts of data from a wide spectrum of sources (possibly including both published and unpublished data). By assessing the data of multiple single studies, systematic reviewers can make a more balanced assessment of an intervention. This is an important issue, because serious adverse effects may occur rarely and might be missed in single studies. However, epidemiological research showed that the seeking and reporting of adverse effects of interventions and the methods used to identify and synthesize them [17–21] were also poor in systematic reviews. The Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) harms checklist was published in 2016 [22] to improve harms reporting in systematic reviews, but its consequences are still largely unknown.

We performed 2 cross-sectional studies on assessing and reporting of adverse effects in systematic reviews of orthodontic interventions. In this study (part 1), we assessed whether adverse effects were sought and reported and what findings on these adverse effects were reported in systematic reviews of orthodontic interventions published in the Cochrane Database of Systematic reviews [2] and in 5 leading orthodontic journals. In a second study (part 2) we assessed the reporting on adverse effects and the presence of spin on adverse effects in the abstracts of these reviews [23]. Adverse effects of orthodontic interventions refer to for example, pain associated with orthodontic tooth movement, root resorption, decalcifications, periodontal problems, relapse, and undesired health experiences [24]. Recent (November 22 2021) scoping searches confirmed that our research objectives have not been addressed previously.

Table 1 Glossary of terms

Term	Definition
Systematic review	Cochrane [5] defines a systematic review as follows: 'A systematic review attempts to identify, appraise and synthesize all the empirical evidence that meets pre-specified eligibility criteria to answer a specific research question. Researchers conducting systematic reviews use explicit, systematic methods that are selected with a view aimed at minimizing bias, to produce more reliable findings to inform decision making.'
Intervention review	Cochrane [5] defines an intervention review as follows: Intervention reviews assess the benefits and harms of interventions used in healthcare and health policy.
Orthodontic interventions	Steegmans et al. [6] defined orthodontic interventions as follows: 'Orthodontic interventions refer to the use of any type of orthodontic appliance to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodontic treatment, for example retainers.'
Adverse effect	Cochrane [3, 4], defines an adverse effect as an adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility'

Objectives

The objectives of this research study are formulated in the following four research questions:

1) Was seeking of adverse effects of interventions defined as a research objective of the review?

2) Did the review seek any findings related to adverse effects of interventions in the included studies?

3) Did the review report findings related to adverse effects of interventions sought in the included studies?

4) Were potential adverse effects of the intervention considered, discussed (weighed) anywhere in the review?

We also assessed what adverse effects of interventions were defined as research objectives and what adverse effects of interventions were sought and reported in the review.

Methods

The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement [25] and the PRISMA 2020 statement [26, 27] were consulted for reporting this cross-sectional study. The STROBE checklist of items for reporting cross-sectional studies was presented in Additional file 1. The methods for this cross-sectional study were explained in our published protocol [6] and can be consulted through the following link https://systematicreviewsjournal.biome dcentral.com/articles/10.1186/s13643-019-1000-1. We adopted the framework of this protocol to report the methods section of this study and its additional files. Raw data are recorded in Open Science Framework (https://osf.io/ka7mp/). Differences between methods originally planned in the protocol and those implemented in the final research study were given with the rationales for these differences in Additional file 2. No patients were involved in the development of the protocol or in the conduct of this study.

Eligibility criteria

The eligibility criteria have been published previously [6, 28] and are presented again in Table 2 [29].

Table 2 Eligibility criteria

Item	Included	Excluded
Study designs	Systematic reviews of orthodontic interventions. The definition of systematic review, intervention review, and orthodontic interven- tions listed in the Glossary of terms will be used to assess whether a review is eligible (Table 1)	 Non-interventional reviews such as, "Methodology." Diagnostic, "Qualitative", "Prognostic" etc. Rapid and scoping reviews Systematic reviews with Bayesian network meta-analysis Systematic reviews of interventions that did not find any eligible studies (empty reviews)
Participants	Systematic reviews on any type of patients undergoing ortho- dontic interventions, i.e., patients of any health status, sex, age, and demographics, and socio-economic status	 Intervention reviews that focus exclusively on patients with congenital anomalies, for example with cleft lip and palate Systematic reviews of animal or laboratory studies
Interventions	 Systematic reviews that assessed the effects of clinical ortho- dontic interventions. Clinical orthodontic interventions refer to the use of any type of orthodontic appliance that are used to move teeth or change the jaw size or position for orthodontic purposes Systematic reviews of interventions with appliances to maintain or stabilize the outcomes of orthodontic treatment, for example retainers Systematic reviews of orthodontic interventions that compared the effects of orthodontic treatment with or without additional interventions such as pharmacological or small surgical interven- tions, e.g., periodontal or implant surgery No exclusion criteria were applied to the characteristics of the operator who conducted the interventions. 	 Systematic reviews in which patients receive orthodontic treatment, but in which the effects of other interventions, e.g., peri- odontal surgery, were compared and not the effects of orthodontic interventions. Systematic reviews of interventions in which orthodontic appli- ances were specifically used for other purpose, e.g., changing jaw positions to treat respiration or temporomandibular disorders. Systematic reviews of interventions in terventions that included orthognathic surgery. Systematic reviews that focussed exclusively on adverse effects of interventions. Systematic reviews that did not assess a specific orthodontic intervention, but referred to orthodontic treatment as a whole
Outcomes	1) Any adverse effect of orthodontic interventions scored at any endpoint or timing 2) The effects of orthodontic interventions did not refer just to outcomes related to tooth and jaw size and positions, but also to broader outcomes such as periodontal health, esthetic changes, the health of the temporomandibular joint, patient health experi- ences, and economic issues associated with the interventions 3) The reporting of outcomes on adverse effects did not determine eligibility of reviews for this cross-sectional study, i.e., reviews were not excluded because they did not report meas- ured outcome data in a 'usable' way [29]	No exclusion oriteria
Stetting	Any type of setting in which the interventions were conducted, i.e., university or private practice	No exclusion criteria

Information sources and search strategy

The information sources for this study were the Cochrane Database of Systematic Reviews [2] and the websites of 5 leading orthodontic journals. The selection of these 5 orthodontic journals was based on having been published at least 10 years and the highest impact factor [30]. The impact factor in 2018, i.e., the year when the protocol was developed, was used to select these journals. The 5 selected orthodontic journals are: European Journal of Orthodontics [EJO], American Journal of Orthodontics and Dentofacial Orthopedics [AJODO], Angle Orthodontist (AO), The Korean Journal of Orthodontics (KJO), and Orthodontics and Craniofacial Research (O&CR). The impact factors of these journals are listed in Additional file 2. August 1 2009 was chosen as the inception date for searching the information sources, because it coincides with the publication of the PRISMA statement and guidance document on 21 July 2009 [31, 32]. Eligible systematic reviews were manually searched in these information sources from the inception date until July 31 2021.

Study records

Data management

All study selection and data extraction procedures were conducted independently by 2 authors (PS and RMR). Pilot tests were done a priori to train and calibrate these operators [33]. Disagreements between these reviewers during these study selection and data collection were resolved in the following order: Firstly, through discussions; secondly, through rereading the article in question; and thirdly, through contacting of the authors of the pertinent manuscript by email to obtain additional information that could help with decision-making [27]. Persistent disagreements were resolved through discussions with a methodologist (SB). All eligible systematic reviews with their supplementary files were downloaded as PDFs and merged in binder files [34, 35]. Data were collected in an Excel spreadsheet [36].

Study selection and data collection procedures

Titles and abstracts were screened for eligible reviews in the websites of the 5 selected orthodontic journals. Eligible Cochrane reviews were searched in the 'Dentistry and Oral health' section of the Cochrane Database of Systematic Reviews [2]. When Cochrane reviews were updated, we only considered the latest published version. A PRISMA flow diagram was presented to illustrate the selection process of the eligible reviews [26, 27]. All included studies and excluded studies were reported and the rationales for exclusion were given in Additional file 3. Contacting of authors was not necessary to clarify eligibility or data extraction issues. We used our pilot tested data collection forms for the extraction of all pertinent data items. These forms are presented in Additional file 2. The entire eligible review except the abstract and protocol were searched, i.e., the main text, tables, figures, and supplemental files. This strategy was implemented for all eligible reviews. In Cochrane systematic reviews, we also did not search data items in the plain language summary.

Assigning adverse effects of orthodontic interventions

Cochrane defines an adverse effect as 'an adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility' [3, 4]. These events can have a permanent or temporary adverse effect on the health condition of the patient. Root resorption, decalcifications of enamel or caries around orthodontic appliances are well known permanent adverse effects of orthodontic interventions, while pain and discomfort during tooth movement are generally temporary adverse effects. Events associated with orthodontic interventions that could have an adverse effect on the health condition were also labeled as adverse events, e.g., breaking of appliances, failure to complete treatment, and tolerability of orthodontic appliances.

According to our protocol we adopted the framework of known orthodontic adverse effects as reported previously by Preoteasa et al. [24] (Additional file 2) and made some changes in labeling the headings of the various categories of adverse effects (Additional file 2). A total of 12 categories of adverse effects were defined. Additional adverse effects identified during our data extraction procedures were also included in this framework and when ambiguous

the rationale for including these adverse effects was given. The following types of adverse events were not labeled as adverse effects: (1) effects that do not refer to health conditions and could be ambiguous, e.g., costs, duration of treatment, number of appointments etc. (2) effects that refer to pre-existing health problems that can actually improve as a result of the intervention, e.g., respiratory problems as a result of maxillary expansion or self-esteem as a result of the retraction of protruding maxillary incisors.

Power calculation

Epitools epidemiological software was used to calculate the required sample size of eligible systematic reviews of orthodontic interventions [37]. We calculated the required sample size of 73 reviews based on the following input: estimated proportion 0.25, desired precision 0.1, and confidence level 0.95. The estimated proportion was based on the findings in our pilot tests as reported in our protocol [6]. These pilot test showed that findings related to adverse effects were sought in 3 of 12 systematic reviews on orthodontic interventions representing the estimated proportion of 0.25 (3/12).

Outcomes and statistical analyses

We reported the number of retrieved systematic reviews and eligible reviews and calculated the prevalence proportions that addressed our research questions. All outcomes were calculated as originally planned in our published protocol [6]. Prevalence proportions were calculated for: (1) all journals together (2) each journal separately and (3) the group of 5 leading orthodontic journals together and the Cochrane reviews separately. Univariable logistic regression models were built to determine the association between each one of four outcomes and the journal in which the systematic review was published, using the Cochrane Database of Systematic reviews as reference. The strength of association was quantified using odds ratios (OR), and 95% confidence intervals (95% CI). Analyses were performed with the use of commercial software (IBM SPSS 22.0, SPSS Inc, Chicago, IL). A two-sided *P* value of 0.05 was considered to be statistically significant.

Results

Results of the search

Through our searches in the databases of the Cochrane Database of Systematic reviews and the 5 leading orthodontic journals we identified 324 reports. One Cochrane review was excluded, because it was later updated leaving 323 reports for screening. A total of 180 papers was excluded during the title and abstract screening and 45 during full text screening. A total of 98 systematic reviews fulfilled the eligibility criteria of this study. The results of the individual selection steps are presented in a PRISMA flow diagram (Fig. 1) [26, 27]. All included studies are listed in Additional file 3 and excluded studies with the rationale for their exclusion are given in Additional file 4.




Fig. 1 Flow diagram for the selection of systematic reviews of orthodontic interventions

Included studies

Figure 2 presents the number of eligible systematic reviews of orthodontic interventions published during the eligible time span. Table 3 presents the number of eligible reviews for each selected journal and shows that 72.4% (71/98) of the included reviews came from the EJO, AJODO, and AO. Table 3 also gives the types of orthodontic interventions for each of these journals, which are divided in three categories. Category 1 refers to orthodontic interventions to move teeth modify jaws such as fixed orthodontic appliances or palatal expansion appliances. Category 2 refers to orthodontic interventions that also include additional surgical, pharmacological or vibrational interventions such as mini-implants, prostaglandins, piezo surgery, or vibratory stimulation. Category 3 refers to orthodontic interventions with appliances to maintain or stabilize orthodontic treatment results such as retainers. The majority of included reviews, 70.4% (69/98), assessed orthodontic interventions with additional surgical, pharmacological or vibratory simulation.




Fig. 2 Line chart of the number of eligible systematic reviews published between August 1 2009 and July 31 2021

Table 3 Characteristics of included reviews

Type of orthodontic intervention	Cochrane	EJO	AJODO	AO	KJO	O&CR	Total
Category 1. Orthodontic interventions to move teeth or modify jaws	7	21	14	15	2	10	69
Category 2. Orthodontic interventions with additional surgical, pharmacological or vibratory interventions	3	6	6	8	1	4	28
Category 3. Orthodontic interventions to maintain or stabilize orthodontic results	0	1	0	0		0	1
Total	10	28	20	23	3	14	98

Outcomes to the research questions

Figure 3 presents the answers to each individual research question and Table 4 gives the proportions. We reported the proportions in answering the four research questions over time in Table 5. The prevalence of reviews that defined seeking of adverse effects of interventions as a research objective was low, i.e., 35.7% in the 98 eligible reviews. Instead, the proportions that addressed the other 3 research question were 85% and higher indicating that seeking and reporting of findings related to adverse effects of interventions in the included studies and considering or discussing potential adverse effects anywhere in the review were implemented in most of the eligible reviews. As compared to the Cochrane Database of Systematic Reviews, the journal of Orthodontics and Craniofacial research had approximately 7 times the odds (OR 7.20, 95%CI 1.08 to 47.96) to report that adverse effects were sought in the research objectives (Table 6). For the other 3 outcomes, no statistical analysis was performed considering the low variability in the response scored (prevalence of 'no' ranging from 9.2 to 15.3%) and the overall small sample sizes (Table 4).

Labeling adverse effects of orthodontic interventions

The type of adverse effects most frequently defined as research objectives were adverse effects related to (1) tooth structures, (2) periodontal tissues, (3) undesired treatment results, (4) relapse and stability, and (5) negative qualitative experiences by the patient or carer(s) (Table 7). These were also the most prevalent types of adverse effects sought in the included studies and reported in the review and accounted for 83.1% (162/195) of all adverse effects sought and reported (Table 8). We were able to categorize all 195 adverse effects except one and labeled it 'Additional adverse effects' (Table 8).




Fig. 3 Seeking adverse effects in systematic reviews of orthodontic interventions

Table 4 Outcomes on seeking adverse el	cts in systematic reviews of orthodontic interventions
--	--

Description of outcomes	Cochrane	EDO	AJODO	AO	KIO	O&CR	All orthodontic journals	All journals
The number of identified systematic reviews	29	100	68	61	11	53	293	322
The number of eligible systematic reviews of orthodontic interven- tions	10	28	20	23	3	14	88	98
The prevalence of eligible systematic reviews of orthodontic interventions that defined seeking of adverse effects of interventions as a research objective of the review [®]	20.0% (2/10)	32.1% (9/28)	45% (\$/20)	21.7 (5/23)	33.3% (1/3)	64.3% (9/14)	47.7% (33/88)	35.7% (35/98
The prevalence of eligible systematic reviews of orthodontic interventions that sought any findings related to adverse effects of interven- tions in the included studies ⁶	100.0% (10/10)	92.9% (26/28)	75.0% (15/20)	78.3% (18/23)	100.0% (3/3)	85.7% (12/14)	84.1% (74/88)	85.7% (84/98
The prevalence of eligible systematic reviews of orthodontic interventions that reported findings related to the adverse effects of interventions sought in the included studies"	100.0% (10/10)	92.9% (26/28)	70.0% (14/20)	78.3% (18/23)	100.0% (3/3)	85.7% (12/14)	83.0% (73/88)	84.7% (83/98
The prevalence of eligible systematic reviews of orthodontic interventions that considered, discussed (weighed) potential adverte effects of the intervention anywhere in the review *	100.0% (10/10)	92.9% (26/28)	85.0% (17/20)	91.3% (21/23)	100.0% (3/3)	85.7% (12/14)	89.8% (79/88)	90.8% (89/98

* The denominator for calculating the proportions for each journal is the number of included reviews for that journal. The denominator for calculating the proportions for all journals together is the total of included reviews, i.e., 98

Table 5 Outcomes on seeking adverse effects in systematic reviews of orthodontic intervention	ns by year of publication
---	---------------------------

		effects of defined a	ing of adverse interventions s a research of the review	Did the review seek any findings related to adverse effects of interventions in the included studies?		Did the review report findings related to adverse effects of interventions sought in the included studies?		Were potential adverse effects of the intervention considered, discussed (weighed) anywhere in the review?		
Year		Yes	No	Yes	No	Yes	No	Yes	No	Total
2009	Count	0	1	1	0	1	0	1	0	1
	% within Year	0.0%	100.0%	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%
2010	Count	1	2	2	1	2	1	3	0	3
	% within Year	33.3%	66.7%	66.7%	33.3%	66.7%	33.3%	100.0%	0.0%	100.0%
2011	Count	3	4	6	1	6	1	6	1	7
	% within Year	42.9%	57.1%	85.7%	14.3%	85.7%	14.3%	85.7%	14.3%	100.0%
2012	Count	2	0	2	0	2	0	2	0	2
	% within Year	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%
2013	Count	3	6	9	0	9	0	9	0	9
	% within Year	33.3%	66.7%	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%
2014	Count	3	6	7	2	7	2	7	2	9
	% within Year	33.3%	66.7%	77.8%	22.2%	77.8%	22.2%	77.8%	22.2%	100.0%
2015	Count	3	9	9	3	9	3	12	0	12
	% within Year	25.0%	75.0%	75.0%	25.0%	75.0%	25.0%	100.0%	0.0%	100.0%
2016	Count	1	8	6	3	6	3	7	2	9
	% within Year	11,1%	88.9%	66.7%	33.3%	66.7%	33.3%	77.8%	22.2%	100.0%
2017	Count	2	7	8	1	7	2	8	1	9
	% within Year	22.2%	77.8%	88.9%	11,1%	77.8%	22.2%	88.9%	11.1%	100.0%
2018	Count	4	9	10	3	10	3	10	3	13
	% within Year	30.8%	69.2%	76.9%	23.1%	76.9%	23.1%	76.9%	23.1%	100.0%
2019	Count	3	1	4	0	4	0	4	0	4
	% within Year	75.0%	25.0%	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%
2020	Count	7	5	12	0	12	0	12	0	12
	% within Year	58.3%	41,796	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%
2021	Count	3	5	8	0	8	0	8	0	8
	% within Year	37.5%	62.5%	100.0%	0.0%	100.0%	0.0%	100.0%	0.0%	100.0%
Total	Count	35	63	84	14	83	15	89	9	98
	% within Year	35.7%	64.3%	85.7%	14.3%	84.7%	15.3%	90.8%	9,2%	100.0%

 Table 6
 Results
 of
 univariable
 logistic
 regression
 including
 journal as a predictor variable of seeking adverse effects as an objective of the systematic review
 including
 <t

Journal	Odds ratio	Lower 95% CI	Upper 95% CI	Pvalue
Cochrane	1	(4)	-	-
AJODO	3.27	0.55	19.45	0.19
AO	1.11	0.18	6.99	0.91
EJO	1.90	0.33	10.80	0.47
KJO	2.00	0.12	34.82	0.63
OSC	7.20	1.08	47.96	0.04

Table 7 Type of adverse effects defined as research objectives of the review

Adverse effects related to	Prevalence
Tooth structures	16.7% (8/48)
Periodontal tissues	12.5% (6/48)
Intraoral (non-tooth or periodontal) tissues	0.0% (0/48)
Extraoral tissues (non-temporomandibular tissues)	0.0% (0/48)
Temporomandibular tissues and disorders	2.1% (1/48)
Undesired treatment results	18.8% (9/48)
Relapse and stability	20.8% (10/48)
Negative qualitative experiences by the patient or carer(s)	18.8% (9/48)
Appliance failure	0.0% (0/48)
Gastro-intestinal	0.0% (0/48)
Non-defined	10.4% (5/48)
Additional adverse effects	0.0% (0/48)

Table 8 Type of adverse effects sought and reported in the review

Adverse effects related to	Prevalence
Tooth structures	12.8% (25/195
Periodontal tissues	13.8% (27/195
Intraoral (non-tooth or periodontal) tissues	2,6% (5/195)
Extraoral tissues (non-temporomandibular tissues)	0.5%(1/195)
Temporomandibular tissues and disorders	4.1% (8/195)
Undesired treatment results	22.1% (43/195
Relapse and stability	18.5% (36/195)
Negative qualitative experiences by the patient or carer(s)	15.9% (31/195
Appliance failure	7,7% (15/195)
Gastro-intestinal	0.5%(1/195)
Non-defined	1.0% (2/195)
Additional adverse effects*	0.5%(1/195)

Discussion

Principal findings of the study

This cross-sectional study showed that in 35.7% (35/98) of reviews of orthodontic interventions seeking of adverse effects was defined as an objective. In 85.7% (84/98) of these reviews, findings related to adverse effects of interventions were sought and in 84.7% (83/98) the reviewers reported on these findings. In more than 90% (89/98) of included systematic reviews, the reviewers discussed (weighed) potential adverse effects of interventions somewhere in the review. Five types of adverse effects accounted for 83.1% (162/195) of adverse effects that were sought and reported in the eligible reviews.

Comparisons with other studies

The proportion of included reviews that defined seeking of adverse effects as a research objective was low, i.e., 35.7% (35/98) in both Cochrane and non-Cochrane systematic reviews (Table 4). Assessing potential adverse effects of interventions is considered a mandatory item when setting the research question for Cochrane intervention reviews [1]. Not defining seeking of adverse effects as a research objective can mislead end-users of systematic reviews. Authors therefore need to include this item in their research objectives and editors and peer reviewers should verify its implementation.

The proportions of reviews that reported findings related to adverse effects of interventions were higher in this sample of orthodontic reviews (84.7%(83/98) compared with gastroenterology reviews (66.7% (52/78) [18], Cochrane reviews of interventions (75.6% (59/78), and Database of Abstracts of Reviews of Effects (DAREs) reviews (48.1% (38/79) [38]. Explanations for these higher proportions could be: (1) the time period of inclusion of reviews (2) the research design and type of interventions of the studies included in the reviews (3) the field of research. Orthodontic research could be more focused on assessing adverse effects of interventions than other fields, because this assessment is an integral part of routine clinical practice. For example, assessing adverse effects such as undesired

treatment results and relapse and stability are part of everyday problems in orthodontic practice and accounted for 40.5% (79/195) of adverse effects sought and reported in this sample of systematic reviews of orthodontic interventions (Table 8).

Strengths and limitations

This cross-sectional study has the following strengths: (1) scoping searches were conducted to identify knowledge gaps, (2) pilot studies were conducted to calibrate researchers and fine-tune research questions and methodology, (3) a protocol was developed and published a priori [6], and (4) all raw data were included with this manuscript or recorded in Open Science Framework (https://osf.io/ka7mp/). This study also has limitations. First, the findings of this cross-sectional study are expected to be better than those reported in the entire body of orthodontic literature, because we assessed reviews published in the five leading orthodontic journals and those listed in the Cochrane Database of Systematic Reviews. Second, the risk of selective (non) reporting bias regarding adverse effects in the eligible reviews. Third, only reviews published in a pre-established period (August 1 2009 until July 31 2021) were eligible, instead of having considered a larger sample, e.g., by having included reviews prior to the inception date. However, we chose this inception date, because it coincides with the launch of the PRISMA statement [31, 32], which provides reviewers better guidance on reporting.

Implications and future research

Several of our findings seem promising at a first glance. For example, the proportion of reviews that sought and reported adverse effects was relatively high, i.e., (84.7% (83/98), but a variety of issues has to be considered when interpreting this finding. First, this proportion only refers to whether or not reviewers implemented this item, but not how. For example, the reviewers could have reported on just one or a selection of all adverse effects assessed and reported in the eligible studies for their reviews. Second, this proportion also does not give any information on the magnitude, and duration of adverse effects nor on the time points for assessing them. Third, we do not know whether all adverse effects were indeed sought and reported as originally planned in the registered protocols of the included reviews. For example, Parsons et al. [39] showed that this was not the case in their sample of systematic reviews of health care interventions. In 35% (51/146) of these reviews they found discrepancies between what was planned in the protocol as registered in PROSPERO and what was reported on adverse effects in the final published reviews. Fourth, a wide body of evidence has shown that adverse events were often assessed inconsistently and reported inadequately in clinical trials and that most results on these events were not available in public sources [8, 40–42]. If these limitations also apply to the clinical trials that fed the reviews of this study one should further question the validity of the findings on adverse effect of systematic reviews of orthodontic interventions.

Strategies to improve the validity of what is reported on adverse effects of orthodontic interventions in systematic reviews include developing tailored core outcome sets on these effects [43] as well as guidelines for assessing and reporting them in both primary research and systematic reviews. Additional strategies on synthesizing adverse effects in systematic reviews at multiple levels were published in a recent paper by Qureshi et al. [19]. By

implementing such strategies progress on the assessing and reporting of adverse effects of orthodontic interventions in both primary studies and systematic reviews can be made.

In conclusion the promising findings of this study should be interpreted with caution by its end users, because they could be jeopardized by numerous uncertainties. Much research is ahead to create valid and usable knowledge on adverse effects of orthodontic interventions involving a wide body of stakeholders.

Abbreviations

AJODO American Journal of Orthodontics and Dentofacial Orthopedics AO Angle Orthodontist CONSORT Consolidated Standards of Reporting Trials EJO European Journal of Orthodontics KJO Korean Journal of Orthodontics O&CR Orthodontics and Craniofacial Research PRISMA Preferred reporting items for systematic reviews and meta-analyses PROSPERO Prospective Register of Systematic Reviews STROBE Strengthening the Reporting of Observational Studies in Epidemiology

Supplementary Information

The online version contains supplementary material available at https://doi. org/10.1186/s13643-023-02273-7.

Additional file 1. STROBE Statement: Checklist of items that should be included in reports of cross-sectional studies [25].

Additional file 2. A. Differences between the protocol and the completed cross-sectional study. B. Selected journals and their 2018 impact factor (Clarivate Analytics 2021). C. Data collection forms*. D. Adverse effects hypothetically linked to orthodontic interventions according to Preoteasa et al. [24]*. E. Adverse effects hypothetically linked to orthodontic interventions*.

Additional file 3. Included reviews.

Additional file 4. Excluded studies.

Acknowledgements

Not applicable.

Authors' contributions

PS and RMR conceived and designed this cross-sectional study and RMR is the guarantor. PS and RMR conducted the study selection procedures and data extraction. PS, RMR, and NDG conducted the data analyses and statistical analyses. SB provided support on methodological

issues and assisted in the overall fine-tuning of this cross-sectional study. All four authors revised the various draft versions of this paper and read and approved the final submitted manuscript.

Funding

All expenses for preparing and conducting of the submitted research study were paid evenly by each author.

Availability of data and materials

Not applicable.

Declarations Ethics approval and consent to participate Not applicable.

Consent for publication Not applicable.

Competing interests

Reint Meursinge Reynders and Shandra Bipat are both Associate Editors for Systematic Reviews. All four authors declare that they have no further competing interests.

Author details

¹Department of Orthodontics, Academisch Centrum Tandheelkunde Amsterdam (ACTA), University of Amsterdam, Gustav Mahlerlaan 3004, Amsterdam 1081 LA, The Netherlands. ²Department of Clinical Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY, USA. ³EBMVet, Via Sigismondo Trecchi 20, Cremona, CR 26100, Italy. ⁴Department of Radiology and Nuclear Medicine, Amsterdam University Medical Center (Amsterdam UMC), Location AMC, Meibergdreef 9, Amsterdam 1105 AZ, The Netherlands. ⁵Department of Oral and Maxillofacial Surgery, Amsterdam University Medical Center (Amsterdam UMC), Location AMC, Meibergdreef 9, Amsterdam 1105 AZ, The Netherlands. ⁶Studio Di Ortodonzia, Via Matteo Bandello 15, Milan 20123, Italy.

Received: 11 February 2022 Accepted: 15 June 2023

Published online: 03 July 2023

References

- Higgins JPT, Lasserson T, Chandler J, Tovey D, Thomas, J, Flemyng E, et al. Methodological expectations of cochrane intervention reviews. Cochrane: London. Version February 2022. Available from: https://commu nity.cochrane.org/mecir-manual. Accessed 27 June 2023.
- The Cochrane Database of Systematic Reviews (CDSR). Available from: https://www.cochranelibrary.com/cdsr/about-cdsr. Accessed 4 Dec 2021.
- 3. Glossary of terms in the Cochrane Collaboration. Version 4.2.5. Updated May 2005. Available from: http://aaz.hr/resources/pages/57/7.%20Coc hrane%20glossary.pdf. Accessed 4 Dec 2021.
- Peryer G, Golder S, Junqueira D, Vohra S, Loke YK. Chapter 19: Adverse effects. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. Cochrane handbook for systematic reviews of Interventions version 6.2 (updated February 2021). Cochrane; 2021. Available from www.training.cochrane.org/handbook.
- Cochrane library. Available from: https://www.cochranelibrary.com/ about/about-cochrane-reviews. Accessed 27 June 2023.
- Steegmans PAJ, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross- sectional study. Syst Rev. 2019;8(1):89. https://doi.org/10.1186/s13643- 019-1000-1. PMID: 30953538; PMCID: PMC6449933.
- Bagul NB, Kirkham JJ. The reporting of harms in randomized controlled trials of hypertension using the CONSORT criteria for harm reporting. Clin Exp Hypertens. 2012;34(8):548–54. https://doi.org/10.3109/10641963. 2012.681724. Epub 2012 May 9.
- Golder S, Loke YK, Wright K, Norman G. Reporting of adverse events in published and unpublished studies of health care interventions: a systematic review. PLoS Med. 2016;13(9):e1002127. https://doi.org/10.1371/ journal.pmed.1002127. eCollection 2016 Sep.
- Hodkinson A, Kirkham JJ, Tudur-Smith C, Gamble C. Reporting of harms data in RCTs: a systematic review of empirical assessments against the CONSORT harms extension. BMJ Open. 2013;3(9):e003436. https://doi. org/10.1136/bmjopen-2013-003436.
- Péron J, Maillet D, Gan HK, Chen EX, You B. Adherence to CONSORT adverse event reporting guidelines in randomized clinical trials evaluating systemic cancer therapy: a systematic review. J Clin Oncol. 2013;31(31):3957–63. https://doi.org/10.1200/JCO.2013.49.3981. Epub 2013 Sep 23.
- Pradhan R, Singh S. Comparison of data on serious adverse events and mortality in ClinicalTrials.gov, corresponding journal articles, and FDA medical reviews: cross-sectional analysis. Drug Saf. 2018;41(9):849–57. https://doi.org/10.1007/s40264-018-0666-y. PMID: 29644579.
- Smith SM, Chang RD, Pereira A, Shah N, Gilron I, Katz NP, Lin AH, McDermott MP, Rappaport BA, Rowbotham MC, Sampaio C, Turk DC, Dworkin RH. Adherence to CONSORT harms-reporting recommendations in publications of recent analgesic clinical trials: an ACTTION systematic review. Pain. 2012;153(12):2415–21. https://doi.org/10.1016/j.pain.2012.08.009. Epub 2012 Sep 15.
- 13. Tang E, Ravaud P, Riveros C, Perrodeau E, Dechartres A. Comparison of serious adverse events posted at ClinicalTrials.gov and published in cor- responding journal articles. BMC Med. 2015;14(13):189. https://doi.org/10. 1186/s12916-015-0430-4. PMID: 26269118; PMCID: PMC4535304.
- Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, Schulz K, Moher D, CONSORT Group. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med. 2004;141(10):781–8.
- Haidich AB, Birtsou C, Dardavessis T, Tirodimos I, Arvanitidou M. The quality of safety reporting in trials is still suboptimal: survey of major general medical journals. J Clin Epidemiol. 2011;64(2):124–35. https://doi.org/10.1016/j.jclinepi.2010.03.005. Epub 2010 Jun 17.
- Moskowitz A, Andersen LW, Holmberg MJ, Grossestreuer AV, Berg KM, Granfeldt A. Identification, collection, and reporting of harms among non-industry-sponsored randomized clinical trials of pharmacologic interventions in the critically ill population: a systematic review. Crit Care. 2020;24(1):398. https://doi.org/10.1186/s13054-020-03113-z. PMID:32641 148;PMCID:PMC7346341.
- Li L, Xu C, Deng K, Zhou X, Liu Z, Busse JW, Ren Y, Zou K, Sun X. The reporting of safety among drug systematic reviews was poor before the implementation of the PRISMA harms checklist. J Clin Epidemiol. 2019;105:125–35. https://doi.org/10.1016/j.jclinepi.2018.09.014. Epub 2018 Sep 29 PMID: 30278212.

- Mahady SE, Schlub T, Bero L, Moher D, Tovey D, George J, Craig JC. Side effects are incompletely reported among systematic reviews in gastroenterology. J Clin Epidemiol. 2015;68(2):144–53.
- 19. Qureshi R, Mayo-Wilson E, Rittiphairoj T, McAdams-DeMarco M, Guallar E, Li T. Summaries of harms in systematic reviews are unreliable Paper 2: Methods used to assess harms are neglected in systematic reviews of gabapentin. J Clin Epidemiol. 2021. https://doi.org/10.1016/j.jclinepi. 2021.10.024.
- Qureshi R, Mayo-Wilson E, Rittiphairoj T, McAdams-DeMarco M, Guallar E, Li T. Summaries of harms in systematic reviews are unreliable Paper 3: Given the same data sources, systematic reviews of gabapentin have different results for harms. J Clin Epidemiol. 2021. https://doi.org/10.1016/j. jclinepi.2021.10.025.
- Zorzela L, Golder S, Liu Y, Pilkington K, Hartling L, Joffe A, Loke Y, Vohra S. Quality of reporting in systematic reviews of adverse events: systematic review. BMJ. 2014;8(348):f7668. https://doi.org/10.1136/bmj.f7668. PMID: 24401468; PMCID: PMC3898583.
- Zorzela L, Loke YK, Ioannidis JP, Golder S, Santaguida P, Altman DG, Moher D, Vohra S, PRISMA Harms Group. PRISMA harms checklist: improving harms reporting in systematic reviews. BMJ. 2016;1(352):i157. https://doi.org/10.1136/bmj.i157.
- Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin on adverse effects in abstracts of systematic reviews of orthodontic interventions: a cross-sectional study (part 2). Syst Rev. 2023;12(1):99. https://doi.org/10. 1186/s13643-023-02269-3.
- Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F, editor. Orthodontics-basic aspects and clinical considerations. March 9, 2012 under CC BY 3.0 license. Available from: www.intechopen.com. https://cdn.intec hopen.com/pdfs/31388/InTech-Risks_and_complicationsassociated_ with_orthodontic_treatment.pdf. Accessed 4 Dec 2021.
- 25. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Ann Intern Med. 2007;147(8):573–7.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald

S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021;29(372):n71. https://doi.org/10.1136/ bmj.n71. PMID: 33782057; PMCID: PMC8005924.

 Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch

VA, Whiting P, McKenzie JE. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. BMJ. 2021;29(372):n160. https://doi.org/10.1136/bmj.n160. PMID: 33781993.

- Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews. Res Integr Peer Rev. 2019;4:27. https://doi.org/10.1186/ s41073-019-0084-4.
- 29. Lefebvre C, Glanville J, Briscoe S, Littlewood A, Marshall C, Metzendorf M-I, Noel-Storr A, Rader T, Shokraneh F, Thomas J, Wieland LS. Chapter 4: searching for and selecting studies. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. Cochrane handbook for systematic reviews of Interventions version 6.2 (updated February 2021). Cochrane; 2021. Available from www.training.cochrane.org/handbook.
- 30. Clarivate Analytics. Available from: https://clarivate.com/. Accessed 4 Dec 2021.
- 31. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. PLoS Med. 2009;6:e1000100 Epub 2009 Jul 21.
- 32. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med. 2009;6:e1000097.
- Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. the PRISMA-P Group: Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;349:g7647.

- Adobe Merge PDFs, combine files into one PDF. Available from: http:// www.wikihow.com/Merge-PDF-Files. Accessed 4 Dec 2021.
- 35. Acrobat X action: find & highlight words & phrases. Available from: https://acrobatusers.com/assets/uploads/actions/Find_and_Highlight_Words_and_Phrases.pdf . Accessed 4 Dec 2021.
- 36. Microsoft Excel. Available from: https://products.office.com/it-IT/excel. Accessed 4 Dec 2021.
- Sergeant ESG. Sergeant, ESG, 2018. Epitools epidemiological calculators. Ausvet. Available from: http://epitools.ausvet.com.au. Accessed 4 Dec 2021.
- Hopewell S, Wolfenden L, Clarke M. Reporting of adverse events in systematic reviews can be improved: survey results. J Clin Epidemiol. 2008;61(6):597–602. https://doi.org/10.1016/j.jclinepi.2007.10.005. Epub 2008 Apr 14 PMID: 18411039.
- Parsons R, Golder S, Watt I. More than one-third of systematic reviews did not fully report the adverse events outcome. J Clin Epidemiol. 2019;108:95–101. https://doi.org/10.1016/j.jclinepi.2018.12.007. Epub 2018 Dec 13 PMID: 30553831.
- Golder S, Loke YK, Wright K, Sterrantino C. Most systematic reviews of adverse effects did not include unpublished data. J Clin Epidemiol. 2016;77:125–33. https://doi.org/10.1016/j.jclinepi.2016.05.003. Epub 2016 Jun 1 PMID: 27259470.
- Mayo-Wilson E, Fusco N, Li T, Hong H, Canner JK, Dickersin K, MUDS investigators. Harms are assessed inconsistently and reported inadequately part 1: systematic adverse events. J Clin Epidemiol. 2019;113:20–7. https://doi.org/10.1016/j.jclinepi.2019.04.022. Epub 2019 May 2 PMID: 31055175.
- Mayo-Wilson E, Fusco N, Li T, Hong H, Canner JK, Dickersin K, MUDS investigators. Harms are assessed inconsistently and reported inadequately Part 2: nonsystematic adverse events. J Clin Epidemiol. 2019;113:11–9. https://doi.org/10.1016/j.jclinepi.2019.04.020. Epub 2019 May 2 PMID: 31055176.
- Tsichlaki A, O'Brien K, Benson PE, Marshman Z, Johal A, Colonio-Salazar FB, Harman NL, Fleming PS. Development of a core outcome set for use in routine orthodontic clinical trials. Am J Orthod Dentofacial Orthop. 2020;158(5):650–60. https://doi.org/10.1016/j.ajodo.2020.05.010. Epub 2020 Sep 16 PMID: 32950336.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



CHAPTER 4

Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews

This is a non-edited version of the manuscript:

Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross- sectional study of systematic reviews. Res Integr Peer Rev. 2019 Dec 19;4:27.

Abstract

Background: Titles and abstracts are the most read sections of biomedical papers. It is therefore important that abstracts transparently report both the beneficial and adverse effects of health care interventions and do not mislead the reader. Misleading reporting, interpretation, or extrapolation of study results is called "spin". In this study, we will assess whether adverse effects of orthodontic interventions were reported or considered in the abstracts of both Cochrane and non-Cochrane reviews and whether spin was identified and what type of spin.

Methods: Eligibility criteria were defined for the type of study designs, participants, interventions, outcomes, and settings. We will include systematic reviews of clinical orthodontic interventions published in the five leading orthodontic journals and in the Cochrane Database. Empty reviews will be excluded. We will manually search eligible reviews published between 1 August 2009 and 31 July 2019. Data collection forms were developed a priori. All study selection and data extraction procedures will be conducted by two reviewers independently. Our main outcomes will be the prevalence of reported or considered adverse effects of orthodontic interventions in the abstract of systematic reviews and the prevalence of "spin" related to these adverse effects. We will also record the prevalence of three subtypes of spin, i.e., misleading reporting, misleading interpretation, and misleading extrapolation- related spin. All statistics will be calculated for the following groups: (1) all journals individually, (2) all journals together, and (3) the five leading orthodontic journals and the Cochrane Database of Systematic Reviews separately. Generalized linear models will be developed to compare the various groups.

Discussion: We expect that our results will raise the awareness of the importance of reporting and considering of adverse effects and the presence of the phenomenon of spin related to these effects in abstracts of systematic reviews of orthodontic interventions. This is important, because an incomplete and inadequate reporting, interpretation, or extrapolation of findings on adverse effects in abstracts of systematic reviews can mislead readers and could lead to inadequate clinical practice. Our findings could result in policy implications for making judgments about the acceptance for publication of systematic reviews of orthodontic interventions.

Keywords: Orthodontics, Reporting, Systematic review, Intervention, Spin, Misleading reporting, Misleading interpretation, Misleading extrapolation, Adverse effect, Adverse event, Harm, Safety

Background

Readers of the biomedical literature mostly just screen the title and the abstract of an article without assessing the full publication [1]. The beneficial and adverse effects of interventions should therefore be transparently reported in these summaries and should not mislead its readers. Misleading reporting, interpretation, or extrapolation of study results is called "spin" [2–4]. We will assess in abstracts of both Cochrane and non-Cochrane reviews whether adverse effects of orthodontic interventions were reported or considered and whether spin was identified and what type of spin.

Titles and abstracts are the most read sections of biomedical papers [1], because assessing the full research article is often conditioned by paywalls or because of a lack of time or language issues of the readers [1]. Abstracts should therefore clearly and truthfully reflect the objectives, methods, results, and the interpretation of research findings. The standard for Methodological Expectations of Cochrane Intervention Reviews (MECIR) [5] has listed a series of highly desirable and mandatory items that should be consulted by reviewers when preparing the abstract of their reviews. Item R13 of the MECIR standard states that: "The Abstract of the review should aim to reflect a balanced summary of the benefits and harms of the intervention." This mandatory item is particularly crucial for presenting adverse effects of health care interventions, because these effects are often poorly reported in systematic reviews [6]. Numerous epidemiological studies have also shown that the assessment and reporting of adverse effects of interventions in primary research studies is often suboptimal [7–11]. We adopted Cochrane's definition of adverse effects: "An adverse event for which the causal relation between the intervention and the event is at least a reasonable possibility" [12, 13]. This definition and other key terminology in this manuscript are summarized in Table 1 [2-4, 12-15].

When presenting information on adverse effects in the abstract, it is also crucial that it does not mislead the reader. A distorted presentation of study results has been defined as "spin" [3], but more elaborate definitions are also used (Table 1). The term spin was first used in 1995 in the medical literature by Horton [16] and has been further subdivided into three categories [4]: misleading reporting-related spin, misleading interpretation-related spin, and misleading extrapolation-related spin (Table 1). Yavchitz et al. [17] have ranked the various types of spin according to their severity. The severest form of spin in abstracts of systematic reviews and meta-analyses was scored for "conclusions that contain recommendations for clinical practices that were not supported by findings" [17]. A high prevalence of the various types of spin has been identified in multiple epidemiological studies [4, 18-22]. Boutron et al. [18] found spin in 50% (36/72) of the conclusions sections of the main text of parallelgroup RCTs and in 58.3% (42/72) of the conclusions sections of the abstracts. Spin was also common in diagnostic accuracy studies published in journals with high impact factors [22]. Lockyer et al. [21] showed that spin is a frequent phenomenon in abstracts of RCTs of wound treatments, and Lazarus et al. [4] identified at least one example of spin in 84% (107/128) of the abstracts of non-randomized intervention studies. Spin is in strong conflict with the Declaration of Helsinki [23] that states that: "Authors have a duty to make publicly available the results of their research on human subjects and are accountable for the completeness and accuracy of their reports."

In this study, we will assess whether potential adverse effects of orthodontic interventions were reported or considered (i.e., discussed, weighed, etc.) in the abstract of systematic reviews. We will further assess whether spin was introduced regarding information on these adverse effects in the abstract, and we will categorize the types of spin (Table 1). We will assess these issues in the five leading orthodontic journals and those included in the Cochrane Database of Systematic Reviews. In these reviews, we will assess adverse effects such as pain as a result of tooth movement and the adverse effects defined by Preoteasa et al. (Table 2) [24]. Scoping searches in the orthodontic literature confirmed the knowledge gaps on our research questions. Our pilot studies quantified these gaps and confirmed the need to address these questions. We will assess these issues in systematic reviews, because they are increasingly consulted by patients [25] and when well-conducted systematic reviews are considered among the information sources with the highest level of evidence [26]. Our research questions are important, because incomplete or misleading information on adverse effects of interventions may have detrimental effects on the treatment of orthodontic patients.

Table 1 Glossary of terms

Term	Definition			
Systematic review	The Cochrane glossary [12] defines a systematic review as "A review of a clearly formulated question that uses systematic and explicit methods to identify, select, and critically appraise relevant research, and to collect and analyse data from the studies that are included in the review. Statistical methods (meta-analysis) may or may not be used to analyse and summarise the results of the included studies."			
Intervention review	Cochrane [14] defines an intervention review as follows: "Intervention reviews assess the benefits and harms of interventions used in healthcare and health policy."			
Orthodontic interventions	Steegmans et al. [15] define orthodontic interventions as follows: "Orthodontic interventions refer to the use of any type of orthodontic appliance that are used to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodonti treatment, for example retainers."			
Adverse effect	Cochrane [12, 13] defines an adverse effect as 'an adverse event for which the caus relation between the intervention and the event is at least a reasonable possibility.			
Spin [3]	"Distorted presentation of study results."			
Spin [3]	"A misrepresentation of study results, regardless of motive (intentionally or unintentionally) that overemphasizes the beneficial effects of the intervention and overstates safety compared with that shown by the results."			
Spin [2]	"A specific intentional or unintentional reporting that fails to faithfully reflect the nature and range of findings and that could affect the impression the results produce in readers."			
Misleading reporting related-spin [4]	"Incomplete reporting of the study results that could be misleading for the reader."			
Misleading interpretation related-spin (4)	Inadequate interpretation of the study results overestimating the beneficial effect of the intervention.			
Misleading extrapolation related-spin [4]	Inappropriate generalization of the study results by inadequate (1) extrapolation from the population, interventions, or ourcome actually assessed in the study to a larger population, different interventions, or ourcomes, or (2) inadequate implications for clinical practice.			
Spin (in the abstract) on adverse effects of interventions	Incomplete or inadequate reporting, interpretation, or extrapolation (or a combination of these variables) of findings on adverse effects of interventions in the abstract that could be misleading for the reader.			

Subgroup	Description
Local adverse effects	
Dental	 Crown: decalcifications, decays, tooth wear, enamel cracks and fractures; discolorations, deterioration of prosthetic crown (as fracturing a ceramic one during debonding);
	 Root: root resorption, early closure of root apex, ankylosis;
	Pulp: ischemia, pulpitis, necrosis;
Periodontal	 Gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscences, fenestrations, interdental fold, dark triangles;
Temporomandibular joint	 Condylar resorption, temporomandibular dysfunction;
Soft tissues of the oral and maxillofacial region	 Trauma (e.g., long archwires, headgear related), mucosal ulcerations or hyperplasia, chemical burns (e.g., etching related), thermal injuries (e.g., overheated burs), stomatiti- clumsy handling of dental instruments;
Unsatisfactory treatment outcome	 Inadequate morpho-functional, aesthetic or functional final result, relapse, failure to complete treatment due to treatment dropout.
Systemic adverse effects	
Psychological	 Teasing, behavioral changes of patients and parents; discomfort associated with pain presence and aesthetic look discontents during orthodontic appliance usage;
Gastro-intestinal	Accidental swallowing of small parts of the orthodontic device (tubes, brackets);
Allergies	To nickel or latex;
Cardiac	 Infective endocarditis;
Chronic fatigue syndrome	
Cross infections	- From doctor to patient, patient to doctor, patient to patient.

Table 2 Adverse effects hypothetically linked to orthodontic interventions [24]

Objectives


The objectives of this research study are summarized in the following research questions:

Research questions

- In abstracts of systematic reviews of orthodontic interventions, were potential adverse effects of these interventions reported or considered (i.e., discussed, weighed, etc.)?
- In abstracts of systematic reviews of orthodontic interventions, was spin identified in the reporting, interpretation, and extrapolation of adverse effects?
- What was the prevalence of each type of spin?

Methods

This protocol is reported according to the guidance of the Preferred Reporting Items for Systematic Review and Meta- Analysis Protocols (PRISMA-P) 2015 statement, and the PRISMA-P checklist is included as Additional file 1 [27, 28]. We adopted the same flow of research methods as reported in our published protocol on seeking adverse effects in systematic reviews of orthodontic interventions (Fig. 1) and con-ducted our pilot tests on the same sample of systematic reviews as was described in our previous protocol [15]. Our sample size of 14 reviews for the pilot test was calculated with the formula reported by Viechtbauer et al. [29]. Further details on the methods of our pilot test are reported in Additional file 2. This pilot test found that the reviewers in only 35.7% (5/14) of the abstracts reported or considered (i.e., dis- cussed, weighed) potential adverse effects of orthodontic interventions. This sample identified an overall prevalence of 14.3% (2/14) of spin in the abstract on adverse effects of orthodontic interventions. Both cases of spin were "misleading reportingrelated spin." The following sections describe our planned methods based on these pilot tests. We will not start the selection of eligible reviews and data extraction prior to the complete acceptance of this protocol for publication.



Eligibility criteria

We will adopt the same eligibility criteria that were defined for our published protocol on seeking adverse effects in systematic reviews of orthodontic interventions [15]. To avoid misinterpretation, we copied and pasted these eligibility criteria into Table 3 [15, 30].

Information sources

We will manually search the Cochrane library [14] and the websites of the five leading orthodontic journals to identify eligible systematic reviews published between 1 August 2009 and 31 July 2019. We chose this starting date because the first of August 2009 coincides with the launch of the Preferred Reporting Items for Systematic Reviews and

Meta-Analyses (PRISMA) statement and its guidance paper on 21 July 2009 [27, 28]. Journal selection was based on two criteria: (1) the journal has been published for 10 years or more and (2) the impact factor. The journal citation reports by Clarivate Analytics were consulted to identify the five leading orthodontic journals based on impact factor [31]. The following five orthodontic journals fulfilled both criteria: European Journal of Orthodontics (EJO), American Journal of Orthodontics and Dentofacial Orthopedics (AJODO), Angle Orthodontist, The Korean Journal of Orthodontics, and Orthodontics and Craniofacial Research.

ltem	Included	Excluded
Study designs	Systematic reviews of orthodontic interventions. The definition of systematic review, intervention review, and orthodontic interventions listed in the Glossary of terms will be used to assess whether a review is eligible (Table 1).	(1) Non-interventional reviews such as, "Methodology," 'Diagnostic," "Qualitative," and "Prognostic" (2) Rapid and scoping reviews (3) Systematic reviews that focus exclusively on adverse effects of interventions (5) Systematic reviews of interventions that did not find any eligible studies (empty reviews)
Participants	Systematic reviews on any type of patients undergoing orthodontic interventions, i.e., patients of any health status, sex, age, demographics, and socio-economic status.	 Intervention reviews that focus exclusively on patients with congenital anomalies, for example with cleft lip and palate (2) Systematic reviews of animal or laboratory studies
Interventions	(1) Systematic reviews that assess the effects of clinical orthodontic interventions. Clinical orthodontic interventions refer to the use of any type of orthodontic appliance that is used to move teeth or change the jaw size or position for orthodontic purposes (2) Systematic reviews of interventions with appliances to maintain or stabilize the outcomes of orthodontic treatment, for example, retainers (3) Systematic reviews of orthodontic interventions that compare the effects of orthodontic treatment with or without additional interventions such as pharmacological or small surgical interventions, e.g., periodontal or implant surgery (4) No exclusion criteria will be applied to the characteristics of the operator who conducted the interventions	(1) Systematic reviews in which patients receive orthodontic treatment, but in which the effects of other interventions, e.g., periodontal surgery, are compared and not the effects of orthodontic interventions (2) Systematic reviews of interventions in which orthodontic appliances are specificall used for other purposes, e.g., changing jaw positions to treat respiration or temporomandibular disorders (3) Systematic review of orthodontic interventions that included orthograthic surgery
Outcomes	(1) Any adverse effect of orthodontic interventions scored at any endpoint or timing	No exclusion criteria
	(2) The effects of orthodontic interventions do not refer just to outcomes related to tooth and juw size and positions, but also to broader outcomes such as periodontal health, esthetic changes, the health of the temporomandibular joint, patient health experiences, and economic issues associated with the interventions	
	(3) The reporting of outcomes on adverse effects will not determine the eligibility of reviews for this cross-sectional study, i.e., reviews will not be excluded because they did not provide "usable" data [30]	
Stetting	Any type of setting in which the interventions were conducted, i.e., university or private practice, etc.	No exclusion criteria

Study records

Walter & Plantation and and

Data management

- Two authors (PS and RMR) will conduct all study selection and data extraction procedures independently.
- Pilot tests were conducted to train both reviewers in applying these methods consistently and for calibration purposes [28].
- We will apply the following strategies in the case of disagreement between the two authors on the eligibility of a paper or the extraction of data: (1) discussions between

reviewers, (2) rereading the paper, (3) or if necessary contacting its authors [32]. Persistent disagreements will be resolved through the consultation and arbitration of a methodologist (NDG).

• All eligible systematic reviews will be downloaded, and all extracted data will be collected in an Excel spreadsheet.

Selection process

- All titles and abstracts in the websites of the five orthodontic journals will be handsearched to identify eligible reviews. The section "Dentistry and Oral health" will be searched in the Cochrane library for eligible Cochrane reviews [14].
- We will only include the latest version of a review when updates have been published.
- Authors will be contacted in the case of doubt regarding multiple publications of the same review. We plan to include the first publication, but will make this decision on a case by case basis and will report the rationale for this choice.
- Our selection procedures will be presented in a PRISMA flow diagram [32, 33].
- All included and excluded studies will be presented in tables, and the rationale for exclusion will be given for each excluded review.

Data collection process

- All eligible studies together with their supplemental files will be merged into binder PDFs, and pertinent search terms are linked to these documents to facilitate data extraction [34, 35].
- Eligible search terms were identified through searches in thesauri and in key articles on adverse effects [13, 36–40]. These terms are given in Additional file 3 and are identical to those used in our protocol on seeking adverse effects in systematic reviews of orthodontic interventions [15].
- Our pilot-tested data collection forms will be used for all data extraction procedures (Additional file 4). The PRISMA [32, 33] and the PRISMA-P [27, 28] checklists and guidance were consulted to develop these forms. The criteria for scoring pertinent data items are defined in these forms.
- The entire eligible review of both orthodontic and Cochrane reviews will be searched for data items, i.e., the text, tables, figures, and all supplemental files. The plain language summary in the eligible Cochrane reviews will not be searched for data items.
- When during the data extraction procedure changes are made in the data collection forms, we will present this with rationale in the section "Differences between the protocol and review."

Scoring adverse effects of orthodontic interventions

• We will use the framework of categories of known orthodontic adverse effects as defined by Preoteasa et al. [24] (Table 2). We will also include pain as a result of tooth movement as an adverse effect. Potential adverse effects that are identified

during data extraction will be discussed between the two reviewers (PS and RMR). We will report the rationale when including additional adverse effects and will categorize them.

 Ambiguous outcomes that could be interpreted as either a beneficial or an adverse outcome will not be scored as "adverse." The rationale for this score will be given. We will only consider ambiguous outcomes as "adverse" when the review authors define these outcomes as such and make a strong case for this classification.

Scoring spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions

- We will assess three types of spin, i.e., misleading reporting, misleading interpretation, and misleading extrapolation on adverse effects of orthodontic interventions in the abstract (Table 4). Each type of spin will be assessed separately for reviews that either did or did not seek adverse effects of interventions.
- To facilitate our scoring procedures and to reduce the risk of misinterpretation, we subdivided each type of spin into categories and defined each category (Table 4). We will score the presence of spin when spin is identified for one or more of these categories. The scoring procedures are summarized in Additional file 4. Pilot tests were conducted to assess the validity of these procedures.

Table 4 Types of spin in reviews that did or did not seek adverse effects of interventions.

Definitions of the three types of spin	Reviews that sought adverse effects of interventions	Reviews that did not seek adverse effects of interventions
Misleading reporting (in the abstract) on adverse effects of interventions: "Incomplete or inadequate reporting in the abstract on the results of adverse effects compared with what is reported in the main text of the manuscript, which could be misleading for the reader."	Categories: (1) Not reporting in the abstract on the results of the adverse effects that were reported in the main text of the review. (2) Selective reporting in the abstract on the results of the adverse effects that were reported in the main text of the review.	Categories: (1) Reporting on results of adverse effects in the abstract when adverse effects were not sought. (2) Reporting in the abstract that adverse effects were sought when they were not sought.
Misleading interpretation (in the abstract) on adverse effects of interventions: "Interpretation in the abstract on the results of adverse effects that is not consistent with what is reported in the main text of the manuscript and underestimates the adverse effects of the intervention."	Categories: (1) Claiming in the abstract that the intervention is safe (has no or minimal adverse effects), despite concerning results on the adverse effects in the main text of the review, e.g., based on non-statistically significant results on adverse effects with wide confidence intervals [17]. (2) Downgrading in the abstract the importance of the adverse effects, despite concerning results on the adverse effects, despite concerning results on the adverse effects in the main text of the review. (3) Recommendations are made in the abstract for clinical practice that are not congruent with the concerning results on the adverse effects in the main text of the review [17].	Categories: (1) Claiming in the abstract that the intervention is safe (has no or minimal adverse effect) despite not having sought adverse effects. (2) Downgrading in the abstract the importance of the adverse effects, despite not having sought adverse effects. (3) Recommendations are made in the abstract for clinical practice despite not having sought adverse effects.
Misleading extrapolation (in the abstract) on adverse effects of interventions: "Overgeneralisation in the abstract of the study results to different populations, interventions, outcomes or settings than were assessed in the study despite evidence in the main text on concerning adverse effects on a different population, intervention, outcome or setting."	Categories: (1) Results are extrapolated in the abstract to another population, intervention, outcome, or setting than were assessed in the review despite evidence in the main text on concerning adverse effects on a different population, intervention, outcome or setting.	Categories: (1) Results are extrapolated in the abstract to another population, intervention, outcome, or setting than were assessed in the review despite not having sought adverse effects.

Outcomes and statistical analyses

- Figure 2 a and b present all research questions in a flow diagram, and Table 5 lists all planned outcomes.
- We will calculate and report all prevalence data with their 95% confidence levels.
- We calculate the prevalence statistics for (1) all journals as one group, (2) the group of five leading orthodontic journals and the Cochrane Database of Systematic Reviews separately, and (3) each individual journal separately. Generalized linear models will be developed having the following outcomes for the abstracts of systematic reviews of orthodontic interventions: the reporting or considering of potential adverse effects of interventions/no reporting or considering of potential adverse effects of interventions (binary); presence of SPIN/absence of "SPIN" (binary); and misleading reporting/misleading interpretation/misleading extrapolation/no SPIN (categorical). The models will account for journal category (Cochrane Database of Systematic Reviews vs others), individual journals, and the geographical location of the study. Statistical significance will be based on a p value < 0.05. Stata software (Stata Corporation, College Station, TX, USA) version 15 will be used for all the statistical analyses [41].
- All outcomes that will be introduced or eliminated post hoc will be reported together with the rationale for inclusion or exclusion.

Reporting of the research study and data management

- The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement will be used as the guideline for reporting the completed cross-sectional study [42].
- A data management plan was prepared for the long-term storage of our research data [43] in the case that the publisher of our completed research study will not or will only partly store our raw data. We consulted the Registry of Research Data Repositories [44] to identify an appropriate repository for our type of research data. We selected Dryad [45] for two reasons: (1) it is an international repository of data of peer-reviewed scientific and medical research and (2) it also includes data sets for which no specific data repository exist such as meta- epidemiological research data of systematic reviews in orthodontics. Our data management plan implies that (1) all our research data will be made freely available, (2) our completed article will present a link to a repository in which all raw data of the study will be deposited, (3) the repository is registered in the Registry of Research Data Repositories [44], (4) our research data will be reported in a format that permits other researchers to understand, cite, and reuse these data, (5) all sensitive data will be protected, and (6) it will be reassessed frequently and also updated if necessary [43, 44].

Differences between the protocol and the completed study

- We will report all modifications between the protocol and the final research study. The rationale for each of these changes will be given.
- We will also report the consequences of these modifications on the magnitude, direction, and the validity of the outcomes [46].




Table 5 Summary of findings	
Description of outcomes from the main text	Statistic
The number of retrieved systematic reviews	Number
The number of eligible systematic reviews	Number
The prevalence of eligible systematic reviews	Prevalence
The prevalence of eligible systematic reviews that did seek any findings related to adverse effects of interventions in the included studies	Prevalence
The prevalence of eligible systematic reviews in which potential adverse effects of these interventions were reported or considered (i.e., discussed, weighted, etc.) in the abstract*	Prevalence
The prevalence of eligible systematic reviews in which spin was identified on adverse effects of orthodontic interventions in the abstract*	Prevalence
The prevalence of misleading reporting-related spin in the abstract*	Prevalence
The prevalence of misleading interpretation-related spin in the abstract*	Prevalence
The prevalence of misleading extrapolation-related spin in the abstract*	Prevalence

All prevalence data will be presented with their 95% confidence intervals

"This statistic will be reported for reviews that sought and did not seek any findings related to adverse effects of interventions in the included studies

Discussion

Strengths

Key strengths of this research study include the following: (1) we conducted extensive scoping searches and pilot studies to fine-tune our research questions and methods. These activities confirmed the importance of our questions. (2) Our research team consists of two topic experts (PS and RMR) and two methodologists (RMR and NDG). (3) All study selection and data collection procedures will be undertaken independently by two authors (PS and RMR). Calibration of these operators was done during the pilot studies. (4) To guarantee reproducibility and full access to our data, we will publish our protocol a priori and will include all raw data of the completed research study in additional files or will deposit them in an open-access repository [43–45, 47].

Limitations

Including only orthodontic intervention reviews published in the five leading orthodontic journals and in the Cochrane Database of Systematic Reviews could be a limitation, but we expect that the findings in this subgroup of journals will underestimate the true severity of spin on adverse effects of interventions in the abstracts of these reviews. Including only reviews published in the last 10 years could also be a limitation. However, we chose this period because it brings the current knowledge status on our research questions to the foreground and these 10 years coincide with the launch in 2009 of the checklist of Preferred Reporting Items for Systematic Reviews and Meta- Analyses (PRISMA) [32, 33].

Importance and beneficiaries

In this research study, we will address three key questions in abstracts of systematic reviews of orthodontic interventions: whether potential adverse effects of these interventions were reported or considered, whether spin was identified regarding information on these adverse effects, and the type of spin. These issues are important, because (1) the assessment and

reporting of adverse effects of interventions is often suboptimal [7–11], (2) titles and abstracts are the most read sections of papers in the biomedical literature [1], (3) a high prevalence of spin has been identified in abstracts of both randomized and non-randomized studies [4, 21], and (4) incomplete or inadequate reporting, interpretation, or extrapolation of findings on adverse effects in the abstract can mislead readers and could lead to inadequate practice [4]. Our results will raise the awareness of considering adverse effects and the phenomenon of spin regarding these effects in abstracts of systematic reviews of orthodontic interventions. Patients, clinicians, researchers, editors, peer-reviewers, guideline developers, policy makers, and research funders will all be beneficiaries of the findings of this research study.

Supplementary information

Supplementary information accompanies this paper at <u>https://doi.org/10.1186/s41073-019-0084-4</u>.

Additional file 1. Checklist for the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P) 2015 statement. Additional file 2. Pilot tests.

Additional file 3. Search terms and their derivatives. Additional file 4. Data collection forms.

Abbreviations

MECIR: Methodological Expectations of Cochrane Intervention Reviews; PRISMA: Preferred reporting items for systematic reviews and meta-analyses; PRISMA-P: Preferred Reporting Items for Systematic review and Meta- Analysis-Protocols

Acknowledgements

Not applicable

Authors' contributions

PS and RMR conceived and designed the study protocol for this cross-sectional study. RMR is the guarantor. PS and RMR conducted the pilot testing of the study selection procedures and data extraction forms and fine-tuned the research protocol after the pilot testing. NDG provided support on methodological and statistical issues and assisted in the overall finetuning of this protocol. All three authors read and approved the final protocol.

Funding

All expenses for preparing this protocol and for conducting the subsequent research study will be paid evenly by each author.

Availability of data and materials

Not applicable

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Orthodontics, Academisch Centrum Tandheelkunde Amsterdam (ACTA), University of Amsterdam, Gustav Mahlerlaan 3004, 1081, LA, Amsterdam, The Netherlands. ²Center for Veterinary Health Sciences, Oklahoma State University, 2065 W Farm Road, Stillwater, OK 74078, USA.

³EBMVet, Via Sigismondo Trecchi 20, 26100 Cremona, CR, Italy.

⁴Department of Oral and Maxillofacial Surgery, Academic Medical Center, University of Amsterdam, Meibergdreef 9, 1105, AZ, Amsterdam, The Netherlands. ⁵Studio di Ortodonzia, Via Matteo Bandello 15, 20123 Milan, Italy.

Received: 13 April 2019 Accepted: 10 October 2019

Published online: 19 December 2019

References

- Dogan RI, Murray GC, Névéol A, Lu Z. Understanding PubMed user search behavior through log analysis. Database (Oxford). 2009;2009:bap018 Epub 2009 Nov 27.
- 2. Boutron I, Ravaud P. Misrepresentation and distortion of research in biomedical literature. Proc Natl Acad Sci U S A. 2018;115(11):2613–9. https:// doi.org/10.1073/pnas.1710755115.
- Haneef R, Yavchitz A, Ravaud P, Baron G, Oranksy I, Schwitzer G, Boutron I. Interpretation of health news items reported with or without spin: protocol for a prospective meta-analysis of 16 randomised controlled trials. BMJ Open. 2017;7(11):e017425. <u>https://doi.org/10.1136/bmjopen-2017-017425</u>.
- Lazarus C, Haneef R, Ravaud P, Boutron I. Classification and prevalence of spin in abstracts of nonrandomized studies evaluating an intervention. BMC Med Res Methodol. 2015;15:85. <u>https://doi.org/10.1186/s12874-015-0079-x.</u>
- Higgins JPT, Lasserson T, Chandler J, Tovey D, Churchill R. Methodological Expectations of Cochrane Intervention Reviews (MECIR). Standards for the conduct and reporting of new Cochrane Intervention Reviews, reporting of protocols and the planning, conduct and reporting of updates. Cochrane: London, 2016. Available from: https:// methods.cochrane.org/sites/default/files/public/uploads/mecir_printed_ booklet_final_v1.02.pdf

methods.cochrane.org/sites/default/files/public/uploads/mecir_printed_booklet_final_v1.02.pdf (accessed 18 Feb 2019).

- Parsons R, Golder S, Watt I. More than one-third of systematic reviews did not fully report the adverse events outcome. J Clin Epidemiol. 2019;108:95–101. <u>https://doi.org/10.1016/j.jclinepi.2018.12.007</u>.
- Bagul NB, Kirkham JJ. The reporting of harms in randomized controlled trials of hypertension using the CONSORT criteria for harm reporting. Clin Exp Hypertens. 2012;34(8):548–54. <u>https://doi</u>.org/10.3109/10641963.2012.681724 Epub 2012 May 9.
- Hodkinson A, Kirkham JJ, Tudur-Smith C, Gamble C. Reporting of harms data in RCTs: a systematic review of empirical assessments against the CONSORT harms extension. BMJ Open. 2013;3(9):e003436. <u>https://doi</u>.org/10. 1136/bmjopen-2013-003436.
- Péron J, Maillet D, Gan HK, Chen EX, You B. Adherence to CONSORT adverse event reporting guidelines in randomized clinical trials evaluating systemic cancer therapy: a systematic review. J Clin Oncol. 2013;31(31): 3957–63. <u>https://doi.org/10.1200/JCO.2013.49.3981</u> Epub 2013 Sep 23.
- 10. Pitrou I, Boutron I, Ahmad N, Ravaud P. Reporting of safety results in published reports of randomized controlled trials. Arch Intern Med. 2009; 169(19):1756–61.
- Smith SM, Chang RD, Pereira A, Shah N, Gilron I, Katz NP, Lin AH, McDermott MP, Rappaport BA, Rowbotham MC, Sampaio C, Turk DC, Dworkin RH. Adherence to CONSORT harms-reporting recommendations in publications of recent analgesic clinical trials: an ACTTION systematic review. Pain. 2012;153(12):2415–21. <u>https://doi.org/10.1016/j.pain.2012.08.009</u> Epub 2012 Sep 15.
- 12. Glossary of terms in the Cochrane Collaboration. Version 4.2.5. Updated May 2005. Available from: <u>http://aaz</u>.hr/resources/pages/57/7.%20Cochrane%2 0glossary.pdf (accessed 27 Sept 2019).
- Loke YK, Price D, Herxheimer A. Chapter 14: adverse effects. In: Higgins JPT, Green S. editors. Cochrane handbook for systematic reviews of interventions version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from: <u>http://handbook</u>-5-1.cochrane.org/ (accessed 29 Aug 2019).
- 14. Cochrane library. Available from: <u>https://www</u>.cochranelibrary.com/about/ about-cochrane-reviews (accessed 24 Sept 2019).
- Steegmans PAJ, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross- sectional study. Syst Rev. 2019;8(1):89. <u>https://doi</u>.org/10.1186/s13643-019- 1000-1.
- 16. Horton R. The rhetoric of research. BMJ. 1995;310(6985):985-7.
- Yavchitz A, Ravaud P, Altman DG, Moher D, Hrobjartsson A, Lasserson T, Boutron I. A new classification of spin in systematic reviews and meta- analyses was developed and ranked according to the severity. J Clin Epidemiol. 2016;75:56–65. <u>https://doi</u>.org/10.1016/j.jclinepi.2016.01.020 Epub 2016 Feb 2.
- Boutron I, Dutton S, Ravaud P, Altman DG. Reporting and interpretation of randomized controlled trials with statistically nonsignificant results for primary outcomes. JAMA. 2010;303(20):2058–64. <u>https://doi</u>.org/10.1001/ jama.2010.651.

- Haneef R, Lazarus C, Ravaud P, Yavchitz A, Boutron I. Interpretation of results of studies evaluating an intervention highlighted in Google health news: a cross-sectional study of news. PloS One. 2015;10(10):e0140889. <u>https://doi</u>. Org/10.1371/journal.pone.0140889 eCollection 2015.
- Latronico N, Metelli M, Turin M, Piva S, Rasulo FA, Minelli C. Quality of reporting of randomized controlled trials published in Intensive Care Medicine from 2001 to 2010. Intensive Care Med. 2013;39(8):1386–95. <u>https://doi</u>.org/10.1007/s00134-013-2947-3 Epub 2013 Jun 7.
- Lockyer S, Hodgson R, Dumville JC, Cullum N. "Spin" in wound care research: the reporting and interpretation of randomized controlled trials with statistically non-significant primary outcome results or unspecified primary outcomes. Trials. 2013;14:371. <u>https://doi.org/10.1186/1745-6215-14-371.</u>
- Ochodo EA, de Haan MC, Reitsma JB, Hooft L, Bossuyt PM, Leeflang MM. Overinterpretation and misreporting of diagnostic accuracy studies: evidence of "spin". Radiology. 2013;267(2):581–8. <u>https://doi.org/10.1148</u>/ radiol.12120527 Epub 2013 Jan 29.
- 23. World Medical Association Declaration of Helsinki Ethical Principles for Medical Research Involving Human Subjects. Research registration and publication and dissemination of results. Principle 36. 64th WMA General Assembly, Fortaleza, Brazil, October 2013. Published Online: October 19, 2013. Doi:10.1001/jama. 2013.281053. Available from: <u>https://www</u>.wma.net/wp-content/uploads/2 016/11/DoH-Oct2013-JAMA.pdf (accessed 28 Sept 2019).
- Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F. (editor). Orthodontics-Basic aspects and clinical considerations. March 9, 2012 under CC BY 3.0 license. <u>www.intechopen.com</u>. Available from: <u>https://www</u>. Intechopen.com/books/orthodontics-basic-aspects-and-clinical- considerations/risks-andcomplications-associated-with-orthodontic- treatment (accessed 29 Sept 2019).
- 25. Reeves BC. Reporting of harms in systematic reviews and their primary studies. BMJ. 2014;349:g6819. <u>https://doi</u>.org/10.1136/bmj.g6819.
- 26. Ioannidis JP. The mass production of redundant, misleading, and conflicted systematic reviews and meta-analyses. Milbank Q. 2016;94(3):485–514.
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. Preferred Reporting Items for Systematic Review and Meta- Analysis Protocols (PRISMA-P) 2015 statement. Syst Rev. 2015;4(1):1.
- Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA, the PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;349:g7647.
- Viechtbauer W, Smits L, Kotz D, Budé L, Spigt M, Serroyen J, Crutzen R. A simple formula for the calculation of sample size in pilot studies. J Clin Epidemiol. 2015;68(11):1375–9. https://doi.org/10.1016/j.jclinepi.2015.04.014 Epub 2015 Jun 6.
- O'Connor D, Green S, Higgins JPT (editors). Chapter 5: Defining the review question and developing criteria for including studies. In: Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. [online] Available from: <u>www.cochrane</u>-handbook.org. (accessed 29 Sept 2019).
- 31. Clarivate Analytics. Available from: https://clarivate.com/ (accessed 29th Sept 2019).
- 32. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. PloS Med. 2009;6:e1000100 Epub 2009 Jul 21.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PloS Med. 2009;6:e1000097.
- Adobe Merge PDFs, combine files into one PDF. Available from: http:// <u>www.wikihow</u>.com/Merge-PDF-Files (accessed 29 Sept 2019).
- Acrobat for legal professionals. Searching and marking multiple words. [online] Available from: <u>http://blogs</u>.adobe.com/acrolaw/2010/04/searching- and-marking-multiple-words-in-a-pdf/ (accessed 29 Sept 2019).

- Golder S, Loke YK, Zorzela L. Some improvements are apparent in identifying adverse effects in systematic reviews from 1994 to 2011. J Clin Epidemiol. 2013;66(3):253–60. https://doi.org/10.1016/j.jclinepi.2012.09.013.
- Saini P, Loke YK, Gamble C, Altman DG, Williamson PR, Kirkham JJ. Selective reporting bias of harm outcomes within studies: findings from a cohort of systematic reviews. BMJ. 2014;349:g6501. <u>https://doi.org/10.1136/bmj.g6501.</u>
- Zorzela L, Golder S, Liu Y, Pilkington K, Hartling L, Joffe A, Loke Y, Vohra S. Quality of reporting in systematic reviews of adverse events: systematic review. BMJ. 2014;348:f7668. <u>https://doi</u>.org/10.1136/bmj.f7668 Review.
- Zorzela L, Loke YK, Ioannidis JP, Golder S, Santaguida P, Altman DG, Moher D, Vohra S, PRISMA Harms Group. PRISMA harms checklist: improving harms reporting in systematic reviews. BMJ. 2016;352:i157. <u>https://doi.org/10.1136/ bmj.i157</u>.
- 41. StataCorp. Stata Statistical Software: Release 15. College Station: StataCorp LLC.; 2017.
- 42. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Ann Intern Med. 2007;147(8):573–7.
- 43. Schiermeier Q. For the record. Making project data freely available for open science. Nature. 2018;555:403–5.
- 44. Registry of Research Data Repositories. [online] Available from: <u>https://www</u>. Re3data.org/ (accessed 29 Sept 2019).
- 45. Dryad. [online] Available from: https://datadryad.org/stash (accessed 28 Sept 2019).
- 46. Higgins JPT, Deeks JJ, Altman DG (editors). Chapter 16: Special topics in statistics. In: Higgins JPT, Green S (editors), Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. [online] Available from: http:// handbook-5-1.cochrane.org/ (accessed 29 Sept 2019).
- Goodman SN, Fanelli D, Ioannidis JP. What does research reproducibility mean? Sci Transl Med. 2016;8(341):341ps12. <u>https://doi.org/10.1126/</u> scitranslmed.aaf5027.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



CHAPTER 5

Spin on adverse effects in abstracts of systematic reviews of orthodontic interventions: a cross-sectional study (part 2)

This is a non-edited version of the manuscript:

Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin on adverse effects in abstracts of systematic reviews of orthodontic interventions: a cross-sectional study (part 2). Syst Rev. 2023 Jun 20;12(1):99.

Abstract

Background It is critical that abstracts of systematic reviews transparently report both the beneficial and adverse effects of interventions without misleading the readers. This cross-sectional study assessed whether adverse effects of interventions were reported or considered in abstracts of systematic reviews of orthodontic interventions and whether spin on adverse effects was identified when comparing the abstracts with what was sought and reported in these reviews.

Methods This cross-sectional study (part 2 of 2) used the same sample of 98 systematic reviews orthodontic interventions as used in part 1. Eligible reviews were retrieved from the Cochrane Database of Systematic Reviews and the 5 leading orthodontic journals between August 1 2009 and July 31 2021. Prevalence proportions were sought for 3 outcomes as defined in the published protocol. Univariable logistic regression models were built to explore associations between the presence of spin in the abstract and a series of predictors. Odds ratios (OR) 95% confidence intervals (95% CI) were used to quantify the strength of associations and their precision.

Results 76.5% (75/98) of eligible reviews reported or considered (i.e., discussed, weighted etc.) potential adverse effects of orthodontic interventions in the abstract and the proportion of spin on adverse effects was 40.8% (40/98) in the abstract of these reviews. Misleading reporting was the predominant category of spin, i.e., 90% (36/40). Our explorative analyses found that compared to the Cochrane Database of Systematic Reviews all 5 orthodontic journals had similar odds of the presence of spin on adverse effects in abstracts of systematic reviews of orthodontic interventions. The odds of the presence of spin did not change over the sampled years (OR: 1.03, 95% CI: 0.9 to 1.16) and did not depend on the number of authors (OR: 0.93, 95% CI: 0.71 to 1.21), or on the type of orthodontic intervention (OR: 1.1, 95% CI: 0.45 to 2.67), or whether conflicts of interests were reported (OR: 0.74, 95% CI: 0.32 to 1.68).

Conclusion End users of systematic reviews of orthodontic interventions have to be careful when interpreting results on adverse effects in the abstracts of these reviews, because they could be jeopardized by uncertainties such as not being reported and misleading reporting as a result of spin.

Keywords Orthodontics, Reporting, Systematic review, Intervention, Spin, Misleading reporting, Misleading interpretation, Misleading extrapolation, Adverse effect, Adverse event, Harm, Safety

Background

Abstracts should provide key information on a research study, which helps readers decide whether or not to access the full report [1]. It is therefore critical that abstracts transparently report the results of both the beneficial and adverse effects of interventions without misleading the readers. Misleading reporting, misleading interpretation, and misleading extrapolation of study results has been called "spin" [2, 3]. In this study, we assessed whether adverse effects of interventions were reported or considered in abstracts of both Cochrane and non-Cochrane reviews of orthodontic interventions and whether spin and what type of spin regarding adverse effects was present when comparing the abstracts with what was sought and reported in these reviews.

Titles and abstracts of publications of healthcare interventions are used for multiple purposes such as (1) an initial screening of the study type; (2) clarifying the included type of patients, interventions, comparators, outcomes, and settings; (3) obtaining a summary of the findings; and (4) an initial assessment of the validity of the study [1, 4, 5]. Titles and abstracts are the most and often only read sections of biomedical papers, because of a lack of time of readers, paywalls, or language issues [6]. It is therefore important that abstracts can be used as stand-alone documents that clearly and truthfully reflect what was reported in the full text [7]. The standard for Methodological Expectations of Cochrane Intervention Reviews (MECIR) [8] states under Item R13 that "The abstract of the review should aim to reflect a balanced summary of the benefits and harms of the intervention" and this is a "mandatory" Cochrane review standard. The inclusion of 'adverse effects' in this standard is crucial, because these effects are often poorly assessed and reported in clinical trials and systematic reviews of healthcare interventions [9–16].

In this context, it is important that findings on adverse effects are presented accurately in the abstract without misleading the reader. "A distorted presentation of study results" has been called "spin" [2, 3]. This definition and other commonly used definitions of spin and key terminology used in this article are listed in Table 1 [2, 3, 17–25].

Spin has been subdivided in 3 categories: "misleading reporting," "misleading interpretations," and "misleading extrapolations" of study results [2]. We adopted the definitions by Lazarus et al. [23] for these 3 categories of spin (Table 1). Controlling spin is important, because of its high prevalence and its consequences. For example, a randomized controlled trial (RCT) showed that spin in abstracts can influence the clinician's interpretation of the results of a study [26]. Further, Yavchitz et al. [27] showed that the presence of spin in press releases and the mass media was related with spin in the conclusions of the pertinent abstracts of peer-reviewed RCTs. Our scoping searches showed that a high prevalence of spin has been recorded in abstracts of numerous research studies and for a wide variety of disciplines. For example, spin was present in 84% (107/128) of abstracts of reports of non-randomized studies assessing an intervention [23], 23% (24/105) of abstracts of RCTs in rheumatology [28], 57% (53/93) of abstracts of cardiovascular RCT reports [29], 34.2% (25/73) of abstracts of systematic reviews and meta-analyses related to treatment of proximal humerus fractures [30], 37.6% (27/72) of results, and 58.3% (42/72) of conclusions of abstracts of parallel-group RCTs with statistically non-significant results $(P \ge 0.05)$ [25]. Spin in abstracts of orthodontic studies was assessed in 2 recent publications

[31, 32]. Guo et al. [31] found spin in 62.2% (69/111) of abstracts of parallel-group RCTs with clearly stated statistically non-significant primary outcomes and Makou et al. [32] identified spin in 48.6% (53/109) of abstracts of orthodontic meta-analyses.

This is part 2 of 2 cross-sectional studies on assessing and reporting of adverse effects in systematic reviews of orthodontic interventions published in 5 leading orthodontic journals and in the Cochrane Database of Systematic Reviews. Part 1 focused predominantly on seeking and reporting of adverse effects in the main text and supplementary files of these reviews [18, 33]. In part 2, we assessed whether adverse effects of orthodontic interventions were reported or considered (i.e., discussed, weighed etc.) in abstracts of these reviews. We further measured whether spin was introduced in the abstract regarding information on adverse effects as found and reported in these reviews. We also assessed the different categories of spin. The findings of this research study are important not only for patients and clinicians but also for researchers, peer reviewers, and editors because they have a crucial role in reducing the prevalence of spin [34].

Term	Definition
Systematic review	Cochrane [17] defines a systematic review as follows: 'A systematic review attempts to identify, appraise and synthesize all the empirical evidence that meets pre-specified eli- gibility criteria to answer a specific research question. Researchers conducting systematic reviews use explicit, systematic methods that are selected with a view almed at minimiz- ing bias, to produce more reliable findings to inform decision making."
Intervention review	Cochrane [17] defines an intervention review as follows: "Intervention reviews assess the effectiveness/safety of a treatment, vaccine, device, preventative measure, procedure or policy."
Orthodontic interventions	Steegmans et al. [18] define orthodontic interventions as follows: "Orthodontic interven- tions refer to the use of any type of orthodontic appliance to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodontic treatment, for example retainers."
Adverse effect	Cochrane [19, 20] defines an adverse effect as "an adverse event for which the causal rela- tion between the intervention and the event is at least a reasonable possibility."
Spin [3]	"Distorted presentation of study results."
Spin [3]	"A misrepresentation of study results, regardless of motive (intentionally or unintention- ally) that overemphasizes the beneficial effects of the intervention and overstates safety compared with that shown by the results."
Spin [21]	"A specific intentional or unintentional reporting that fails to faithfully reflect the nature and range of findings and that could affect the impression the results produce in readers"
Spin [22]	"A specific reporting that fails to faithfully reflect the nature and range of findings and that could affect the impression that the results produce in readers, a way to distort sci- ence reporting without actually lying"
Misleading reporting related spin [23]	"Incomplete reporting of the study results that could be misleading for the reader."
Misleading interpretation related spin [23]	Inadequate interpretation of the study results overestimating the beneficial effect of the intervention.
Misleading (inappropriate) extrapolation related spin [23]	Inappropriate generalization of the study results by inadequate (1) extrapolation from the population, interventions, or outcome actually assessed in the study to a larger population, different interventions, or outcomes or (2) inadequate implications for clinical practice.
Spin (in the abstract) on adverse effects of interventions [24]	Incomplete or inadequate reporting, interpretation, or extrapolation (or a combination of these variables) of findings on adverse effects of interventions in the abstract that could be misleading for the reader.

Table 1 Glossary of terms

Objectives

Our objectives were presented in the following 3 research questions [24]. Recent (up to October 31, 2021) scoping searches showed that these questions were not assessed previously.

• Question 1. In abstracts of systematic reviews of orthodontic interventions, were potential adverse effects of these interventions reported or considered (i.e., discussed, weighed etc.)?

• Question 2. Was spin identified on adverse effects of orthodontic interventions in the abstract?

• Question 3. What type of spin was identified on adverse effects of orthodontic interventions in the abstract?

Methods

This manuscript reports the methods and results of part 2 of a cross-sectional study using the same 98 eligible reviews as in part 1 [33]. Additional information on the research methods and the characteristics of the included reviews can be found in part 1 [33] and in the published protocols of parts 1 and 2 [18, 24]. The protocol for this second cross-sectional study was published in "Research Integrity and Peer Review" [24] and can be consulted via the following link:

https://researchintegrityjournal.biomedcentral.com/articles/10.1186/s41073-019-0084-4.

The checklist of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement for cross-sectional studies [35] was included as Additional file 1. The differences between the methods planned in our protocol and those implemented in the final study were reported in Additional file 2. The rationales for these differences were also given. All raw data were reported in the Open Science framework Open Science Framework (https://osf.io/ka7mp/). There was no patient or public involvement during the development of the protocol or in the conduct of this study. The eligibility criteria, information sources, search strategy, and selection process used in part 1 of this cross-sectional study [33] were also used for part 2 of this study. To reduce the need of cross-checking between manuscripts, we reported these sections again.

Eligibility criteria

The eligibility criteria were published previously in our protocol [24] and in part 1 of this study [33] and were developed by two researchers (PS and RMR). These criteria are presented in Table 2 [36] and are further explained under here.

Study designs

• We included systematic reviews of orthodontic interventions. The definitions of the terms "systematic review," "intervention review," and "orthodontic interventions" listed in the Glossary of terms (Table 1) were used to assess eligibility.

The following reviews were excluded: (1) noninterventional reviews such as "Methodology," "Diagnostic," "Qualitative," and "Prognostic"; (2) rapid and scoping reviews; (3) systematic reviews with Bayesian network meta-analysis; and (4) systematic reviews of interventions that did not find any eligible studies (empty reviews).

Participants

- Systematic reviews of interventions on any type of patients undergoing orthodontic interventions, i.e., patients of any health status, sex, age, and demo graphics, and socio-economic status were eligible.
- Intervention reviews that focused exclusively on patients with congenital anomalies, for example, with cleft lip and palate and systematic reviews of animal or laboratory studies were excluded.

Interventions

- Systematic reviews on the following interventions were eligible: (1) systematic reviews that assessed the effects of clinical orthodontic interventions. Clinical orthodontic interventions refer to any type of orthodontic appliance that are used to move teeth or change the jaw size or position for orthodontic purposes; (2) systematic reviews of interventions with appliances to maintain or stabilize the outcomes of orthodontic treatment, for example, retainers; (3) systematic reviews of orthodontic interventions such as pharmacological or small surgical interventions, e.g., periodontal or implant surgery; and (4) no exclusion criteria were applied to the characteristics of the operator who conducted the interventions.
- Systematic reviews on the following interventions were excluded: (1) systematic
 reviews in which patients receive orthodontic treatment, but in which the effects of
 other interventions, e.g., periodontal surgery, were compared and not the effects of
 orthodontic interventions; (2) systematic reviews of interventions in which
 orthodontic appliances were specifically used for other purposes, e.g., changing jaw
 positions to treat respiration or temporomandibular disorders; (3) systematic review
 of orthodontic interventions that included orthognathic surgery; (4) systematic
 reviews that focused exclusively on adverse effects of interventions; and (5)
 systematic reviews that did not assess a specific orthodontic intervention but
 referred to orthodontic treatment as a whole.

Outcomes

• Systematic reviews of orthodontic interventions that assessed any adverse effect of orthodontic interventions scored at any endpoint or timing were eligible. The effects

of orthodontic interventions did not refer just to outcomes related to tooth and jaw size and positions but also to broader outcomes such as periodontal health, esthetic changes, the health of the temporomandibular joint, patient health experiences, and economic issues associated with the interventions. The reporting of outcomes on adverse effects did not determine eligibility of reviews for this cross-sectional study, i.e., reviews were not excluded because they did not report measured outcome data in a "usable" way [36].

• No exclusion criteria regarding the outcomes of systematic reviews of orthodontic interventions were applied.

Setting

• Systematic reviews of orthodontic interventions that reported on interventions conducted in any type of setting, i.e., university or private practice, were eligible.

Item	Included	Excluded
Study designs	Systematic reviews of orthodontic interventions. The defini- tion of systematic review, intervention review, and orthodontic interventions listed in the Glossary of terms will be used to assess whether a review is eligible (Table 1).	 Noninterventional reviews such as "Methodology," Diagnostic," "Qualitative," and "Prognostic" Rapid and scoping reviews Systematic reviews of interventions that did not find any eligible studies (empty reviews)
Participants	Systematic reviews on any type of patients undergoing ortho- dontic interventions, i.e., patients of any health status, sex, age, and demographics, and socio-economic status.	1) Intervention reviews that focus exclusively on patients with congenital anomalies, for example with cleft lip and palate 2) Systematic reviews of animal or laboratory studies
Interventions	 Systematic reviews that assessed the effects of clinical ortho- dontic interventions. Clinical orthodontic interventions refer to any type of orthodontic appliance that are used to move teeth or change the jaw size or position for orthodontic purposes 2) Systematic reviews of interventions with appliances to maintain or stabilize the outcomes of orthodontic treatment, for example retainers Systematic reviews of orthodontic interventions that com- pared the effects of orthodontic interventions that com- pared the effects of orthodontic treatment with or without addi- tional interventions, e.g., periodontal or implant surgery No exclusion criteria were applied to the characteristics of the operator who conducted the interventions 	 Systematic reviews in which patients receive orthodontic treatment, but in which the effects of other interventions, e.g., peri- odontal surgery, were compared and not the effects of orthodon- tic interventions Systematic reviews of interventions in which orthodontic appli- ances were specifically used for other purposes, e.g., changing jaw positions to treat respiration or tempororandibular disorders Systematic review of orthodontic interventions that included orthognathic surgery Systematic reviews that focused exclusively on adverse effects of interventions Systematic reviews that did not assess a specific orthodontic intervention but referred to orthodontic treatment as a whole
Outcomes	 Any adverse effect of orthodontic interventions scored at any endpoint or timing The effects of orthodontic interventions did not refer just to outcomes related to tooth and jaw size and positions but also to broader outcomes such as periodontal health, esthetic changes, the health of the temporomandibular joint, patient health expe- riences, and economic lisues associated with the interventions The reporting of outcomes on adverse effects did not determine eligibility of reviews for this cross-sectional study, i.e., reviews were not excluded because they did not report meas- ured outcome data in a "usable" way [36]. 	No exclusion criteria
Stetting	Any type of setting in which the interventions were conducted, i.e., university or private practice etc.	No exclusion criteria

Information sources and search strategy

The Cochrane Database of Systematic reviews [37] and the websites of 5 leading orthodontic journals were the information sources of this study. The journal selection of the latter journals was based on two criteria: (1) the journal has been published for 10 years or more

and (2) the highest impact factor. The following 5 orthodontic journals fulfilled these criteria: European Journal of Orthodontics [EJO], American Journal of Orthodontics and Dentofacial Orthopedics [AJODO], Angle Orthodontist (AO), The Korean Journal of Orthodontics (KJO), and Orthodontics and Craniofacial Research (OCR). These journals were manually searched from August 1, 2009, until July 31, 2021, for systematic reviews that fulfilled the eligibility criteria. August 1, 2019, was chosen as the starting date, because it coincides with the launch of the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement on 21 July 2009 [38, 39].

Study records

Selection process

Two reviewers (PS and RMR) manually searched systematic reviews that fulfilled the eligible criteria. Pilot tests were conducted a priori to train both reviewers and to calibrate them. All titles and abstracts in the websites of the 5 orthodontic journals were hand-searched for eligible reviews. Eligible Cochrane reviews were searched in the section "Dentistry and Oral health" in the Cochrane library. Only the latest version of a review was eligible when review updates had been published. In the case of disagreement on the selection procedures, the following strategies were implemented and in this sequence: (1) discussions between these operators, (2) rereading the paper, (3) contacting of authors by email to clarify issues regarding a specific manuscript. Persistent disagreements were resolved through consultation with a methodologist. A total of 98 eligible systematic reviews of orthodontic interventions was identified in part 1 of this study [33]. This same sample of 98 studies was also used in this study.

Data collection procedures

All 98 eligible reviews together with their supplemental files were merged into binder PDFs, and according to protocol [24], pertinent search terms were linked to these documents to facilitate data extraction (Additional file 2). Our pilot-tested data collection forms were used to extract data and are given in Additional file 2. Data items were collected from the entire eligible review, i.e., the entire manuscript including the abstract, tables, figures, and additional files. We implemented this procedure for all eligible reviews but did not extract data from the plain language summary of eligible Cochrane reviews. Two calibrated authors (PS and RMR) independently collected data from the 98 eligible reviews to address the research questions. In the case of disagreement, we applied the same strategies as reported in the section "Selection process" and the third author (NDG) was consulted in the case of persistent disagreements.

Assessing adverse effects of orthodontic interventions

Pain and the various categories of adverse effects hypothetically linked to orthodontic interventions as defined by Preoteasa et al. [40] and modified by Steegmans et al. [33] were reported in a table in Additional file 2. This table was consulted as our reference to assess the reporting on adverse effects in the abstract. When additional adverse effects were identified that were not given in this table, we included them with rationale. Effects that

could be labeled either as "beneficial" or "adverse" were not included unless the review authors labeled these ambiguous effects as "adverse." Explanations for such decisions were given. Orthodontic interventions were classified in three types, i.e., type 1: orthodontic interventions to move teeth or change the jaw size or position for orthodontic purposes. type 2: orthodontic interventions with additional surgical, pharmacological, or vibratory interventions, and type 3: orthodontic interventions to maintain or stabilize orthodontic results.

Assigning spin of adverse effects of orthodontic interventions in abstracts of systematic reviews

Spin was assigned by comparing whether what was reported in the abstract on adverse effects of orthodontic interventions was congruent with the findings on these effects in the review. Three types of spin were assigned i.e., misleading reporting, misleading interpretation, and misleading (inappropriate) extrapolation on adverse effects of orthodontic interventions in abstracts of systematic reviews [27]. To facilitate this assignment and to reduce the risk of misinterpretation, each type of spin was subdivided in categories. The presence of spin was assigned when it was identified in one or more of these categories. Spin was assessed in all eligible reviews irrespective of whether these reviews sought adverse effects of interventions or not. Because the pilot tests for our protocol identified only 2 reviews with spin [24] and because assessing spin is not easy [22], we conducted additional pilot tests on 10 RCTs to further calibrate the operators (PS and RMR) that assigned spin and to fine-tune the descriptions of spin and the checklists for assigning spin. These fine-tuned descriptions of the different types of spin and the pertinent data collection forms to identify spin are reported respectively in Table 3 and Additional file 2. Definitions of spin were given for reviews that sought and those that did not seek adverse effects of orthodontic interventions (Table 3).

Table 3 Types of spin in reviews that did or did not seek adverse effects of interventions

Definitions of the 3 types of spin Reviews that sought adverse effects of interventions Reviews that did not seek adverse effects of interventions deading reporting (in the abstract) on adverse effects of 1) Not reporting in the abstract on the results of adverse effects Categories: 1) Reporting orting on results of adverse effects in the abstract when found in the review. 2) Selective reporting in the abstract on the results of adverse adverse effects were not sought. 2) Reporting in the abstract that adverse effects were sought plete or inadequate reporting in the abstract on the of adverse effects that were not supported by the findings of the review" effects found in the review when they were not sought. isleading interpretation (in the abstract) on adverse Categories: Categories: Claming in the abstract that the intervention is safe (has no or 11CL inimal advense effects), despite concerning results on advense fects found in the review, e.g., based on non-statistically sig-Claiming in the abstract that the intervention is safe thas no minimal adverse effects! despite not having sought adverse. ntions spretation in the abstract on the results of adverse effects hat was not supported by the findings of the review."This could ects. inficant results on adverse effects with wide confidence intervals. 2) Downgrading in the abstract the importance of the adverse for example underestimated the adverse effects of the interven-2) Downgrading in the abstract the importance of the adver 3) Recor effects, despite concerning results on adverse effects found in despite not having sought adverse effects. the review

iate) extrapolation (in the abstract)

effects of it

idverse effects of interventions: ingeneralisation in the abstract of the study results to different populations, interventions, outcomes or settings than were sed in the review despite evidence on adverse effects on a different population, intervention, outcome or setting*

Categories 1) Results are extrapolated in the abstract to another population, intervention, outcome, or setting than were assessed in the review despite evidence on adverse effects on a different. population, intervention, outcome, or setting,

Ti Recommendations are made in the abstract for clinical practice that are not supported by the findings in the review on adverse effects [27].

ects, despite not having sought advense effects. Recommendations are made in the abstract for clinical practice

Categories: 1) Results are entropolated in the abstract to another popula-tion, intervenion, outcome, or setting than were assessed in the review dispite not having sought adverse effects.

Power calculation

In our pilot sample, we identified an overall proportion of 14.3% (2/14) of spin of the adverse effects in the abstracts of 14 systematic reviews of orthodontic interventions. For this proportion, the Epitools software [41] calculated a required sample size of 48 studies (precision 0.1 and confidence level 0.95), which was fulfilled by our 98 eligible reviews.

Outcomes and statistical analyses

Outcomes

Prevalence proportions were calculated to quantify the answers to our 3 research questions in the 98 selected reviews. According to our published protocol, these proportions were also calculated separately for reviews that either did (n = 84) or did not (n = 14) seek any findings related to adverse effects of interventions in the included studies [33]. These statistics were calculated for (1) all journals as a one group, (2) the five leading orthodontic journals together and the Cochrane Database of Systematic Reviews separately, and (3) each eligible journal separately.

Explorative analyses

Univariable logistic regression models were built to determine the association between the presence of spin in the abstract and characteristics of the systematic review, i.e., journal, year of publication, number of authors, conflict of interest reported, conflict of interest present, funding reported, and type of orthodontic intervention. These analyses were not registered in the protocol and should therefore be interpreted as exploratory. The strengths of associations were quantified using odds ratios (OR) and 95% confidence intervals (95% CI). Multivariable models were built if multiple significant predictors were found in the univariable analysis. Analyses were performed with the use of commercial software (IBM SPSS 22.0, SPSS Inc, Chicago, IL). A two-sided *P* value of 0.05 was considered to be statistically significant.

Results

The results of the search for eligible reviews were reported previously in part 1 of this crosssectional study [33] and identified 98 eligible reviews. The PRISMA flow diagram and all included reviews and all excluded studies with rationale were given again in Additional file 3. Figures 1 and 2 present the flow diagrams of the answers to the research questions, and Tables 4 and 5 report the pertinent proportion statistics. The number of identified systematic reviews and the number of eligible systematic reviews of orthodontic interventions given in these tables were published previously [33] and were reported again to give context to the outcomes to our research questions. In these tables, outcomes are further subdivided for eligible reviews that did (n = 84) or did not (n = 14) seek any findings related to adverse effects of interventions in the included studies. The initial inter-operator agreement between both operators for assigning spin was high (Cohen's $\kappa = 0.94$), and complete agreement between operators was reached after discussion.




Fig. 1 Reporting or considering adverse effects of orthodontic interventions in abstracts of systematic reviews




Fig. 2 Spin on adverse effects of orthodontic interventions in abstracts of systematic reviews

Results for questions 1a and 1b

The results for questions 1a and 1b combined showed that the majority 76.5% (75/98) of eligible reviews reported or considered (i.e., discussed, weighted etc.) potential adverse effects of orthodontic interventions in the abstract (Fig. 1). This prevalence was much higher in the reviews that sought any findings related to adverse effects of interventions in the included studies (88.1% (74/84) than those reviews that did not seek these findings (7.1% (1/14) (Fig. 1).

Results for questions 2a and 2b

The results for questions 2a and 2b combined showed that the total proportion of the presence of spin on adverse effects in the abstract was 40.8% (40/98) in the eligible reviews (Table 4). This prevalence was considerable higher in the reviews that sought any findings related to adverse effects of interventions in the included studies (Question 2a), i.e., 46.4% (39/84) than those reviews that did not seek such findings 7.1% (1/14) (Fig.2).

Results for questions 3a and 3b

For questions 3a and 3b combined, misleading reporting was the predominant type of spin i.e., 90% (36/40), which was subdivided in the categories of not reporting, 32.5% (13/40), and selective reporting 57.5% (23/40) (Table 5). Misleading interpretation and misleading (inappropriate) extrapolation types of spin were respectively 7.5% (3/40) and 2.5% (1/40).

Table 4 Outcomes on	reporting	or considering	adverse	effects	and	spin	in	abstracts	of	systematic	reviews (of	orthodontic
interventions													

Interventions	P. A.	F10					48	
Description of outcomes	Cochrane	EJO	AJODO	AO	KJO	O&CR	All orthodontic journals	All journals
The number of identified system- atic reviews	n=29	n=100	n=68	n=61	n=11	n=53	n=293	n=322
The number of eligible systematic reviews of ortho- dontic interven- tions	n=10	n=28	n=20	n=23	n=3	n=14	n=88	n=98
The prevalence of eligible systematic reviews of ortho- dontic interven- tions that sought any findings related to adverse effects of interventions in the included studies ⁸	100.0% (10/10)	92.9% (26/28)	75.0% (15/20)	78.3% (18/23)	100.0% (3/3)	85.7% (12/14)	84.1% (74/88)	85.7% (84/98)
Outcome 1: The prevalence of eligible systematic reviews in which potential adverse effects of interven- tions were reported or considered (<i>u.e.</i> , discussed, weighted etc.) in the abstract [®]	100.0% (10/10)	82.1% (23/28)	65.0% (13/20)	65.2% (15/23)	100.0% (3/3)	78.6% (11/14)	73.9% (65/88)	76.5% (75/98)
Outcome 1a: The prevalence of eligible systematic reviews in which potential adverse effects of interven- tions were reported or considered (i.e., discussed, weighted ecc) in the abstract [®]	100% (10/10)	88.5% (23/26)	86.7% (13/15)	77.8% (14/18)	100% (3/3)	91.7% (11/12)	865% (64/74)	88.1% (74/84)
Outcome 1b: The prevalence of eligible systematic reviews in which potential adverse effects of interven- tions were reported or considered (i.e., discussed, weighted etc.) in the abstract*	00.0% (0/0)	0.0% (0/2)	0.0% (0/5)	20.0% (1/5)	0.0% (0/0)	0.0% (0/2)	7.1%(1/14)	7,1% (1/14)
Outcome 2: The prevalence of eligible systematic reviews in which spin was identified on adverse effects of orthodontic interventions in the abstract*	60,0% (6/10)	35.7% (10/28)	30.0% (6/20)	39.1% (9/23)	66.7% (2/3)	50.0% (7714)	38.6% (34/88)	40,8% 40/98

Table 4 (continued)

Description of outcomes	Cochrane	EIO	AJODO	AO	KJO	O&CR	All orthodontic journals	All journals
Outcome 2a: The prevalence of eligible systematic reviews in which spin was identified on adverse effects of orthodontic interventions in the abstract ⁶	60.0% (6/10)	38.5% (10/26)	40.0% (6/15)	44,4% (8/18)	66.7% (2/3)	58.3% (7/12)	44.6% (33/74)	46,4% (39/84)
Outcome 2b: The prevalence of eligible systematic reviews in which spin was identified on adverse effects of orthodontic interventions in the abstract ⁴	0.0% (0/0)	0.0% (0/2)	0.0% (0/5)	20.0% (1/5)	0.0% (0/0)	0.0% (0/2)	7,1% (1/14)	7.1% (1/14)

*Outcome 1a addresses research question 1a, outcome 1b addresses research question 1b, and outcome 1 addresses the answers to questions 1a and 2b combined, outcome 2a addresses research question 2a, outcome 2b addresses research question 2b, and outcome 2 addresses the answers to questions 2a and 2b combined (see Figs. 1 and 2)

Table 5 Outcome 3: types and categories of spin in abstracts of systematic reviews of orthodontic interventions

Type of spin and category	Cochrane	E)O	AJODO	AO	KJO	O&CR	All orthodontic journals	All journals
Misleading reporting Category: Not reporting in the abstract on the results of adverse effects found in the review.	n=2	n=3	n=2	n=5	n=0	<i>n</i> =1	n = 11 Prevalence: 32.4% (11/34)	n = 13 Prevalence: 32.5% (13/40)
Misleading reporting Category: Selective reporting in the abstract on the results of adverse effects found in the review	n=4	n=6	n=4	n=2	n=2	n=5	n = 19 Prevalence: 55.9% (19/34)	n=23 Prevalence: 57.5% (23/40)
Misleading interpretation Category: Claiming in the abstract that the intervention is safe (has no or minimal adverse effects), despite concerning results on adverse effects found in the review		n=1		n=1		n=1	n=3 Prevalence: 8.8% (3/34)	n=3 Prevalence: 7.5% (3/40)
Misleading (inappropriate) extrapolation Category: Results are extrapolated in the abstract to another population, intervention, outcome or setting than were assessed in the review despite evidence on adverse effects on a different population, intervention, outcome or setting.				<i>n</i> =1			n=1 Prevalence: 2.9% (1/34)	n=1 Prevalence: 2.5% (1/40)

Explorative analyses

The findings of our explorative analysis on the presence of spin on adverse effects in abstracts of systematic reviews of orthodontic interventions were reported in Table 6. Compared to the Cochrane Database of Systematic Reviews, the EJO (OR: 0.37, 95% CI: 0.08 to 1.63), the AJODO (OR: 0.29, 95% CI: 0.06 to 1.39), the AO (OR: 0.43, 95% CI: 0.09 to 1.95), the KJO (OR: 1.33, 95% CI: 0.09 to 20.11), and the O&C (OR: 0.67, 95% CI: 0.13 to 3.45) had similar odds of the presence of spin on adverse effects in abstracts of systematic reviews of orthodontic interventions. The odds of the presence of spin on adverse effects in abstracts of systematic reviews of orthodontic interventions did not change over the sampled years (OR: 1.03, 95% CI: 0.9 to 1.16). The odds of the presence of spin on adverse effects in abstracts of systematic reviews of orthodontic interventions did not change depending on

the number of authors (OR: 0.93, 95% CI: 0.71 to 1.21). Compared to systematic reviews that did not report conflicts of interest, systematic reviews that reported conflicts of interest had similar odds (OR: 0.74, 95% CI: 0.32 to 1.68) of the presence of spin on adverse effects in abstracts of systematic reviews of orthodontic interventions. Systematic reviews on type 1 orthodontic interventions, i.e., orthodontic interventions to move teeth or change the jaw size or position for orthodontic purposes compared with systematic reviews on type 3 orthodontic interventions, i.e., orthodontic interventions to maintain or stabilize orthodontic results had similar odds (OR: 1.1, 95% CI: 0.45 to 2.67) of the presence of spin on adverse effects of orthodontic interventions with additional surgical, pharmacological, or vibratory interventions was not included in the analysis since there was only 1 systematic review of this type of intervention.

Item	Variable insertion in the model	Description	Yes (%)	No (%)	OR	Lower 95%CI	Upper 95%Cl	P value
Journal	Categorical	Cochrane	4 (40:0%)	6 (60.0%)	1	40)	4	10
		EJO	18 (64.3%)	10 (35.7%)	0.37	0.08	1.63	0.19
		AJODO	14 (70.0%)	6 (30.0%)	0.29	0.06	1.39	0.12
		AO	14 (60.9%)	9 (39.1%)	0.43	0.09	1.95	0.27
		KJO	1 (33.3%)	2 (66.7%)	1.33	0.09	20.11	0.84
		OBC	7 (50.0%)	7 (50.0%)	0.67	0.13	3.45	0.63
Year of publication	Continuous				1,03	0.9	1.16	0.7
		2009	1 (100.0%)	0 (0.0%)				
		2010	1 (33.3%)	2 (66.7%)				
		2011	5 (71.4%)	2 (28.6%)				
		2012	2 (100.0%)	0 (0.0%)				
		2013	6 (66.7%)	3 (33.3%)				
		2014	3 (33.3%)	6 (66.7%)				
		2015	9 (75.0%)	3 (25.0%)				
		2016	5 (55.6%)	4 (44,4%)				
		2017	5 (55.6%)	4 (44.4%)				
		2018	6 (46.2%)	7 (53.8%)				
		2019	3 (75.0%)	1 (25.0%)				
		2020	7 (58.3%)	5 (41.7%)				
		2021	5 (62.5%)	3 (37.5%)				
Number of authors	Continuous				0.93	0.71	1.21	0.59
		2	3 (50.0%)	3 (50.0%)				
		3	7 (50.0%)	7 (50.0%)				
		4	16 (69.6%)	7 (30.4%)				
		5	15 (55.6%)	12 (44,4%)				
		6	8 (53.3%)	7 (46,7%)				
		7	6 (66.7%)	3 (33.3%)				
		8	2 (100.0%)	0 (0.0%)				
		9	1 (50.0%)	1 (\$0.0%)				
Conflict of interest reported	Categorical	Yes	32 (56.1%)	25 (43.9%)	0.74	0.32	1.68	0.47
	M.2050024	No	26 (63.4%)	15 (36.6%)	1		Contraction of the second	
Conflict of interest present		Not reported	26 (63.4%)	15 (36.6%)	NA	NA.	NA	NA
		No	32 (56.1%)	25 (43.9%)	NA.	NA	NA.	NA
Funding reported	Categorical	Yes	23 (56.1%)	18 (43.9%)	0.8	0.35	1.81	0.6
	5412264466	No	35 (61.4%)	22 (38.6%)	1	1000	-	-
Type of orthodontic intervention*	Categorical	1	41 (59.4%)	28 (40.6%)	1	-	14	*
1922	51	2	1 (100.0%)	0 (0.0%)	NA	NA	NA	NA
		3	16 (57.1%)	12 (42.9%)	1,1	0.45	2.67	0.84

Table 6 Associations between presence of spin in the abstract and characteristics of the systematic review

^{*}Type 1 orthodontic interventions: Orthodontic interventions to move teeth or change the jaw size or position for orthodontic purposes. Type 2 orthodontic interventions: Orthodontic interventions with additional surgical, pharmacological, or vibratory interventions. Type 3 orthodontic interventions: Orthodontic interventions to maintain or stabilize orthodontic results

Discussion

Principal findings of the study

This cross-sectional study showed that the majority, i.e., 76.5% (75/98), of the eligible systematic reviews reported or considered (i.e., discussed, weighted) potential adverse effects of orthodontic interventions in the abstract (Table 4). In 40.8% (40/98), spin on adverse effects was found in the abstract of these reviews (Table 4). Spin related to misleading reporting was the predominant, i.e., 90.0% (36/40), type of spin (Table 5). No association was found between the presence of spin in the abstract and any of the predictors (Table 6).

Comparison with other studies

Item 1 of the 2004 CONSORT (Consolidated Standards of Reporting Trials) Harms extension [42] states that "if the study collected data on harms and benefits, the title or abstract should so state". In this second part of our cross-sectional studies, 76.5% (75/98) of systematic reviews reported or considered adverse effects of interventions in the abstracts. This was lower than the 84.7% (83/98) of reviews that sought and reported findings on adverse effects in the main text or supplementary files of these reviews as we reported in part 1 [33]. A recent overview of systematic reviews by Jungueira et al. [43] found that harms were reported in 47% (258/552) of the abstracts of RCTs published prior to the CONSORT harms statement and in 54% (643/1201) of the abstracts of the RCTs published after the publication of this statement, indicating only a limited improvement in recent years. Qureshi et al. [14] found that most systematic reviews 81.4% (57/70) on interventions with gabapentin reported a statement on harms in the abstract. Different results in reporting of adverse effects in abstracts in this cross-sectional study compared with those in other studies could be the result of variables such as differences in (1) research design, i.e., systematic reviews versus RCTs, (2) sample size, (3) what is reported on harms in the abstract, e.g., specific versus more general statements, and (4) the field of research. For example, the relatively high prevalence of reporting of adverse effects in abstracts of systematic reviews in this study could be the result of having included only reviews of orthodontic interventions. Orthodontists might be more attentive in assessing adverse effects, because assessing adverse events such as undesired outcomes of orthodontic treatment and relapse is part of daily clinical practice.

A wide variety of prevalence proportions on spin has been identified in abstracts of systematic reviews of randomized-and non-randomized studies [23, 25, 28–30]. These studies identified proportions of spin that varied between 23% (24/105) of spin in abstracts of RCTs in rheumatology [28] and 84% (107/128) of spin in abstracts of non-randomized studies that assessed interventions [23]. According to our scoping searches, spin in the field of orthodontics has been assessed only in 2 recent studies [31, 32]. Spin was identified in 62.2% (69/111) of abstracts of parallel-group RCTs with clearly stated statistically non-significant primary outcomes [31] and in 48.6% (53/109) of abstracts of orthodontic meta-analyses [32]. In our study, none of the predictors assessed was associated with the presence of spin in this study. Similar findings were identified by Guo et al. [31] on the overlapping predictors with our study, i.e., "the year of publication" and "the number of 96

authors." Makou et al. [32] also found no association with the presence of spin for the overlapping predictors "journal" and "year of publication" but found a higher risk of spin in studies with a large number of authors (\geq 6). However, direct comparisons of our results on spin and those identified in other studies are often difficult because of differences in variables such as (1) the types and subtypes of spin and definitions of spin, (2) the research design, (3) the locations in the text where spin was assessed, (4) the field of research, (5) the types of interventions, (6) the journals included, and (7) the time point of the publication [42].

Strengths and limitations

This study has the following strengths: (1) the research methods were pilot tested on a series of systematic reviews and RCTs to consistently extract data and to calibrate data extractors; (2) the protocol of this study was published a priori; and (3) according to our scoping searches, this is the first study that assessed spin on adverse effects in abstracts of systematic reviews of interventions. These searches also showed that our protocol was the first article [24] that planned to assess spin in the field of orthodontics. Subsequently, 2 additional studies [31, 32] have assessed other types of spin in the orthodontic literature, indicating a growing interest in this topic. (4) All raw data were either included with this manuscript in additional files or registered in Open Science Framework (https://osf.io/ka7mp/).

This study also has limitations such as (1) the risk of inaccurate findings on reporting adverse effects of interventions in abstracts as a result of the inconsistent assessment and reporting of adverse effects in both primary research and in systematic reviews and (2) the assessment of spin is not completely objective [44]. However, our inter-operator agreement was high as indicated by a high Cohen's κ (0.94), and disagreements were completely resolved through discussions. (3) The wide variety of different types and definitions of spin and the assessment of spin in different contexts often limits comparing findings on spin between studies [44]. (4) This study assessed a variety of proportions exclusively in Cochrane intervention reviews and in the 5 orthodontic journals with the highest impact factor. Our findings therefore probably underestimate the true magnitude of proportions on poor reporting and spin on adverse effects in the abstracts of the wider body of orthodontic systematic reviews, and (5) the true magnitude of some proportions could also be underestimated, because we assessed a recent sample (August 1, 2009, until July 31, 2021) of reviews. In this context, one should consider that poor reporting has decreased over time as was shown in a study that assessed the evolution of poor reporting in 20,920 RCTs included in a sample of Cochrane reviews [45].

Implications and future research

Our results imply that end-users of systematic reviews of orthodontic interventions have to be careful when interpreting the findings on adverse effects in abstracts of both Cochrane reviews and those published in the 5 leading orthodontic journals. This is particularly important, because the title and abstracts are often the only read sections of biomedical papers [6] and spin in abstracts can bias the clinician's interpretations of the results [26]. Reading the full text of research studies is not a solution, because recent studies showed

that the proportions of spin in abstracts are similar to those in the full text of the pertinent RCTs [29] and systematic reviews [46]. Guideline developers, researchers, peer reviewers, and editors have an important role in tackling poor reporting and spin regarding adverse effects in abstracts of systematic reviews of orthodontic interventions. Standards for reporting adverse effects in abstracts of systematic reviews of systematic reviews of interventions have to be developed. Much research is ahead.

Abbreviations

AJODO American Journal of Orthodontics and Dentofacial Orthopedics AO Angle Orthodontist EJO European Journal of Orthodontics KJO Korean Journal of Orthodontics O&CR Orthodontics and Craniofacial Research MECIR Methodological Expectations of Cochrane Intervention Reviews PRISMA Preferred reporting items for systematic reviews and meta-analyses RCT Randomized controlled trial STROBE Strengthening the Reporting of Observational Studies in Epidemiology

Supplementary Information

The online version contains supplementary material available at https://doi. org/10.1186/s13643-023-02269-3.

Additional file 1. STROBE Checklist: Checklist of items that should be included in reports of cross-sectional studies [35].

Additional file 2. 2A. Differences between the protocol and the completed cross-sectional study. 2B. Search terms and their derivatives. 2C. Data collection forms. 2D. Adverse effects hypothetically linked to orthodontic interventions.

Additional file 3. 3A. PRISMA flow diagram. 3B. Included reviews. 3C. Excluded studies with rationale

Acknowledgements

Not applicable.

Authors' contributions

This cross-sectional study was conceived and designed by PS and RMR. These 2 researchers also conducted all data extraction procedures. PS, RMR, and NDG conducted the data analyses and statistical analyses. NDG provided sup- port on methodological issues. All 3 authors revised the various draft versions of this paper and read and approved the final submitted manuscript. RMR is the guarantor.

Funding

All expenses for developing and conducting of this cross-sectional study were paid evenly by each author.

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate Not applicable.

Consent for publication Not applicable.

Competing interests

Reint Meursinge Reynders is an associate editor for *Systematic Reviews*. All three authors declare that they have no further competing interests.

Author details

¹Department of Orthodontics, Academisch Centrum Tandheelkunde Amsterdam (ACTA), University of Amsterdam, Gustav Mahlerlaan 3004, Amsterdam 1081 LA, The Netherlands.

²Department of Clinical Sciences, College

of Veterinary Medicine, Cornell University, 930 Campus Rd, Ithaca, NY 14853, USA.

³Department of Oral and Maxillofacial Surgery, Academic Medical Center, University of

Amsterdam, Meibergdreef 9, Amsterdam 1105 AZ, The Nether- lands. ⁴Studio Di Ortodonzia, Via Matteo Bandello 15, Milan 20123, Italy.

Received: 11 February 2022 Accepted: 8 June 2023

Published online: 20 June 2023

References

- Beller EM, Glasziou PP, Altman DG, Hopewell S, Bastian H, Chalmers I, Gøtzsche PC, Lasserson T, Tovey D, PRISMA for Abstracts Group. PRISMA for Abstracts: reporting systematic reviews in journal and conference abstracts. PLoS Med. 2013;10(4):e1001419. https://doi.org/10.1371/ journal.pmed.1001419. Epub 2013 Apr 9. PMID: 23585737; PMCID: PMC3621753.
- Haneef R, Lazarus C, Ravaud P, Yavchitz A, Boutron I. Interpretation of results of studies evaluating an intervention highlighted in Google health news: a cross-sectional study of news. PLoS One. 2015;10(10):e0140889. https://doi.org/10.1371/journal.pone.0140889. eCollection 2015.
- Haneef R, Yavchitz A, Ravaud P, Baron G, Oranksy I, Schwitzer G, Boutron I. Interpretation of health news items reported with or without spin: protocol for a prospective meta-analysis of 16 randomised controlled trials. BMJ Open. 2017;7(11):e017425. https://doi.org/10.1136/bmjop en-2017-017425.
- 4. Ad Hoc Working Group for Critical Appraisal of the Medical Literature. A proposal for more informative abstracts of clinical articles. Ann Intern Med. 1987;106:598–604.
- Haynes RB, Mulrow CD, Huth EJ, Altman DG, Gardner MJ. More informative abstracts revisited. Ann Intern Med. 1990;113:69–76.
- IslamajDogan R, Murray GC, Névéol A, Lu Z. Understanding Pub-Med user search behavior through log analysis. Database (Oxford). 2009;2009:bap018. https://doi.org/10.1093/database/bap018. Epub 2009 Nov 27. PMID: 20157491; PMCID: PMC2797455.
- Page MJ, Cumpston M, Chandler J, Lasserson T. Chapter III: Reporting the review. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. Cochrane handbook for systematic reviews of interventions version 6.2 (updated February 2021). Cochrane; 2021. Available from www.training.cochrane.org/handbook.
- Higgins JPT, Lasserson T, Chandler J, Tovey D, Thomas J, Flemyng E, Churchill R. Methodological expectations of Cochrane intervention reviews. London: Cochrane; 2021. Available from: MECIR Version February 2021 clean final_1.pdf (cochrane.org). Accessed 4 Dec 2021.
- Bagul NB, Kirkham JJ. The reporting of harms in randomized controlled trials of hypertension using the CONSORT criteria for harm reporting. Clin Exp Hypertens. 2012;34(8):548–54. https://doi.org/10.3109/10641963. 2012.681724. Epub 2012 May 9.
- Carmichael K, Nolan SJ, Weston J, Tudur Smith C, Marson AG. Assessment of the quality of harms reporting in non-randomised studies and randomised controlled studies of topiramate for the treatment of epilepsy using CONSORT criteria. Epilepsy Res. 2015;114:106–13. https://doi.org/ 10.1016/j.eplepsyres.2015.04.019. Epub 2015 May 8. PMID: 26088893.
- Hodkinson A, Kirkham JJ, Tudur-Smith C, Gamble C. Reporting of harms data in RCTs: a systematic review of empirical assessments against the CONSORT harms extension. BMJ Open. 2013;3(9):e003436. https://doi. org/10.1136/bmjopen-2013-003436.
- 12. Parsons R, Golder S, Watt I. More than one-third of systematic reviews did not fully report the adverse events outcome. J Clin Epidemiol. 2019;108:95–101. https://doi.org/10.1016/j.jclinepi.2018.12.007.
- 13. Pitrou I, Boutron I, Ahmad N, Ravaud P. Reporting of safety results in published reports of randomized controlled trials. Arch Intern Med. 2009;169(19):1756–61.
- Qureshi R, Mayo-Wilson E, Rittiphairoj T, McAdams-DeMarco M, Guallar E, Li T. Harms in Systematic Reviews Paper 2: methods used to assess harms are neglected in systematic reviews of gabapentin. J Clin Epidemiol. 2022;143:212–23. https://doi.org/10.1016/j.jclinepi.2021.10.024. Epub 2021 Nov 3. PMID: 34742789; PMCID: PMC9875742.
- Saini P, Loke YK, Gamble C, Altman DG, Williamson PR, Kirkham JJ. Selective reporting bias of harm outcomes within studies: findings from a cohort of systematic reviews. BMJ. 2014;349:g6501. https://doi.org/10. 1136/bmj.g6501. PMID: 25416499; PMCID: PMC4240443.
- Zorzela L, Golder S, Liu Y, Pilkington K, Hartling L, Joffe A, Loke Y, Vohra S. Quality of reporting in systematic reviews of adverse events: systematic review. BMJ. 2014;348:f7668. https://doi.org/10.1136/bmj.f7668. PMID: 24401468; PMCID: PMC3898583.
- 17. Cochrane library. About Cochrane reviews. Available from https://www. cochranelibrary.com/about/about-cochranereviews. Accessed 8 June 2023.

- Steegmans PAJ, Bipat S, MeursingeReynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross- sectional study. Syst Rev. 2019;8(1):89. https://doi.org/10.1186/s13643-019-1000-1. PMID: 30953538; PMCID: PMC6449933.
- 19. Glossary of terms in the Cochrane Collaboration. Version 4.2.5. Updated May 2005. Available from: http://aaz.hr/resources/pages/57/7.%20Coc hrane%20glossary.pdf. Accessed 4 Dec 2021.
- Peryer G, Golder S, Junqueira D, Vohra S, Loke YK. Chapter 19: Adverse effects. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. Cochrane handbook for systematic reviews of interventions version 6.2 (updated February 2021). Cochrane; 2021. Available from www.training.cochrane.org/handbook.
- 21. Boutron I, Ravaud P. Misrepresentation and distortion of research in bio- medical literature. Proc Natl Acad Sci U S A. 2018;115(11):2613–9. https:// doi.org/10.1073/pnas.1710755115.
- Boutron I. Spin in scientific publications: a frequent detrimental research practice. Ann Emerg Med. 2020;75(3):432–4. https://doi.org/10.1016/j. annemergmed.2019.11.002. Epub 2019 Dec 23. PMID: 31874770.
- Lazarus C, Haneef R, Ravaud P, Boutron I. Classification and prevalence of spin in abstracts of nonrandomized studies evaluating an intervention. BMC Med Res Methodol. 2015;13(15):85. https://doi.org/10.1186/ s12874-015-0079-x.
- 24. Steegmans PAJ, Di Girolamo N, MeursingeReynders RA. Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews. Res Integr Peer Rev. 2019;4:27. https://doi.org/10.1186/ s41073-019-0084-4.
- Boutron I, Dutton S, Ravaud P, Altman DG. Reporting and interpretation of randomized controlled trials with statistically nonsignificant results for primary outcomes. JAMA. 2010;303(20):2058–64. https://doi.org/10.1001/ jama.2010.651.
- Boutron I, Altman DG, Hopewell S, Vera-Badillo F, Tannock I, Ravaud P. Impact of spin in the abstracts of articles reporting results of randomized controlled trials in the field of cancer: the SPIIN randomized controlled trial. J Clin Oncol. 2014;32(36):4120–6. https://doi.org/10.1200/JCO.2014. 56.7503. Epub 2014 Nov 17. PMID: 25403215.
- Yavchitz A, Ravaud P, Altman DG, Moher D, Hrobjartsson A, Lasserson T, Boutron I. A new classification of spin in systematic reviews and meta- analyses was developed and ranked according to the severity. J Clin Epidemiol. 2016;75:56–65. https://doi.org/10.1016/j.jclinepi.2016.01.020. Epub 2016 Feb 2.
- Mathieu S, Giraudeau B, Soubrier M, Ravaud P. Misleading abstract conclusions in randomized controlled trials in rheumatology: comparison of the abstract conclusions and the results section. Joint Bone Spine. 2012;79(3):262–7. https://doi.org/10.1016/j.jbspin.2011.05.008. Epub 2011 Jul 5. PMID: 21733728.
- Khan MS, Lateef N, Siddiqi TJ, Rehman KA, Alnaimat S, Khan SU, Riaz H, Murad MH, Mandrola J, Doukky R, Krasuski RA. Level and prevalence of spin in published cardiovascular randomized clinical trial reports with statistically nonsignificant primary outcomes: a systematic review. JAMA Netw Open. 2019;2(5):e192622. https://doi.org/10.1001/jamanetwor kopen.2019.2622. PMID: 31050775; PMCID: PMC6503494.
- Jones C, Rulon Z, Arthur W, Ottwell R, Checketts J, Detweiler B, Calder M, Adil A, Hartwell M, Wright DN, Vassar M. Evaluation of spin in the abstracts of systematic reviews and meta-analyses related to the treatment of proximal humerus fractures. J Shoulder Elbow Surg. 2021:S1058- 2746(21)00016-1. https://doi.org/10.1016/j.jse.2020.11.026. Epub ahead of print. PMID: 33482369.
- Guo F, Fang X, Li C, Qin D, Hua F, He H. The presence and characteristics of 'spin' among randomized controlled trial abstracts in orthodontics. Eur J Orthod. 2021;43(5):576–82. https://doi.org/10.1093/ejo/cjab044. PMID: 34397084.
- Makou O, Eliades T, Koletsi D. Reporting, interpretation, and extrapolation issues (SPIN) in abstracts of orthodontic meta-analyses published from 2000 to 2020. Eur J Orthod. 2021;43(5):567–75. https://doi.org/10.1093/ ejo/cjab009. Epub ahead of print. PMID: 33740054.
- Steegmans PAJ, Di Girolamo N, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1). Accepted for publication in Systematic Reviews on 16 June 2023.
- Yavchitz A, Boutron I, Bafeta A, Marroun I, Charles P, Mantz J, Ravaud P. Misrepresentation of randomized controlled trials in press releases and news coverage: a cohort study. PLoS Med.

2012;9(9):e1001308. https:// doi.org/10.1371/journal.pmed.1001308. Epub 2012 Sep 11. PMID: 22984354; PMCID: PMC3439420.

- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Ann Intern Med. 2007;147(8):573–7.
- 36. Lefebvre C, Glanville J, Briscoe S, Littlewood A, Marshall C, Metzendorf M-I, Noel-Storr A, Rader T, Shokraneh F, Thomas J, Wieland LS. Chapter 4: Searching for and selecting studies. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. Cochrane handbook for systematic reviews of interventions version 6.2 (updated February 2021). Cochrane; 2021. Available from www.training.cochrane.org/handbook.
- The Cochrane Database of Systematic Reviews (CDSR). Available from: https://www.cochranelibrary.com/cdsr/about-cdsr. Accessed 4 Dec 2021.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. J Clin Epidemiol. 2009;62(10):e1-34. https://doi.org/10.1016/j.jclinepi.2009.06.006. Epub 2009 Jul 23. PMID: 19631507.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. J Clin Epidemiol. 2009;62(10):1006–12. https://doi.org/10.1016/j.jclinepi.2009.06.005. Epub 2009 Jul 23. PMID: 19631508.
- Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F, editor. Orthodontics-basic aspects and clinical considerations. 2012. Under CC BY 3.0 license. www.intechopen.com. Available from: https://cdn.intechopen. com/pdfs/31388/InTech-Risks_and_complications_associated_with_ orthodontic_treatment.pdf. Accessed 4 Dec 2021.
- 41. Sergeant ESG. Sergeant, ESG, 2018. Epitools Epidemiological Calculators. Ausvet. Available from: http://epitools.ausvet.com.au. Accessed 4 Dec 2021.
- Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, Schulz K, Moher D, CONSORT Group. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med. 2004;141(10):781–8.
- Junqueira DR, Phillips R, Zorzela L, Golder S, Loke Y, Moher D, Ioannidis JPA, Vohra S. Time to improve the reporting of harms in randomized con- trolled trials. J Clin Epidemiol. 2021;136:216–20. https://doi.org/10.1016/j. jclinepi.2021.04.020. Epub 2021 May 10. PMID: 33984494.
- Chiu K, Grundy Q, Bero L. 'Spin' in published biomedical literature: a methodological systematic review. PLoS Biol. 2017;15(9):e2002173. https://doi.org/10.1371/journal.pbio.2002173. PMID: 28892482; PMCID: PMC5593172.
- 45. Dechartres A, Trinquart L, Atal I, Moher D, Dickersin K, Boutron I, Perrodeau E, Altman DG, Ravaud P. Evolution of poor reporting and inadequate methods over time in 20 920 randomised controlled trials included in Cochrane reviews: research on research study. BMJ. 2017;357:j2490. https://doi.org/10.1136/bmj.j2490. Erratum in: BMJ. 2017 Aug 8:358:j3806. PMID: 28596181.
- 46. Eleftheriadi I, Ioannou T, Pandis N. Extent and prevalence of spin in randomized controlled trials in dentistry. J Dent. 2020;100:103433. https://doi. org/10.1016/j.jdent.2020.103433. Epub 2020 Jul 17. PMID: 32687962.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



CHAPTER 6

Fixed orthodontic retainers and periodontal health: a critical appraisal of a systematic review

This is a non-edited version of the manuscript:

Steegmans PAJ, Meursinge Reynders RA. Fixed orthodontic retainers and periodontal health. Evid Based Dent. 2020 Dec;21(4):146-149.

A commentary on

Arn M L, Dritsas K, Pandis N, Kloukos D. The effects of fixed orthodontic retainers on periodontal health: A systematic review. Am J Orthod Dentofacial Orthop 2020; DOI: 10.1016/j.ajodo.2019.10.010.

Abstract

Data sources: The following electronic databases were searched from 1946 to 31 August 2019: Medline, Embase, the Cochrane Oral Health Group's Trials Register, CENTRAL, ClinicalTrials.gov, the National Research Register and Pro-Quest Dissertation Abstracts and Thesis database. **Study selection** The following study designs were eligible: randomised controlled trials (RCTs), controlled clinical trials, cohort studies of prospective and retrospective design, and cross-sectional studies that reported periodontal measurements on patients who received fixed retention after orthodontic therapy. Studies irrespective of their language were selected by two reviewers independently.

Data extraction and synthesis: Data extraction from the selected studies and risk of bias assessments were performed by two reviewers independently. Specific risk of bias tools were used according to the pertinent research designs of the included studies. Criteria for conducting a meta-analysis were not met and a qualitative synthesis was conducted.

Results: Twenty-nine studies fulfilled the eligibility criteria; that is, 11 RCTs, four prospective cohort studies, one retrospective cohort study and 13 cross-sectional studies. The quality of the evidence was low for most of the studies included in this review. Contrary to the general consensus, two RCTs, one prospective cohort study and two cross-sectional studies identified poorer periodontal health in patients with fixed orthodontic retainers.

Conclusions: The authors of this systematic review concluded that fixed orthodontic retainers in the majority of the 29 included studies seemed to be a method of retention that is rather compatible with periodontal health, or at least not related to severe detrimental consequences for the periodontium. No recommendations on the best type of fixed retainer to use could be given. High-quality evidence from long-term studies is necessary to provide definitive conclusions on the relationship between fixed retainers and periodontal health.

Practice Point

- This review concluded that fixed orthodontic retainers in the majority of the 29 included studies seemed rather compatible with periodontal health or are at least not related to severe detrimental outcomes of the periodontium. No recommendation on the best type of fixed retainer to use in clinical practice could be given.
- The findings of this review should be considered in the context that: (1) five of the included studies reported poorer periodontal health around fixed retainers; (2) most included studies were of low quality; and (3) a variety of additional limitations identified in this critical appraisal could have skewed these results.
- Evidence from high-quality long-term studies is necessary to provide definitive conclusions on the relationship between fixed orthodontic retainers and periodontal health.

Commentary

Objectives and key findings of the systematic review

Arn et al.¹ defined the following objectives for their systematic review: 1) 'to evaluate the potentially deleterious effects of fixed orthodontic retainers on periodontal health'; and 2) 'to compare different kinds of fixed retainers according to their effects on periodontal health, and if possible, to recommend [one] of them'. The majority of the 29 included studies found that orthodontic fixed retainers seemed rather compatible with periodontal health, but five studies reported poorer periodontal status in the presence of these retainers. The quality of evidence for most of these studies was low. No recommendations on the best type of fixed retainer to use in practice could be given.

Methods of our critical appraisal of the systematic review

We applied a three-step critical appraisal strategy for the systematic review. We assessed: 1) how the review was reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist;^{2,3,4} 2) the methodological validity, by applying the AMSTAR 2 tool;⁵ and 3) the risk of bias in the review using the ROBIS tool.^{6,7} These assessments were conducted independently by two reviewers (PS and RMR) and disagreements were resolved through discussions between these operators. We adopted methods of the AMSTAR 2 tool to create an overall quality score.⁵ Seven critical domains (Q1, Q2, Q4, Q8, Q9, Q13, Q14) were assigned for this purpose.

Findings of our critical appraisal

The findings of our critical appraisals are presented in Tables 1, 2 and 3. The overall quality of the review of the AMSTAR 2 assessment was rated as 'moderate' because we identified one unclear critical domain (partial yes) and two non-critical flaws. Table 4 summarises the

limitations identified in Tables 1, 2 and 3, and includes additional shortcomings. Of the ten limitations, two were assigned to the methods and eight to the reporting of the review. The limitations and strengths of this review are further explained below.

Item		Reporting score
Title		
Title	1	Reported
Abstract		
Structured summary	2	Reported
Introduction		
Rationale	3	Reported
Objectives	4	Partially reported
Methods		
Protocol and registration	5	Reported
Eligibility criteria	6	Reported
Information sources	7	Reported
Search	8	Reported
Study selection	9	Reported
Data collection process	10	Reported
Data items	11	Reported
Risk of bias in individual studies	12	Reported
Summary measures	13	Reported
Synthesis of results	14	Reported
Risk of bias across studies	15	Reported
Additional analyses	16	Not applicable
Results		
Study selection	17	Partially reported
Study characteristics	18	Reported
Risk of bias within studies	19	Reported
Results of individual studies	20	Reported
Synthesis of results	21	Reported
Risk of bias across studies	22	Reported
Additional analysis	23	Not applicable
Discussion		
Summary of evidence	24	Reported
Limitations	25	Reported
Conclusions	26	Reported
Funding		
Funding	27	Not reported

AMSTAR questions	Scores
Q1. Did the research questions and inclusion criteria for the review include the components of PICO?	Yes
Q2. Did the report of the review contain an explicit statement that the review methods were established before the conduct of the review and did the report usify any significant deviations from the protocol?	Partial yes
Q3. Did the review authors explain their selection of the study designs for inclusion in the review?	Yes
Q4. Did the review authors use a comprehensive literature search strategy?	Yes
Q5. Did the review authors perform study selection in duplicate?	Yes
Q6. Did the review authors perform data extraction in duplicate?	Yes
Q7. Did the review authors provide a list of excluded studies and justify the exclusions?	No
Q8. Did the review authors describe the included studies in adequate detail?	Yes
Q9. For RCTs*	
Did the review authors use a satisfactory technique for assessing the risk of bias (RoB) in individual studies that were included in the review?	Yes
Q9. For NRSI**	
Did the review authors use a satisfactory technique for assessing the risk of bias (RoB) in individual studies that were included in the review?	Yes
Q10. Did the review authors report on the sources of funding for the studies included in the review?	Yes
Q11. For RCTs*	
If meta-analysis was performed did the review authors use appropriate methods for statistical combination of results?	No meta- analysis conducted
Q11. For NRSI**	
If meta-analysis was performed did the review authors use appropriate methods for statistical combination of results?	No meta- analysis conducted
Q12. If meta-analysis was performed, did the review authors assess the potential impact of RoB in individual studies on the results of the meta-analysis or other evidence symthesis?	No meta- analysis conducted
Q13. Did the review authors account for RoB in individual studies when interpreting/discussing the results of the review?	Yes
Q14. Did the review authors provide a satisfactory explanation for, and discussion of, any heterogeneity observed in the results of the review?	Yes
Q15. If they performed quantitative synthesis did the review authors carry out an adequate investigation of publication bias (small study bias) and discuss its likely impact on the results of the review?	No meta- analysis conducted
Q16. Did the review authors report any potential sources of conflict of interest, including any funding they received for conducting the review?	No

Phase 1: Assessing relevance	Phase 2: Identifying concer	ns with the review proce	55		Phase 3: Judging risk of bias
Does the question addressed by the review match the target question?	Domain 1. Study eligibility criteria	Domain 2. Identification and selection of studies	Domain 3. Data collection and study appraisal	Domain 4. Synthesis and findings	Risk of bias in the review
Not applicable, because we did not formulate a target question	0	0	0	0	0

Table 4 Limitations of the methods and the reporting in the review by Arn et al.¹

Limitation #	Item	Limitation
1	Registration or publication of the protocol	No publication or registration of a protocol a priori
2	Defining objectives	The Patients, Interventions, Comparators, Outcomes, Settings (PICOS) elements were not implemented in the objectives
3	Pilot testing of research methods	Not reported whether research methods such as study selection, data extraction, and bias assessments were pilot tested
4	Search strategy	Not reported whether an information specialist for the health sciences was consulted for the development of the search strategies
5	Information sources	Not reported in which information sources the reviewers had found the included studies
6	Excluded studies	A list of excluded studies with justifications for exclusion was not given
7	Contacting of authors	The reviewers reported that they would contact authors on missing or unclear data, but did not report on contacting authors on study eligibility or unclear items for the risk of bias assessment
8	Differences between the protocol and the review	Differences between the protocol and the review were not reported
9	Repository	It was not reported whether a repository (with name) was used to deposit data extraction forms, raw data etc
10	Funding of the systematic review	Funding sources and the role of the funder for this systematic review were not given

Limitations of the methods of the review

Limitations of the methods were: (1) the protocol was not registered, for example, in PROSPERO⁸ or published a priori which could have introduced risk of bias related to selective reporting of outcomes; 9,10,11 (2) the research objectives were defined, but should have been formulated using the Patients, Interventions, Comparators, Outcomes, Settings (PICOS) acronym. Using this format is important for the development of the search strategy¹² and for future reviewers who want to assess the reproducibility of this review or update it.

Limitations of the reporting of the review

The following eight items were not reported in the review: (1) pilot testing of the research methods. These procedures are important to fine-tune these methods a priori and to calibrate reviewers; (2) whether an information specialist for the health sciences was consulted for the development of the various search strategies; (3) the information source in which each included study was found. For example, end users of systematic reviews want to know which included studies were retrieved in the grey literature; (4) the references of excluded studies with justifications; (5) contacting of authors regarding eligibility of studies and unclear risk of bias issues; (6) differences between the review and the protocol; (7) whether a data repository was used for the deposition of for example: data extraction forms with definitions of variables, lists of excluded studies with rationale, raw data on the scoring of risk of bias with rationale, information on selective reporting within studies etc; and (8) the reviewers reported to have no potential conflicts of interest, but the sources of funding and the role of the funder of this review were not reported.

Not reporting on these items could jeopardise the trust in the outcomes of this review and could compromise its reproducibility. However, not reporting on these eight items does not necessarily imply that some of these reporting issues were not implemented. Contacting the reviewers would be the ideal strategy to verify the status on these reporting items.

Strengths of the review

This systematic review has important strengths: (1) it addressed a research question with a wide external validity, because a broad spectrum of orthodontic patients receives fixed orthodontic retainers; (2) mostly high-quality methodology was implemented; (3) an overall low risk of bias of the review process was assigned using the ROBIS tool; (4) up-to-dateness of the search; and (5) a solid assessment of the strengths and limitations of the review.

Limitations of the included studies

Three limitations of the included studies were identified; (1) most of them were of low quality; (2) a high diversity of patients, interventions, comparators, and outcomes complicated data synthesis; and (3) all were published studies. This latter issue could have skewed the findings of this systematic review, because research has shown that adverse effects of interventions are more frequently reported and are more severe in unpublished

versus published studies.^{13,14} The reviewers correctly searched various sources of grey literature, but did not identify non-published research. Searching a broader spectrum of grey literature for reviews that assess the adverse effects of interventions could be indicated.

Can we implement the findings of this review in daily practice?

The reviewers concluded that the overall consensus of the included studies was that orthodontic fixed retainers seemed rather compatible with periodontal health. However, before implementing these findings in daily practice clinicians should: (1) consider that five of the 29 included studies reported contrary findings; (2) consider that most studies provided low-quality evidence; (3) assess whether the strengths of the review outweigh its limitations; (4) consider all patient-important outcomes regarding retention interventions; (5) weigh the costs of the intervention; and (6) assess whether the patients in the included studies differ from those in front of them.

Conclusions

This review addressed research questions on the adverse effects of fixed orthodontic retainers that are important for clinicians, researchers, policy makers, and a broad spectrum of orthodontic patients. The reviewers concluded that fixed orthodontic retainers in the majority of included studies seemed rather compatible with periodontal health or are at least not related to severe detrimental outcomes for the periodontium. No recommendation could be given on the best type of fixed retainer to use in clinical practice. The findings of this review should be considered in the context that: (1) five of the 29 included studies reported poorer periodontal health around fixed retainers; (2) most included studies were of low quality; and (3) various additional limitations identified in this critical appraisal could have skewed these results.

Author affiliations

¹Department of Orthodontics, Academisch Centrum Tandheelkunde Amsterdam (ACTA), University of Amsterdam, Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands; ²Department of Oral and Maxillofacial Surgery, Academic Medical Centre, University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands; ³Studio di Ortodonzia, Via Matteo Bandello 15, 20123 Milan, Italy. *Correspondence to: Reint A. Meursinge Reynders Email address: reyndersmail@gmail.com

References

- 1. Arn M L, Dritsas K, Pandis N, Kloukos D. The effects of fixed orthodontic retainers on periodontal health: Asystematic review. Am J Orthod Dentofacial Orthop 2020; **157**: 156–164.
- Liberati A, Altman D G, Tetzlaff J et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. PLoS Med 2009; 339: DOI: 10.1136/bmj.b2700.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009; 6: DOI: 10.1371/journal.pmed.1000097.
- Stevens A, Garritty C, Hersi M, Moher D. Developing PRISMA-RR, a reporting guideline for rapid reviews of primary studies (Protocol). Available at http://www.equator-network. org/wpcontent/uploads/2018/02/PRISMA-RR-protocol.pdf (accessed March 2020).
- Shea B J, Reeves B C, Wells G et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non randomised studies of healthcare interventions, or both. BMJ 2017; 358: DOI:10.1136/bmj.j4008.
- University of Bristol. Risk of bias in systematic reviews (ROBIS). ROBIS tool and the ROBIS guidance document. Available at http://www.bristol.ac.uk/social-community- medicine/projects/robis/ (accessed February 2020).
- Whiting P, Savović J, Higgins J P et al. ROBIS: A new tool to assess risk of bias in systematic reviews was developed. J Clin Epidemiol 2016; 69: 225–234.
- 8. PROSPERO: Centre for Reviews and Dissemination. University of York UK. Available at http://www.crd.york.ac.uk/PROSPERO/ (accessed February 2020).
- Kirkham J J, Altman D G, Williamson P R. Bias due to changes in specified outcomes during the systematic review process. PLoS One 2010; 5: DOI: 10.1371/journal. pone.0009810.
- Page M J, McKenzie J E, Kirkham J et al. Bias due to selective inclusion and reporting of outcomes and analyses in systematic reviews of randomised trials of healthcare interventions. Cochrane Database Syst Rev 2014; DOI: 10.1371/journal.pone.0009810.
- 11. Stewart L, Moher D, Shekelle P. Why prospective registration of systematic reviews makes sense. Syst Rev 2012; 1: 7.
- Cooke A, Smith D, Booth A. Beyond PICO: the SPIDER tool for qualitative evidence synthesis. Qual Health Res 2012; 22: 1435–1443.
- Golder S, Loke Y K, Wright K, Norman G. Reporting of adverse events in published and unpublished studies of health care interventions: a systematic review. PLoS Med 2016; 13: DOI: 10.1371/journal.pmed.1002127.

Evidence-Based Dentistry (2020) 21, 146-149. https://doi.org/ 10.1038/s41432-020-0144-0



CHAPTER 7

Which orthodontic retention protocol should I implement? A critical assessment of a randomized controlled trial

This is a non-edited version of the manuscript:

Steegmans PAJ, Cavagnetto D, Meursinge Reynders RA. Which orthodontic retention protocol should I implement? A critical assessment of a randomized controlled trial. Evid Based Dent. 2022 Dec;23(4):162-165.

A commentary on

Krämer A, Sjöström M, Apelthun C, Hallman M, Feldmann I. Post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers-a randomized controlled trial. Eur J Orthod 2022; DOI: 10.1093/ejo/cjac043.

Abstract

Trial design: A single-centre two-arm parallel group randomised controlled trial.

Objectives: To assess differences in dental stability, patient perceptions and compliance and retainer failures in adolescents treated with vacuum-formed retainers (VFR) compared with those receiving bonded canine-to canine retainers after five years in retention.

Methods: In total, 104 eligible adolescents treated with fixed appliances in both jaws in a Swedish orthodontic clinic were randomised to two retention protocols. The intervention protocol consisted of a VFR covering all erupted teeth in the maxilla and a VFR in the mandible covering first premolar to first premolar. The controls received a VFR in the maxilla covering all erupted teeth and a bonded retainer wire to the lingual surfaces of the canines. The primary outcomes were various dental stability measures assessed at: debond (T1); six months (T2); 18 months (T3); and after five years (T4) in retention. Generalised estimating equations were used to quantify the effect of the different interventions on these outcome measures. One operator assessed all outcomes and participants could not be blinded. For the secondary outcomes, the perception and compliance with the retention protocols were assessed and the prevalence and rationale of retainer failure at T4. The trial was registered at ClinicalTrials.gov (NCT03070444) and the research project was supported by the Centre for Research and Development, Region Gävleborg, Sweden.

Results: Of the 104 randomised patients, 30 were not available at T4, leaving 35 patients in the intervention and 39 in the comparator group. An intention-to-treat analysis was used to impute outcomes for the missing patients. Post-treatment changes at T4 were small in both jaws. In the maxilla, the Little's Irregularity Index (LII) increased similar in both retention groups (median difference: 0.3 mm). In the mandible, the median difference for the LLI in the bonded retainer group was 0.1 mm compared with 0.6 mm in VFR group. In both retention protocols, the overjet remained stable, the overbite increased and the arch lengths continued to decrease. Intercanine and intermolar width remained stable in the mandible. Intermolar width decreased significantly in the maxilla. No differences in satisfaction were found between retention protocols after five years. Also, 72% of patients had stopped or rarely wore the VFR appliances at T4. Besides some retainer failures in both groups, no serious adverse effects associated with the retainers were reported.

Conclusions: Most post-treatment changes in both retention protocols were small in both jaws, except for the anterior alignment in the mandible, which was more stable in the bonded retainer group. This difference is possibly not related to the retention technique but to the poor compliance with the VFRs and the inclusion of adolescents only. Satisfaction with both protocols was similar.

Practice Point

- At five years in retention, most post-treatment changes were small in both jaws for both retention protocols, except for the anterior alignment in the mandible, which was more stable in the bonded retainer group.
- The differences in the results are possibly not related to the appliances used, but to the poor compliance identified in the removable retainer group and to the inclusion of adolescents only, which tend to adhere less to wearing removable retainers compared with adults. The satisfaction with both retention protocols was similar.
- Before implementing the findings of this RCT in clinical practice, one should consider the high risk of bias in the results and additional limitations presented in Tables 4 and 5.

Commentary

Research questions

In this commentary we appraised a randomised controlled trial (RCT) by Krämer et al.¹ that assessed the following primary research question: 'in adolescent patients treated with fixed orthodontic appliances in the maxilla and mandible, how do post-treatment changes compare between two different retention protocols at debond (T1), after six months (T2), after 18 months (T3) and after five years (T4)'. The retention protocol in the intervention group was a vacuum-formed retainer (VFR) in the maxilla and mandible. The retention protocol in the comparator group was a VFR in the maxilla and a bonded retainer wire to lingual surfaces of the canines only. A secondary question was: 'what was the patient's perception and compliance regarding the different retention protocols and what was the prevalence and rationale of retainer failure after five years of retention?'. The elements 'participants', 'interventions', 'comparators', 'outcomes', 'time points', 'setting' of the PICOTS acronym of this RCT are reported in Table 1.

Items	Description
Participants	 Adolescent patients treated with fixed orthodontic appliances
Interventions	 VFR covering all erupted teeth in the maxilla and a VFR in the mandible covering first premolar to first premolar. VFR wear was gradually decreased to 1-2 rights per week only after two years of retention and on
Comparators	 VFR in the maxilla covering all erupted teeth and a bonded retainer wire to the lingual surface of the canines. VFR wear was gradually decreased to 1-2 nights per week only after two years of retention and on. The manibular retainer was not removed during the full retention period
Outcomes	 Primary outcomes: post-treatment dental and arch stability measures in the maxilla and mandible
	 Secondary outcomes: patient's perception and compliance regarding the different retention protocols and the prevalence and rationale of retainer failure
Time points	 Primary outcome: start of retention (T1), 6 months (T2), 18 months (T3), and after five years in retention (T4) Secondary outcomes: after five years in retention
Setting	 Orthodontic clinic in Gävle, public dental health service, Region Gävleborg, Sweden

Methods of the RCT

A total of 165 patients were eligible, of which 61 declined to participate in the trial. The remaining 104 eligible patients were then randomised to 52 patients in the intervention and 52 in the comparator arm. Digitised models were used to assess the stability of the dentition at the start of retention (T1), at six months (T2), at 18 months (T3), and after five years in retention (T4). Questionnaires were used to assess the perception and compliance with the retention protocols at T4. The prevalence and rationale of retainer failure after five years of retention was also measured. Treatment with fixed appliances was conducted by different orthodontists but all retention appliances were placed and outcomes were assessed by one orthodontist (the first author) during the entire five-year retention period.

Results of the RCT

At five years in retention, 30 of the 104 randomised patients were not available for outcome assessment, leaving, respectively, 35 patients in the intervention and 39 in the comparator group. An intention-to-treat analysis was implemented, which gave 50 patients in the intervention and 51 in the control group. Post-treatment changes at T4 in both retention protocols were small in both jaws. In the maxilla, the Little's Irregularity Index (LII) increased similarly in both retention groups (median difference: 0.3 mm). In the mandible, the median difference for the LLI in the bonded retainer group was 0.1 mm compared with 0.6 mm in VFR group. In both retention protocols, the overjet remained stable, the overbite increased and the arch lengths continued to decrease. Intercanine and intermolar width remained stable in the mandible. Intermolar width decreased significantly in the maxilla. At T4, patients were very satisfied with both the outcomes of treatment and the assigned retention protocols. Besides

some retainer failures in both groups, no serious adverse effects associated with the retainers were reported.

Methods for the critical appraisal

The checklist of the CONSORT statement² was used to assess how this RCT was reported and the risk of bias 2 (RoB2) tool³ was used to assess the risk of bias. All items in these tools were completed independently by two operators (PS and DC). The third operator (RMR) was consulted in the case of disagreements between these operators. Shortcomings that were not covered by these tools were also reported. All three authors developed and drafted this commentary and approved the final version of this manuscript.

Results of the critical appraisal

Our assessment of the reporting items according to the CONSORT statement² were reported in Table 2. In this table, we only list the items that were either 'not' or 'partially reported'. The results of the risk of bias assessment with the RoB2 tool³ are presented in Table 3 for each outcome separately. The limitations reported by the authors of the RCT are given in Table 4. All limitations identified with the CONSORT checklist², the RoB2 tool³ and additional limitations identified during our critical appraisal of this RCT are explained and summarised in Table 5.

Table 2 Items not reported according to the CONSORT statement ²		
CONSORT item	Type of reporting	
1b. Structured summary of trial design, methods, results and conclusions	Partially reported	
3b. Changes in the eligibility criteria compared with the protocol	Not reported	
6b. Changes in the trial outcomes compared with the protocol	Not reported	
16. Secondary outcome numbers analysed	Partially reported	
19. Harms	Partially reported	
20. Limitations	Partially reported	
21. Generalisability of the trial findings	Not reported	

Outcome	D1	D2	D3	D4	DS	Overall
Outcome 1	Low risk of bias	High risk of bias	Some concerns	High risk of bias	High risk of bias	High risk of bias
Outcome 2	Low risk of bias	High risk of bias	Some concerns	Low risk of bias	High risk of bias	High risk of bias
Outcome 3	Low risk of bias	High risk of bias	Some concerns	Low risk of bias	High risk of bias	High risk of bias
Outcome 4	Low risk of bias	High risk of bias	Some concerns	High risk of bias	High risk of bias	High risk of bias

Outcome 3: prevalence of compliance with wearing the VFR

Outcome 4: prevalence of retention failure in the different retention groups

Domain 1: bias arising from the randomisation process

Domain 2: bias due to deviations from the intended

Domain 4: bias in measurement of the outcome

Domain 5: bias in selection of the reported result

Table 4 Summary of the limitations identified by the authors of the RCT

Limitations reported by the authors	Description		
Binding issues	Complete blinding was not possible because of the type of interventions		
Little's Irregularity Index	The limitations of Little's Inregularity Index, such as not considering spacing or ro-tations with intact contact points		
Selection bias	Selection bias because patients who declined to participate in the trial could have been the poor compliers with the retention protocols		
Compliance issues	Differences in results are possibly not related to the retention technique but to the quality of compliance. Good compliance was required in the VFR group and no compliance in the fixed retainer group		
Hawthorne effect	False-positive bias because of the awareness of the participants being observed		
Duration of the trial	The long duration (five years) of the trial might have influenced compliance		

Table 5 Additional limitations identified during the critical appraisal of the RCT

Additional limitations identified during the critical appraisal	Description			
Misleading reporting related spin in the abstract ⁴	The poor compliance with the VFR was not reported in the abstract, which could mislead the reader			
Differences in the lengths of the retention devices	In the intervention group the VFR in the mandibular arch covered eight teeth, that is, first premolar to first premolar, while the comparator group had a shorter length covering only 6 teeth, that is, from canine to canine			
Deviations from the planned protocol	Differences between the two protocols (2009 and 2017) and with the final study were not reported nor the rationale for these changes. Deviations from the planned protocol refer to: The eligibility criteria of the 2017 protocol refer to child, adult and older adult. In the final RCT only adolescents were included Compliance and retainer failures were not defined as outcomes in the 2017 protocol = The 2009 protocol planned to assess secondary outcomes at four time points and not just at five years in retention as was reported in the final RCT = Primary outcomes were measured on digital models and not on dental casts as planned in the 2009 protocol The assessment of outcomes on adverse effects were not planned in the protocols, but in the final RCT the authors reported on adverse effects.			
Poor reporting on the definitions of secondary outcomes	Definitions of secondary outcomes such as satisfaction, retainer failure etc were not reported			
Poor reporting on adverse effects	In the final RCT, the authors reported that no serious adverse effects associated with the retainers were observed, but did not report which adverse effects were assessed, for example, no information was given regarding caries and periodontal problems			
Unreliable sample size calculation	The sample size calculation was not planned a priori and was based on an arbitrary value			
Incomplete reporting on the selection procedures of patients	It was not reported who selected the patients (preferably done by two operators) and how many patients were assessed for eligibility and how many of these patients did not meet the inclusion criteria and why they were excluded. The authors reported only the number of eligible patients			
Bias in measurement of the outcome	Outcomes were assessed by one outcome assessor only, who also placed all retainers and conducted all follow-up procedures			
Bias due to deviations from the intended intervention	Many patients (72%) had already stopped wearing the VFRs at the last follow-up, due to unknown reasons, which implies poor adherence to the intervention			
Lost to follow up	High prevalence of patients, that is, 29% (30/104), were lost to follow-up			
Caution regarding the generalisability of the trial findings	Generalisability of the trial findings should be done with caution, because: Only adolescents were included. This implies that this study cannot be generalised to adults, because adults possibly comply differently with VFRs Selection bias The incomplete reporting on the selection procedures of patients 			
Raw data	Raw data were not given as additional files or in a repository and could only be obtained on 'reasonable request to the corresponding author'			

appraisal, which were not already reported by the authors of the RCT.

Author affiliations

¹PhD Student, Department of Orthodontics, Academisch Centrum Tandheelkunde Amsterdam, University of Amsterdam, Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands;

²PhD Student, Department of Oral and Maxillofacial Surgery, Amsterdam University Medical Centre, Location AMC, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands;

³Studio di Ortodonzia, Via Matteo Bandello 15, 20123 Milan, Italy;

⁴Senior Scientist and Associate Editor, Department of Oral and Maxillofacial Surgery, Amsterdam University Medical Centre, Location AMC, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands.

*Correspondence to: Reint A. Meursinge Reynders

Ethics declaration

The authors declare no conflicts of interest.

References

- Krämer A, Sjöström M, Apelthun C, Hallman M, Feldmann I. Post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers-a randomized controlled trial. Eur J Orthod 2022; DOI: 10.1093/ejo/cjac043.
- Schulz K F, Altman D G, Moher D, CONSORT Group. CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. Ann Intern Med 2010; 152: 726–732.
- Sterne J A C, Savović J, Page M J et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ 2019; DOI: 10.1136/bmj.l4898.
- 4. Lazarus C, Haneef R, Ravaud P, Boutron I. Classification and prevalence of spin in abstracts of nonrandomized studies evaluating an intervention. BMC Med Res Methodol 2015; 15: 85.

Evidence-Based Dentistry (2022) 23, 162-165. https://doi.org/10.1038/s41432-022-0845-7



CHAPTER 8

Fixed flexible spiral wire retainers and unwanted tooth movements: a Case Report

This is a non-edited version of the manuscript:

Steegmans PAJ, Jonkman REG, de Lange J. Fixed Flexible Spiral Wire Retainers and Unwanted Tooth movements: A Case Report. Appl. Sci. 2023, 13(2), 922.

Abstract

This case report presents a study of unwanted tooth movements during the retention phase after orthodontic treatment. The early recognition of these unwanted tooth movements is paramount for patients and clinicians to prevent the associated negative consequences. A 21-year-old male presented with aesthetic complaints regarding his upper front teeth. He underwent orthodontic treatment at the age of 9 years and 11 months and finished his treatment 2 years and 11 months later. Flexible spiral wires (FSW) were bonded to the anterior segment of the upper and lower jaws to stabilize the end result. The failure of the fixed retainers had never occurred previously. The diagnostic assessment demonstrated a previously orthodontically treated class I malocclusion with excessive angulation and torque differences in the maxillary anterior segment. To correct the position of the maxillary anterior segment and prevent further misalignment, the patient received orthodontic retreatment. Thereafter, the result was retained with fixed braided-rectangular-wire (BRW) retainers located at 12–22 and 33–43 and a vacuum-formed retainer (VFR) in the maxilla. The end result appeared to be stable after 28 months of retention. Unwanted tooth movements can occur during the orthodontic retention phase and might result from the use of fixed flexible spiral wire retainers. Follow-up appointments are recommended to monitor the stability and recognize these movements.

Keywords: orthodontics; dentistry; adverse effects; stability; fixed retainers; retention; unwanted tooth movements

1.Introduction

The retention phase of an orthodontic treatment aims to maintain the end result of orthodontic treatment and is of great importance for both patients and orthodontists. In 1934, Oppenheim stated: "Retention is one of the most difficult problems in orthodontia; in fact, it is the problem" [1]. Without retention, teeth have the possibility of (1) returning to their initial position, also known as 'relapse' [2], or (2) displacement as a result of growth and aging [3]. Orthodontic retention can be performed in two ways: (1) by placing fixed or removable appliances on the teeth or (2) by additional treatments of the teeth and periodontal structures in order to achieve stability. Fixed retainers are usually bonded to the palatal or lingual side of the anterior teeth [4]. Hawley or vacuum-formed retainers (VFR) can be taken off by the patient. Additional treatments used to prevent relapse are (1) small surgical procedures that cut the supra-crestal periodontal fibers around the teeth to reduce the chance of relapse [5] or (2) interproximal enamel reduction of the lower front teeth to create space for the corrected crowding and to compensate in advance for the expected reduction in the inter-canine width during aging [6].

Currently, fixed retention is commonly used in orthodontic practice [7,8]. Previous research on the long-term outcomes showed that when using a flexible spiral wire (FSW) retainer bonded to the mandibular anterior segment, the alignment is stabilized in 90.5% (200/221) of cases after 5 years of retention [9]. However, these fixed retention appliances may also fail or have adverse effects. Recently, a systematic review was conducted to evaluate the available evidence on the failure of fixed retainers and reported that fixed retainers fail in a range of 7.3% to 50%, according to which detachment at the adhesiveenamel interface was the most commonly reported type of failure [10]. Other observed adverse effects of fixed retainers in the orthodontic literature are detachment at the adhesive-wire interface [11], wire fracture [12], wire untwisting [13], and calculus accumulation [14,15]. Wire untwisting may lead to unwanted changes in the tooth position and can be associated with the development of gingival recessions [13,16,17]. Other causes of unwanted tooth movements can be tongue thrust or personal habits [18,19]. A clinical case of unwanted tooth movements during the retention phase is presented in this case report. This case showed that the unwanted tooth movement was the direct result of an untwisting FSW, because the maxillary teeth were displaced in a different direction from the original tooth position (Figure 1), which clearly showed that this was not a case of simple relapse. The early recognition of these unwanted tooth movements is paramount for patients and clinicians to prevent the associated negative consequences [20].



Figure 1. Extra-oral and intra-oral photographs before first orthodontic treatment.

2. Materials and Methods

This case was reported according to the Case Report (CARE) guidelines [21]. The CARE checklist of items used for this case report is presented in Supplementary File S1.

Patient information

A 21-year-old Caucasian male presented with aesthetic complaints regarding his upper front teeth. He previously underwent orthodontic treatment for a deep bite at the age of 9 years and 11 months for 2 years and 11 months. He was treated with high-pull headgear (Headgear, Dentsply GAC International, NY, USA) followed by full fixed appliances (3M Victory Series APC conventional twin brackets, 3M Health Care Division, London, Canada), and flexible spiral wires (0.0195-inch, 3-strand, heat-treated twist wire, Wildcat, GAC International, Bohemia, NY, USA) were bonded to all the anterior teeth between 13–23 and 33–43 to stabilize the end result (Figure 2). Over the last few years (the moment of onset was unknown), he noted a continuing shift in the position of his upper front teeth (Figure 3A). Aside from his aesthetic complaints, he was concerned that the situation would deteriorate. With regard to his medical history, he only used antihistamines for hay fever if necessary. There was no record of dental trauma, and he was unfamiliar with oral parafunctions. The failure of the fixed retainers had never occurred, and he did not experience any pain or functional constraints.



Figure 2. Extra-oral and intra-oral photographs after first orthodontic treatment.



(A)





(C)



(D)

Figure 3. Documentation before orthodontic re-treatment. (A) Extra-oral and intra-oral photographs. (B) Dental casts. (C) Lateral cephalometric radiograph. (D) Panoramic radiograph.

Clinical findings

Figure 3A shows the case at the first presentation. The photographs demonstrate that the 13 shows excessive palatal root torque, and the 23 shows excessive buccal root torque. A cant of the upper incisors can be observed. Moreover, cross-bites are present between the 14 and the 44 and between the 23 and the 34.

Timeline

The timeline for this patient is presented in Supplementary File S2.

Diagnostic Assessment

The patient presented with a Class I profile and a chin point deviation to the right side (Figure 3A). The intra-oral assessment showed good oral hygiene, a thin gingival biotype, group function on the left and right side, and fixed retainers (0.0195-inch, 3-strand, heattreated twist wire, Wildcat, GAC International, Bohemia, NY, USA) in the upper anterior segment and lower anterior segment (Figure 3A). The dental cast assessment showed a Class I molar and canine occlusion, an overjet of 2 mm, and an overbite of 3 mm. A cant of the upper incisors was present with palatal root torgue of the 13 and buccal root torgue of the 23. Crossbites were present between the 14/44 and the 23/34. The arch length discrepancy measurement resulted in values of 0 mm in the case of the maxillary arch and -4 mm in the case of the mandibular arch. No tooth size discrepancy was present. The PAR index resulted in a score of 12 points [22] (Figure 3B). The lateral cephalometric radiograph showed a Class I intermaxillary relationship with an ANB angle of 3.2° (VistaPano S Ceph, Dürr Dental imaging software, Dürr Dental SE, Bietigheim-Bissingen, Germany) (Figure 3C). The panoramic radiograph (VistaPano S Ceph, Dürr Dental imaging software, Dürr Dental SE, Bietigheim-Bissingen, Germany) showed a maxillary sinus mucosal cyst (MSMC) on the right side and external apical root resorption located at the 12, 11, and 21 (Figure 3D).

Diagnosis

This clinical, radiographic, and dental cast examination contributed to the following orthodontic diagnosis: A 21-year-old male with an Angle Class I malocclusion, angulation, and torque differences in the maxillary anterior segment, with cross-bites located at 14/44 and 23/34, a thin gingival biotype, external apical root resorption located at 12, 11, and 21, and the presence of fixed retainers (0.0195-inch, 3-strand, heat-treated twist wire, Wildcat, GAC International, Bohemia, NY, USA) located between 13–23 and 33–43. 2.2.

Prognosis

The torque and angulation differences in the maxillary anterior segment were not observed before (Figure 1) or after the orthodontic treatment from 2006 to 2009 (Figure 2). From 2009 to 2017, no failure of the fixed retainers had occurred, and no oral parafunctions were present. It was hypothesized that the FSW retainer caused these unwanted tooth movements [9,16]. Since the patient identified a continued worsening of the position of the upper front teeth, it is likely that this process of unwanted tooth movement was persistent

8

and consequently resulted in a worse prognosis. For example, the development of gingival recessions may occur [17]. A recent case report showed a similar situation that resulted in the exposure of the apex of a canine [20]. A pilot study based on a retrospective data analysis showed that the removal of the FSW retainer led to the cessation of further unwanted tooth movement [23].

Therapeutic intervention

To correct the position of the maxillary anterior segment and prevent further misalignment, the patient received orthodontic re-treatment for 2 years and 4 months. The fixed FSWs (0.0195-inch, 3-strand, heat-treated twist wire, Wildcat, GAC International, Bohemia, NY, USA) in the maxilla and mandible were removed. Afterward, fixed appliances were placed (3M Victory Series APC conventional twin brackets, 3M Health Care Division, London, Canada). To correct the excessive palatal root torgue of the 13, we incorporated the 13 into the fixed appliances with a 0.012-inch NiTi overlay wire (Sentalloy, Dentsply GAC international, Bohemia, NY, USA) on a 0.016 × 0.022-inch SS base wire (Stainless Steel Ideal® Form, Dentsply GAC international, Bohemia, NY, USA). In this way, the risk of unwanted reactionary tooth movements was reduced. During the active orthodontic treatment, a panoramic radiograph (VistaPano S, Dürr Dental imaging software, Dürr Dental SE, Bietigheim-Bissingen, Germany) was performed to monitor the root resorption observed in the maxillary anterior region in the initial stage of the treatment [24] (Figure 4). It appeared to be stable. However, it was difficult to reach a conclusion in this regard, since (1) a great difference in the tooth position between the two panoramic radiographs and (2) variations in quality between the panoramic radiographs were observed due to the use of different radiographic devices. The oral hygiene was checked at every orthodontic appointment. If necessary, instructions were given to maintain the quality of the patient's oral hygiene during the orthodontic treatment. After active orthodontic treatment over 2 years and 4 months, the treatment result was stabilized with fixed braided-rectangular-wire (BRW) retainers (Forestaflex, Forestadent[®], Bernhard Förster GmbH, Pforzheim, Germany) located at 12–22 and 33–43. We opted for the BRW retainers, which achieve better torque control compared to flexible spiral wires (FSW) [25]. In addition, a VFR was prescribed for approximately 10 h a day to cover all the maxillary teeth. According to the current literature, the part-time wear of the VFR should be sufficient enough to generate stability [26]. Table 1 provides a detailed overview of the interventions.




Figure 4. Panoramic radiograph during orthodontic treatment for monitoring root resorption.

 Table 1. Interventions for the orthodontic re-treatment.

Intervention	Description
Removal of the fixed flexible spiral wires	Maxilla: 0.0195-inch, 3-strand, heat-treated twist wire (Wildcat, GAC International, Bohemia, NY, USA) located on the palatal surface 13-12-11-21-22-23 Mandible: 0.0195-inch, 3-strand, heat-treated twist wire (Wildcat, GAC International, Bohemia, NY, USA) located on the lingual surface 33-32-31-41-42-43
Fixed appliance therapy	Type: 3M Victory Series APC conventional twin brackets (3M Victory Series APC conventional twin brackets, 3M Health Care Division, London, Canada) Prescription: MBT Slot system: 0.022-inch Wires: 0.012-inch NITI *, 0.014-inch NITI *, 0.016-inch NITI *, 0.018-inch NITI *, 0.016 × 0.022-inch NITI *, 0.017 × 0.025-inch NITI *, 0.016 × 0.022-inch SS **, 0.017 × 0.025-inch SS ** Overlay wires: 0.012-inch NITI *, 0.014-inch NITI *, 0.016-inch NITI *, 0.018-inch NITI *
Retention therapy	Maxilla: 0.016 X 0.016-inch, 8-stranded, braided rectangular wire (Forestaflex, Forestadent [®] Bernhard Förster GmbH, Pforzheim, Germany) located on the palatal surface of 12-11-21-22 in combination with a vacuum-formed retainer Mandible: 0.016 × 0.022-inch, 8-stranded, braided rectangular wire (Forestaflex, Forestadent [®] , Bernhard Förster GmbH, Pforzheim, Germany) located on the lingual surface of 33-32-31-41-42-43

: Dentspiy GAC Sentalloy (Dentspiy GAC international, Bohemia, NY, USA), **: Dentspiy GAC Stamless Ideal[®] Form (Dentspiy GAC international, Bohemia, NY, USA).

3. Results

3.1. Follow-Up and Outcomes

3.1.1. Clinician- and Patient-Assessed Outcomes

The end result of the orthodontic re-treatment is shown in Figure 5. The cant of the upper incisors, the excessive palatal and buccal root torgue of the 13 and 23, and the crossbites were corrected. In addition to aesthetic improvement, cuspid guidance on the left and right sides was achieved. However, the panoramic radiograph showed an increase in root resorption (Figure 4), and a buccal gingival recession was observed at the 23 (Figure 5). To assess the outcome of this orthodontic re-treatment, the Peer Assessment Rating (PAR) index was consulted and resulted in a PAR score of 2 [22]. Compared to the PAR score measured before the orthodontic re-treatment (PAR score of 12), the malocclusion improved by 83% (10/12). The patient participated in the ESAS patient satisfaction evaluation guestionnaire to assess the outcome of the orthodontic retreatment. ESAS (EFOSA Self Assessment System) is a quality assessment system available to all orthodontists in Europe [27]. This patient satisfaction evaluation questionnaire rates different aspects of the treatment, e.g., patient satisfaction with the orthodontist, assistants, practice, and overall treatment. The completed questionnaire is provided (Supplementary File S3). The answers indicated that the patient was satisfied with nearly all aspects. Nevertheless, the treatment duration was slightly longer than expected, and he believed that the waiting room was not comfortable enough.



Figure 5. Extra-oral and intra-oral photographs after orthodontic re-treatment

3.1.2. Important Follow-Up Test Results

Since the patient went abroad for 3 years, the first follow-up appointment was scheduled for April 2022 (28 months after the end of the orthodontic re-treatment). At that moment, the bonded retainers were still fixed and had not been displaced. He still wore the VFR for approximately 10 h a day. An intra-oral examination showed a class I occlusion with an overjet of 2 mm and an overbite of 3 mm (Figure 6). These findings implied that the end result of the orthodontic re-treatment seemed to be stable. Unfortunately, the patient's oral hygiene was insufficient because, plaque and calculus were present around the fixed retainer in the mandibular anterior segment.



Figure 6. Extra-oral and intra-oral photographs 28 months after orthodontic re-treatment.

3.1.3 Intervention Adherence and Tolerability

The patient was compliant with the intervention. The oral hygiene was sufficient during the orthodontic treatment, and all the check-up appointments were undertaken. According to the patient, his aesthetic concerns and awareness of the need for orthodontic treatment could be attributed to this adherence.

3.1.4. Adverse and Unanticipated Events

During the active orthodontic re-treatment, an increase in root resorption occurred, and a buccal gingival recession was observed at the 23.

4. Discussion

This case report illustrates an adverse effect in orthodontics. This concerned unwanted tooth movement during the retention phase, probably due to the fixed retainer in the upper jaw. Even though fixed retainers are effective in preventing relapse, previous research has shown that fixed FSWs may also lead to unwanted effects and can complicate oral hygiene, with negative consequences for the periodontium [9,16,28–30]. Katsaros et al. described two possible effects of fixed FSWs during the retention phase: (1) the X-effect, a torque difference between two adjacent mandibular incisors, and (2) the twist-effect, an increase in the buccal inclination of the canine [16]. The X-effect or twist effect was observed in 2.7% (6/221) of the evaluated patients during a retention period of 5 years [9]. Recently, Singh described a case in which a combination of the X-effect and twist-effect was present, leading to the avulsion of the canine [20]. These reported 'twist-effects' are similar to the tooth movements described in this case report. However, in the current case report, the fixed FSW retainer was located in the upper jaw, while previous studies investigated fixed FSWs in the lower jaw [9,16,20,28]. It is hypothesized that the forces that contribute to the etiology of these unwanted tooth movements might be generated in three ways: (1) by the untwisting of the round flexible spiral wires (tooth movements generated by these forces might also be the reason for the observed root resorption localized at the 12, 11, and 21 on the panoramic radiograph from July 2017 (Figure 3D) [31]), (2) the mechanical deformation of the wire as a result of masticatory forces, and (3) the elastic deflection of the wire due to an inadequate passive bonding procedure [25]. One or more of the aforementioned etiologic factors might explain the situation described in this case report. To prevent a recurrence of these unwanted tooth movements, BRWs were bonded to the upper and lower jaws after the second orthodontic treatment. Fiber-reinforced composite or polyethylene splints could represent a viable alternative to conventional metallic bonded retention [32]. However, these materials have been shown to be less clinically reliable over time than stainless steel retainers [33]. A rectangular chain retainer could also have been a good alternative for the fixed retention in the upper and lower jaws. However, according to Arnold et al., BRWs achieve higher torque control [25]. Therefore, we chose to apply BRWs. Due to occlusal interferences with the lower canine, the retainer in the upper jaw was bonded to the incisors alone. However, it has been shown that significantly fewer rotational changes occur when fixed retainers are bonded from 13 to 23 compared to fixed retainers bonded from 12 to 22 [34]. Therefore, a VFR was prescribed to prevent rotational changes of the 13 and 23. According to a systematic review conducted by Bellini-Pereira et al., fixed retainers and VFRs in the upper jaw are equally effective in maintaining the end result of an orthodontic treatment (with a moderate level of certainty) [14]. Hence, a VFR placed in the upper jaw should stabilize the cuspids adequately. It has been shown that the failure of fixed orthodontic retainers occurs more frequently in the upper jaw compared to the lower jaw [10]. Therefore, if the fixed retainer fails from 12 to 22, the VFR will stabilize the alignment until the retainer can be repaired. The limitations of this case report include (1) the delay of the first follow-up appointment, which was eventually scheduled after 28 months due to the

patient's departure abroad and the COVID-19 pandemic. Fortunately, the fixed retainers did not fail in this period. However, the first follow-up appointment should have been scheduled earlier, since most failures of fixed retention occur in the first 6 months after the bonding of fixed retainers. In addition, failure is more frequent in the upper jaw [10]. (2) The existing root resorption should have been monitored earlier and more often during the active orthodontic re-treatment. (3) Moreover, case reports are considered to represent the lowest level of evidence. Implications for future research could include the design of studies with a long-term follow-up that assess variables that could influence the stability of fixed orthodontic retention, such as the periodontal and dental status of the patient, the type and dimensions of the retainer wire, the type and quantity of the composite used, and the placement technique.

5. Conclusions

In conclusion, fixed retainers appear to be effective in maintaining alignment after orthodontic treatment. However, unwanted tooth movements might result from the use of fixed flexible spiral wire retainers, requiring orthodontic re-treatment. Follow-up appointments are recommended to monitor the stability and to recognize unwanted tooth movements at an early stage. The results of this case report are of great importance for researchers, patients, and clinicians. In particular, dentists should be aware of these possible adverse effects of fixed flexible spiral wire retainers, since they regularly perform dental check-ups after orthodontic treatment. Therefore, dentists play an important role in recognizing these unwanted tooth movements.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/app13020922/s1, Supplementary File S1: CARE Checklist of information to include when writing a case report. Supplementary File S2: Timeline. Supplementary File S3: ESAS patient satisfaction evaluation part.

Author Contributions: This case report was conceived and designed by P.A.J.S. The final submitted manuscript was reviewed, revised, and approved by all three authors. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding. The APC was funded by the Academic Center for Dentistry Amsterdam.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Written informed consent was obtained from the patient to publish this paper.

Data Availability Statement: The dataset used for this case report is available from the corresponding author.

Acknowledgments: We thank the patient for his participation in this case report.

Conflicts of Interest: The authors declare no conflict of interest.

136

References

- 1. Oppenheim, A. Crisis in orthodontia. Part I. Tissue changes during retention. Int. J. Orthod. 1934, 20, 639–644.
- Thilander, B. Orthodontic relapse versus natural development. Am. J. Orthod. Dentofac. Orthop. 2000, 117, 562–563. [CrossRef] [PubMed]
- Massaro, C.; Miranda, F.; Janson, G.; Rodrigues de Almeida, R.; Pinzan, A.; Martins, D.R.; Garib, D. Maturational changes of the normal occlusion: A 40-year follow-up. Am. J. Orthod. Dentofac. Orthop. 2018, 154, 188–200. [CrossRef] [PubMed]
- 4. Knierim, R.W. Invisible lower cuspid to cuspid retainer. Angle Orthod. 1973, 43, 218–220. [PubMed]
- Al-Jasser, R.; Al-Subaie, M.; Al-Jasser, N.; Al-Rasheed, A. Rotational relapse of anterior teeth following orthodontic treatment and circumferential supracrestal fiberotomy. Saudi Dent. J. 2020, 32, 293–299. [CrossRef] [PubMed]
- 6. Aasen, T.O.; Espeland, L. An approach to maintain orthodontic alignment of lower incisors without the use of retainers. Eur. J. Orthod. 2005, 27, 209–214. [CrossRef] [PubMed]
- 7. Fudalej, P.S.; Renkema, A.M. A brief history of orthodontic retention. Br. Dent. J. 2021, 230, 777–780. [CrossRef] [PubMed]
- Padmos, J.A.D.; Fudalej, P.S.; Renkema, A.M. Epidemiologic study of orthodontic retention procedures. Am. J. Orthod. Dentofac. Orthop. 2018, 153, 496–504. [CrossRef] [PubMed]
- Renkema, A.M.; Renkema, A.; Bronkhorst, E.; Katsaros, C. Long-term effectiveness of canine-to-canine bonded flexible spiral wire lingual retainers. Am. J. Orthod. Dentofac. Orthop. 2011, 139, 614–621. [CrossRef] [PubMed]
- Jedliński, M.; Grocholewicz, K.; Mazur, M.; Janiszewska-Olszowska, J. What causes failure of fixed orthodontic retention?— Systematic review and meta-analysis of clinical studies. Head Face Med. 2021, 17, 32. [CrossRef]
- Gelin, E.; Seidel, L.; Bruwier, A.; Albert, A.; Charavet, C. Innovative customized CAD/CAM nickeltitanium lingual retainer versus standard stainless-steel lingual retainer: A randomized controlled trial. Korean J. Orthod. 2020, 50, 373–382. [CrossRef] [PubMed]
- 12. Radlanski, R.J.; Zain, N.D. Stability of the Bonded Lingual Wire Retainer? A Study of the Initial Bond Strength. J. Orofac. Orthop. 2004, 65, 321–335. [CrossRef] [PubMed]
- 13. Shaughnessy, T.G.; Proffit, W.R.; Samara, S.A. Inadvertent tooth movement with fixed lingual retainers. Am. J. Orthod. Dentofac. Orthop. 2016, 149, 277–286. [CrossRef] [PubMed]
- Bellini-Pereira, S.A.; Aliaga-Del Castillo, A.; Dos Santos, C.C.O.; Henriques, J.F.C.; Janson, G.; Normando, D. Treatment stability with bonded versus vacuum-formed retainers: A systematic review of randomized clinical trials. Eur. J. Orthod. 2022, 44, 187–196. [CrossRef]
- Juloski, J.; Glisic, B.; Vandevska-Radunovic, V. Long-term influence of fixed lingual retainers on the development of gingival recession: A retrospective, longitudinal cohort study. Angle Orthod. 2017, 87, 658–664. [CrossRef]
- 16. Katsaros, C.; Livas, C.; Renkema, A.M. Unexpected complications of bonded mandibular lingual retainers. Am. J. Orthod. Dentofac. Orthop. 2007, 132, 838–841. [CrossRef]
- Renkema, A.M.; Fudalej, P.S.; Renkema, A.A.P.; Abbas, F.; Bronkhorst, E.; Katsaros, C. Gingival labial recessions in orthodontically treated and untreated individuals: A case-control study. J. Clin. Periodontol. 2013, 40, 631–637. [CrossRef]
- Seo, Y.J.; Kim, S.J.; Munkhshur, J.; Chung, K.R.; Ngan, P.; Kim, S.H. Treatment and retention of relapsed anterior open-bite with low tongue posture and tongue-tie: A 10-year follow-up. Korean J. Orthod. 2014, 44, 203–216. [CrossRef]
- Sfondrini, M.F.; Pascadopoli, M.; Beccari, S.; Beccari, G.; Rizzi, C.; Gandini, P.; Scribante, A. Orthodontic Fixed Retainer and Unwanted Movements of Lower Anterior Teeth: Case report. Case Rep. Dent. 2022, 2, 3100360. [CrossRef]
- 20. Singh, P. Canine avulsion: An extreme complication of a fixed mandibular lingual retainer. Am. J. Orthod. Dentofac. Orthop. 2021, 160, 473–477. [CrossRef]
- Riley, D.S.; Barber, M.S.; Kienle, G.S.; Aronson, J.K.; von Schoen-Angerer, T.; Tugwell, P.; Kiene, H.; Helfand, M.; Altman, D.G.; Sox, H.; et al. CARE guidelines for case reports: Explanation and elaboration document. J. Clin. Epidemiol. 2017, 89, 218–235. [CrossRef]

- Richmond, S.; Shaw, W.C.; O'Brien, K.D.; Buchanan, I.B.; Jones, R.; Stephens, C.D.; Roberts, C.T.; Andrews, M. The development of the PAR Index (Peer Assessment Rating): Reliability and validity. Eur. J. Orthod. 1992, 14, 125–139. [CrossRef] [PubMed]
- Knaup, I.; Bartz, J.R.; Schulze-Späte, U.; Craveiro, R.B.; Kirschneck, C.; Wolf, M. Side effects of twistflex retainers—3D evaluation of tooth movement after retainer debonding. J. Orofac. Orthop. 2021, 82, 121–130. [CrossRef] [PubMed]
- Årtun, J.; Van 't Hullenaar, R.; Doppel, D.; Kuijpers-Jagtman, A.M. Identification of orthodontic patients at risk of severe apical root resorption. Am. J. Orthod. Dentofac. Orthop. 2009, 135, 448–455. [CrossRef] [PubMed]
- 25. Arnold, D.T.; Dalstra, M.; Verna, C. Torque resistance of different stainless steel wires commonly used for fixed retainers in orthodontics. J. Orthod. 2016, 43, 121–129. [CrossRef]
- Littlewood, S.J.; Millett, D.T.; Doubleday, B.; Bearn, D.R.; Worthington, H.V. Retention procedures for stabilising tooth position after treatment with orthodontic braces. Cochrane Database Syst. Rev. 2016, 1, 1–139. [CrossRef]
- 27. EFOSA's Self Assessment System Quality Development in Orthodontics. Available online: https://esas.nu/ (accessed on 6 June 2022).
- Kučera, J.; Marek, I. Unexpected complications associated with mandibular fixed retainers: A retrospective study. Am. J. Orthod. Dentofac. Orthop. 2016, 149, 202–211. [CrossRef]
- 29. Pazera, P.; Fudalej, P.; Katsaros, C. Severe complication of a bonded mandibular lingual retainer. Am. J. Orthod. Dentofac. Orthop. 2012, 142, 406–409. [CrossRef]
- 30. Kartal, Y.; Kaya, B. Fixed Orthodontic Retainers: A review. Turk. J. Orthod. 2019, 32, 110–114. [CrossRef]
- 31. Segal, G.R.; Schiffman, P.H.; Tuncay, O.C. Meta analysis of the treatment-related factors of external apical root resorption. Orthod. Craniofac Res. 2004, 7, 71–78. [CrossRef]
- Karaman, A.I.; Kir, N.; Belli, S. Four applications of reinforced polyethylene fiber material in orthodontic practice. Am. J. Orthod. Dentofac. Orthop. 2002, 121, 650–654. [CrossRef] [PubMed]
- Scribante, A.; Sfondrini, M.F.; Broggini, S.; D'Allocco, M.; Gandini, P. Efficacy of Esthetic Retainers: Clinical Comparison between Multistranded Wires and Direct-Bond Glass Fiber-Rein forced Composite Splints. Int. J. Dent. 2011, 2011, 548356. [CrossRef] [PubMed]
- Naraghi, S.; Ganzer, N.; Bondemark, L.; Sonesson, M. Stability of maxillary anterior teeth after 2 years of retention in adolescents: A randomized controlled trial comparing two bonded and a vacuumformed retainer. Eur. J. Orthod. 2021, 43, 152–158. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.



CHAPTER 9

General discussion

This chapter is divided into three sections. Section 1 presents the principal findings of this dissertation and how they compare with other studies. Section 2 presents the strengths and limitations of this dissertation and section 3 presents the meaning of the findings of this dissertation (possible explanations and implications for clinicians and policymakers), reported unanswered questions, and future research.

Section 1. Principal findings of this dissertation and how they compare with other studies

Chapter 2 presents the protocol for the cross-sectional study on defining, seeking and reporting of adverse effects in systematic reviews of orthodontic interventions. The pilot studies in this protocol confirmed the importance of our research questions. The cross-sectional study (chapter 3) identified 98 eligible systematic reviews of orthodontic interventions. The results showed that a small proportion (36% (35/98)) of these systematic reviews defined seeking adverse effects as an objective. Findings related to adverse effects of interventions were sought in 86% (84/98) of reviews and reviewers reported on these effects in 85% (83/98) of reviews. In 91% (89/98) of the included systematic reviews, the reviewers discussed (weighed) potential adverse effects of interventions anywhere in the review. These outcomes were higher compared to those identified in similar investigations in gastroenterology (67% (52/78)) [1], drug intervention systematic reviews (76% (59/78)) [2], and in Database of Abstracts of Reviews of Effects (DAREs) reviews (48% (38/79)) [2]. The research design, type of intervention, the field of research, and different periods of inclusion of systematic reviews, can explain the differences in the reported proportions between these studies and our results (chapter 3).

A wide variety of different types of adverse effects (n=195) were identified. Based on these findings, we modified the framework of known orthodontic adverse effects reported by Preoteasa et al. [3] and developed a new framework for defining adverse effects of orthodontic interventions (Table 3). Systematic reviews of orthodontic interventions sought and reported predominantly (83% (162/195)) on five types of adverse effects, i.e., (1) tooth structures, (2) periodontal tissues, (3) undesired treatment results, (4) relapse and stability, and (5) negative qualitative experiences by the patient or carer(s).

Adverse effects related to	Description
Tooth structures	Tooth crown
	 decalcifications, decays, tooth wear, enamel cracks and fractures; discolorations, deterioration of prosthetic crown (as fracturing a ceramic one during debonding); iatrogenic damage to the crown, e.g., fracture as a result of trauma tooth root root resorption, early closure of root apex, ankylosis; iatrogenic damage to the root, e.g., fracture as a result of trauma tooth pulp ischemia, pulpitis, necrosis iatrogenic damage to the pulp, e.g., fracture as a result of trauma
Periodontal tissues	 gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscences, fenestrations, interdental fold, dark triangles; tooth mobility, plague retention, bacterial count
Intraoral (non-tooth or periodontal) tissues	 intraoral tissue irritations and inflammation such as mucosal ulcerations or hyperplasia or irritations of the tongue (as a result of trauma by appliances, e.g., breakage, failure, loosening etc. of appliances or long arch wires) Scar formation after suturing chemical burns (e.g., etching related) thermal injuries (e.g., overheated burs) nerve damage tooth eruption, i.e., eruption disturbances (e.g. impactions) caused by orthodontic appliances
Extraoral tissues (non- temporomandibular tissues)	 cutting of lips or cheeks, eye injury (e.g., as a result of trauma by appliances, e.g., breakage, failure, loosening etc. of appliances or long arch wires or headgear-related trauma) discomfort on the lip
Temporomandibular tissues and disorders	temporomandibular tissues and disorders
Appliance failure	 breakage, failure, loosening etc. of appliances long archwires, headgear-related trauma
Undesired treatment results	 inadequate morpho-functional, aesthetic or functional final result inaccuracy of the treatment result non predictability of the treatment result Dental side effects e.g. unwanted tipping of teeth, anchorage loss etc. Skeletal side effects, e.g., unwanted backward rotation of the mandible
Relapse and stability	 Relapse and stability of the obtained treatment result
Undesired qualitative experiences by the patient or carer(s)	 Pain and discomfort orthodontic tooth movement-related pain and discomfort appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g., tension or pressure of the appliances (constriction of appliances), speech difficulties, eating difficulties, swallowing difficulties, food accumulation, bad tastes and smells additional intervention-related pain and discomfort, e.g., surgical and non-surgical adjunctive interventions to accelerate tooth movement Tolerability/acceptance/stress issues with the treatment procedures

Table 3. Adverse effects hypothetically linked to orthodontic interventions*

	this table
	could not be labeled under any of the categories of adverse effects given in
Additional adverse effects	Additional adverse effects that were identified during data extraction that
Non defined	referring to 'any adverse effect', 'any side effect' etc.
Non-defined	Adverse effects that were not defined by the authors of the review:
Cross infections	 from doctor to patient, patient to doctor, patient to patient.
Chronic fatigue	
Cardio	infective endocarditis;
Allergy	Allergies to nickel or latex;
	brackets);
Gastro-intestinal	accidental swallowing parts of the orthodontic device (tubes,
	measured, i.e., during or after)
	 was measured, i.e., during or after) not satisfied with the final treatment result (Check in text what was
	 not satisfied with the treatment procedures (Check in text what was measured i.e. during or after)
	Satisfaction with the treatment procedures and final result
	Catisfaction with the twenty at more during and final years'
	development of mannerisms
	removal of appliance during sleep
	sleeping difficulties
	 reduced enjoyment of food and change in taste
	concentration difficulties
	aesthetic look discontents during orthodontic appliance usage
	relationships
	 behavioral changes of patients and parents, impaired family
	 embarrassment to wear the appliance
	social discomfort
	 being teased
	 patient anxiety
	 collaboration (compliance) issues or failure to complete treatment, e.g., dropout

*Modified from Preoteasa et al. [3]

Chapter 4 presents the protocol for the cross-sectional study described in chapter 5. The pilot studies in this protocol confirmed the need to address our research questions on reporting or considering adverse effects in abstracts of systematic reviews of orthodontic interventions and the presence of spin in these abstracts.

The results of this cross-sectional study (chapter 5) showed that 77% (75/98) of the included systematic reviews of orthodontic interventions reported or considered (i.e., discussed, weighted, etc.) potential adverse effects in the abstract. This prevalence was lower i.e., 77% (75/98) than what was reported or considered on adverse effects in the main manuscript of these reviews i.e., 85% (83/98) [4] (chapter 3). The CONSORT (Consolidated Standards of Reporting Trials) harms extension [5], PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) harms extension [6], and the MECIR (Methodological Expectations of Cochrane Intervention Reviews) [7] have all incorporated statements that reporting on adverse effects or findings related to adverse effects is mandatory in abstracts when such findings were sought or reported in the main text. A recent overview of reviews analyzed the changes in reporting adverse effects in abstracts of RCTs over time and found that adverse effects were reported in 47% (258/552) of the abstracts of RCTs before the

publication of the CONSORT harms extension and in 54% (643/1201) of the abstracts of the RCTs published after the publication of this statement [8]. However, caution should be applied when comparing the findings of other studies with our results because of possible differences such as; (1) research designs; (2) sample size; (3) the field of research.

Our research showed that spin on adverse effects in abstracts of systematic reviews of orthodontic interventions was present in 41% (40/98) of the included reviews (chapter 5). Other epidemiolocal research on abstracts of systematic reviews, randomized- and nonrandomized studies in different research fields also identified a high prevalence of spin [9, 10, 11, 12, 13, 14, 15, 16]. In the field of orthodontics, spin was found in 49% (53/109) of abstracts of orthodontic meta-analyses [17] and in 62% (69/111) of abstracts of parallelgroup RCTs with clearly stated statistically non-significant primary outcomes [18]. Both studies [17, 18] also assessed the association of predictors with the presence of spin. In our study, no associations were found between the presence of spin in the abstracts and any of the predictors (Predictors were: journal, publication year, number of authors, conflict of interest reported, conflict of interest present, funding reported, and type of orthodontic intervention) (Chapter 5). The study by Guo et al. [18] supported our findings regarding the overlapping predictors i.e., 'the year of publication' and 'the number of authors'. In the study by Makou et al. [17] overlap was found for the predictors 'the year of publication' and 'the journal'. However, Makou et al. [17] reported a higher risk of spin in studies with six or more authors. Several variables should be considered when comparing our results with these previous studies [17, 18]. The main variable to consider is that our study only assessed spin on adverse effects in abstracts of systematic reviews. Other variables that could explain differences are: (1) different types and subtypes of spin and the definitions of spin; (2) the location in the text where spin was assessed; (3) the research design; (4) the field of research; (5) the types of interventions; (6) the journals included; and (7) the time point of publication [5].

Our cross-sectional study further showed that misleading reporting (90% (36/40)) on adverse effects was the predominant category of spin in abstracts of systematic reviews of orthodontic interventions. Nascimento et al. [19] reported similar findings, i.e., misleading reporting was the most common type of spin (73% (48/66)) in abstracts of systematic reviews of physiotherapy interventions for low back pain.

Chapter 6 presents the critical appraisal of a systematic review of fixed orthodontic retainers and periodontal health [20]. This systematic review addressed research questions on the possible adverse effects of fixed orthodontic retainers on periodontal health. According to the AMSTAR-2 assessment [21], this review was classified as 'moderate' quality, and overall low risk of bias was assigned using the ROBIS tool [22, 23]. However, ten additional limitations, not covered by the AMSTAR-2 and ROBIS tools, were identified, i.e., two for the methods and eight for the reporting of the systematic review. The critical appraisal pointed out that no recommendations could be made on the best type of fixed retainer to use in clinical practice because (1) the quality of evidence was low for most of the included studies in this review; (2) five out of the 29 included studied reported on poorer periodontal health around fixed retainers; (3) the additional identified limitations in this critical appraisal could have influenced the results of this systematic review. Chapter 7 presents the critical appraisal of a randomized controlled trial that assessed differences in dental stability, patient perceptions and compliance, and retainer failure after five years of retention, comparing vacuum-formed retainers (VFR) and canine-to canine bonded retainers. According to the Risk of bias 2 tool (RoB2) [24], an overall high risk of bias was assigned, and seven items were not or partially reported according to the CONSORT statement [25]. Additional limitations were identified during the critical appraisal of this RCT. The authors of this RCT [26] reported that most changes were small in both jaws after five years of retention, and only the anterior alignment in the mandible was more stable in the canine-to-canine bonded retainer group. However, this finding is possibly not the result of the retention technique but due to the poor compliance with the VFRs and the inclusion of adolescents only. This critical appraisal showed that the results reported by Krämer et al. [26] should be carefully considered before implementation.

The case report described in chapter 8 presents an adverse effect in orthodontics, which concerns unwanted tooth movements during the retention phase after orthodontic treatment. These movements were probably not the result of dental relapse, but caused by the fixed flexible spiral retainer wire because data at the start of the first orthodontic treatment showed that the original position of the right upper canine differed significantly from the position at the beginning of the second orthodontic treatment phase. This type of unwanted tooth movement combined with a fixed flexible spiral retainer wire has been described previously in orthodontic research in the lower anterior segment [27, 28, 29]. According to Arnold et al., the etiology of these tooth movements consists of forces generated in various ways, i.e., (1) untwisting of the round flexible spiral wires, (2) mechanical deformation of the wire or (3) elastic deflection of the wire [30]. One or more of these generated forces could be the reason for the unwanted tooth movements in this case report. At the completion of the second orthodontic treatment phase, we chose to place fixed braided rectangular retainer wires in the upper and lower jaw since they achieve better torque control [30]. This case report presents a crucial adverse effect in orthodontics. Researchers, patients, and clinicians should be aware of this and recognize this type of unwanted tooth movement early on by organizing regular control visits in the retention phase following orthodontic treatment.

Section 2. Strengths and limitations of this dissertation

This dissertation has the following strengths. First, we developed a new framework for defining the adverse effects of orthodontic interventions (Table 3). Second, pilot studies were conducted to calculate the needed sample size, calibrate researchers, and improve our research questions and methodology (chapters 2 and 4). Third, a protocol for each cross-sectional study was developed and published a priori (chapters 2 and 4), which (1) permitted a careful development and planning, including pilot tests, of various research steps, thereby avoiding potential methodological issues; (2) prevented ad hoc decisions during data extraction; (3) reduced the risk of biases, in particular selective (non) reporting bias; and (4) improved transparent reporting and reproducibility [31, 32]. Fourth, all raw data are available in the additional files or were registered in Open Science Framework. Fifth, the protocol presented in chapter 4 was the first publication on spin on adverse effects in systematic reviews of interventions. Sixth, to achieve accurate, complete, and transparent

reporting, we consulted reporting guidelines for both protocols (PRISMA-P 2015 statement [33, 34]) and both cross-sectional studies (STROBE statement [35], PRISMA 2020 statement [36, 37]) and the case report (CARE-guideline [38, 39]) (chapter 2,3,4,5 and 8).

Limitations in this dissertation can be classified as those related to the methods of our crosssectional studies and those related to the quality of the primary studies included in the eligible systematic reviews of orthodontic interventions. Limitations to the methodology of our cross-sectional studies are: First, the findings reported in the complete orthodontic literature are expected to be worse than those reported in our cross-sectional studies [4, 40] because we only included systematic reviews published in the five leading orthodontic journals and the Cochrane Database of Systematic Reviews. Second, the true magnitude of certain prevalence data (chapters 3 and 5) could be underestimated because we only included recent eligible systematic reviews, i.e., those published between August 1, 2009, and July 21, 2021. However, this period was selected because the inception date coincides with the launch of the PRISMA statement [41, 42]. Third, the assessment of spin is not entirely objective [43]. However, our initial inter-operator agreement for assigning spin was high (Cohen's $\kappa = 0.94$), and complete disagreement was obtained through discussions. Limitations related to the quality of the primary studies included in the eligible systematic reviews of orthodontic interventions are: First, the risk of selective (non) reporting bias because research has shown that adverse effects in both primary research and systematic reviews were poorly assessed and reported [44, 45, 46, 47, 48, 49]. Second, the wide variety of types and definitions of reported adverse effects (chapter 3), made it at times hard to analyze and classify them. This was also reported by Qureshi et al. [50].

Section 3. The meaning of the findings of this dissertation (possible explanations and implications for clinicians and policymakers), reported unanswered questions and future research.

This dissertation provides insights into (1) seeking and reporting of adverse effects in systematic reviews of orthodontic interventions, (2) spin on adverse effects in the abstract of systematic reviews of orthodontic interventions, and (3) classifying and defining adverse effects of orthodontic interventions according to our new framework (Table 3). However, more research is needed to investigate the methodology of how reviewers sought and reported adverse effects. This was confirmed in a recent study, which showed that none of the analyzed systematic reviews and meta-analyses followed any guidelines for assessing adverse effects [45]. Research studies should include mandatory information on adverse effects such as definitions of adverse effects, their duration, and time points for assessing them. Research on developing core outcomes for adverse effects [51] and specific guidelines for defining, assessing, and reporting them in both primary studies and systematic reviews are indicated. In addition, systematic reviews that focus exclusively on adverse effects regardless of the condition for which the intervention was performed are necessary [52, 53]. Research is indicated to investigate whether all adverse effects were indeed sought and reported as initially planned in the registered protocols of the included studies. Discrepancies between what was planned for the assessment of adverse effects in protocols registered in PROSPERO and what was reported on these effects in the final review have been reported [54].

The findings in this dissertation could have policy implications for making judgments on accepting or rejecting a systematic review of orthodontic interventions for publication. For example, editors and peer-reviewers should be instructed to adopt various items on adverse effects defined in the Methodological Expectations of Cochrane Intervention Reviews (MECIR) standards [7] and trained to recognize the presence of spin.

References

- 1. Mahady SE, Schlub T, Bero L, Moher D, Tovey D, George J, Craig JC. Side effects are incompletely reported among systematic reviews in gastroenterology. J Clin Epidemiol. 2015 Feb;68(2):144-53.
- Hopewell S, Wolfenden L, Clarke M. Reporting of adverse events in systematic reviews can be improved: survey results. J Clin Epidemiol. 2008 Jun;61(6):597-602. doi: 10.1016/j.jclinepi.2007.10.005. Epub 2008 Apr 14. PMID: 18411039.
- Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F. (editor). Orthodontics-Basic aspects and clinical considerations. March 9, 2012 under CC BY 3.0 license. www.intechopen.com. [online] Available from: https://cdn.intechopen.com/pdfs/31388/InTech-Risks and complicationsassociated with orthodontic treatment.pdf">https://cdn.intechopen.com/pdfs/31388/InTech-Risks and complicationsassociated with orthodontic treatment.pdf (Accessed 4 December 2021).
- Steegmans PAJ, Di Girolamo N, Bipat S, Reynders RAM. Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1). Syst Rev. 2023 Jul 3;12(1):112. doi: 10.1186/s13643-023-02273-7. PMID: 37400925
- Ioannidis JP, Evans SJ, Gøtzsche PC, O'Neill RT, Altman DG, Schulz K, Moher D; CONSORT Group. Better reporting of harms in randomized trials: an extension of the CONSORT statement. Ann Intern Med. 2004 Nov 16;141(10):781-8.
- Zorzela L, Loke YK, Ioannidis JP, Golder S, Santaguida P, Altman DG, Moher D, Vohra S; PRISMA_Harms Group. PRISMA harms checklist: improving harms reporting in systematic reviews. BMJ. 2016 Feb 1;352:i157. doi: 10.1136/bmj.i157.
- 7. Higgins JPT, Lasserson T, Chandler J, Tovey D, Thomas, J, Flemyng E, Churchill R. Methodological Expectations of Cochrane Intervention Reviews. Cochrane: London, Version February 2022
- Junqueira DR, Phillips R, Zorzela L, Golder S, Loke Y, Moher D, Ioannidis JPA, Vohra S. Time to improve the reporting of harms in randomized controlled trials. J Clin Epidemiol. 2021 Aug;136:216-220. doi: 10.1016/j.jclinepi.2021.04.020. Epub 2021 May 10. PMID: 33984494.
- Boutron I, Dutton S, Ravaud P, Altman DG. Reporting and interpretation of randomized controlled trials with statistically nonsignificant results for primary outcomes. JAMA. 2010 May 26;303(20):2058-64. doi: 10.1001/jama.2010.651.
- Jones C, Rulon Z, Arthur W, Ottwell R, Checketts J, Detweiler B, Calder M, Adil A, Hartwell M, Wright DN, Vassar M. Evaluation of spin in the abstracts of systematic reviews and meta-analyses related to the treatment of proximal humerus fractures. J Shoulder Elbow Surg. 2021 Jan 19:S1058-2746(21)00016-1. doi: 10.1016/j.jse.2020.11.026. Epub ahead of print. PMID: 33482369.
- Khan MS, Lateef N, Siddiqi TJ, Rehman KA, Alnaimat S, Khan SU, Riaz H, Murad MH, Mandrola J, Doukky R, Krasuski RA. Level and Prevalence of Spin in Published Cardiovascular Randomized Clinical Trial Reports With Statistically Nonsignificant Primary Outcomes: A Systematic Review. JAMA Netw Open. 2019 May 3;2(5):e192622. doi: 10.1001/jamanetworkopen.2019.2622. PMID: 31050775; PMCID: PMC6503494.
- Lazarus C, Haneef R, Ravaud P, Boutron I. Classification and prevalence of spin in abstracts of nonrandomized studies evaluating an intervention. <u>BMC Med Res Methodol.</u> 2015 Oct 13;15:85. doi: 10.1186/s12874-015-0079-x.
- Mathieu S, Giraudeau B, Soubrier M, Ravaud P. Misleading abstract conclusions in randomized controlled trials in rheumatology: comparison of the abstract conclusions and the results section. Joint Bone Spine. 2012 May;79(3):262-7. doi: 10.1016/j.jbspin.2011.05.008. Epub 2011 Jul 5. PMID: 21733728.
- Reddy AK, Shepard S, Ottwell R, Thompson J, Price CM, Arthur W, Hanson C, Ebert A, Wright DN, Hartwell M, Vassar M. Over 30% of Systematic Reviews and Meta-analyses Focused on Rotator Cuff Tear Treatments Contained Spin in the Abstract. Arthroscopy. 2021 Sep;37(9):2953-2959. doi: 10.1016/j.arthro.2021.03.066. Epub 2021 Apr 19.PMID: 33887409
- Wang D, Chen L, Wang L, Hua F, Li J, Li Y, Zhang Y, Fan H, Li W, Clarke M. Abstracts for reports of randomised trials of COVID-19 interventions had low quality and high spin. J Clin Epidemiol. 2021 Jul 2;139:107-120. doi: 10.1016/j.jclinepi.2021.06.027. Online ahead of print.PMID: 34224834
- 16. Wise A, Mannem D, Arthur W, Ottwell R, Greiner B, Srouji D, Wildes D, Hartwell M, Wright DN, Khojasteh J, Vassar M. Spin within systematic review abstracts on antiplatelet therapies after acute

coronary syndrome: a cross-sectional study. BMJ Open. 2022 Aug 2;12(8):e049421. doi: 10.1136/bmjopen-2021-049421.PMID: 35918107

- Makou O, Eliades T, Koletsi D. Reporting, interpretation, and extrapolation issues (SPIN) in abstracts of orthodontic meta-analyses published from 2000 to 2020. Eur J Orthod. 2021 Mar 19:567-575. doi: 10.1093/ejo/cjab009. Epub ahead of print. PMID: 33740054.
- Guo F, Fang X, Li C, Qin D, Hua F, He H. The presence and characteristics of 'spin' among randomized controlled trial abstracts in orthodontics. Eur J Orthod. 2021 Oct 4;43(5):576-582. doi: 10.1093/ejo/cjab044. PMID: 34397084.
- Nascimento DP, Gonzalez GZ, Araujo AC, Moseley AM, Maher CG, Costa LOP. Eight in Every 10 Abstracts of Low Back Pain Systematic Reviews Presented Spin and Inconsistencies With the Full Text: An Analysis of 66 Systematic Reviews. J Orthop Sports Phys Ther. 2020 Jan;50(1):17-23. doi: 10.2519/jospt.2020.8962. Epub 2019 Aug 23.PMID: 31443622
- Arn ML, Dritsas K, Pandis N, Kloukos D. The effects of fixed orthodontic retainers on periodontal health: A systematic review. Am J Orthod Dentofacial Orthop. 2020;157(2):156–164.e17. doi:10.1016/j.ajodo.2019.10.010
- 21. Shea B J, Reeves B C, Wells G et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non randomised studies of healthcare interventions, or both. BMJ 2017; 358: DOI:10.1136/bmj.j4008.
- University of Bristol. Risk of bias in systematic reviews (ROBIS). ROBIS tool and the ROBIS guidance document. Available at http://www.bristol.ac.uk/social-community- medicine/projects/robis/ (Accessed February 2020)
- Whiting P, Savović J, Higgins JP, Caldwell DM, Reeves BC, Shea B, Davies P, Kleijnen J, Churchill R; ROBIS group. ROBIS: A new tool to assess risk of bias in systematic reviews was developed. J Clin Epidemiol. 2016 Jan;69:225-34.
- 24. Sterne J A C, Savović J, Page M J et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ 2019; DOI: 10.1136/bmj.l4898.
- Schulz K F, Altman D G, Moher D, CONSORT Group. CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. Ann Intern Med 2010; 152: 726–732.
- Krämer A, Sjöström M, Apelthun C, Hallman M, Feldmann I. Post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers-a randomized controlled trial. Eur J Orthod 2022; DOI: 10.1093/ejo/cjac043.
- Renkema, A.M.; Renkema, A.; Bronkhorst, E.; Katsaros, C. Long-term effectiveness of canine-to-canine bonded flexible spiral wire lingual retainers. Am J Orthod Dentofacial Orthop. 2011, 139(5), 614– 621. doi:10.1016/j.ajodo.2009.06.04
- Katsaros, C.; Livas, C.; Renkema, A.M. Unexpected complications of bonded mandibular lingual retainers. Am J Orthod Dentofacial Orthop. 2007, 132(6), 838-41. doi:10.1016/j.ajodo.2007.07.011.PMID: 18068606
- 29. Singh, P.; Canine avulsion: An extreme complication of a fixed mandibular lingual retainer. Am J Orthod Dentofacial Orthop. 2021, 160(3), 473–477. doi:10.1016/j.ajodo.2021.04.013
- Arnold, D.T.; Dalstra, M.; Verna, C. Torque resistance of different stainless steel wires commonly used for fixed retainers in orthodontics. J Orthod. 2016, 43(2), 121-9. doi: 10.1080/14653125.2016.1155814. Epub 2016 Apr 22.PMID: 27104351
- Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A Deceased, Petticrew M, Shekelle P, Stewart LA; the PRISMA-P Group: Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;349:g7647.
- 32. Lasserson TJ, Thomas J, Higgins JPT. Chapter 1: Starting a review. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.3 (updated February 2022). Cochrane, 2022. Available from <u>www.training.cochrane.org/handbook</u> (Accessed 9 August 2023)
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. Preferred reporting items for systematic review and meta- analysis protocols (PRISMA-P) 2015 statement. Syst Rev. 2015;4(1):1.
- 34. Shamseer L, Moher D, Clarke M, Ghersi D, Deceased LA, Petticrew M, Shekelle P, Stewart LA, the PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;349:g7647.

- 35. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE<u>Initiative</u>. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. <u>Ann Intern Med.</u> 2007 Oct 16;147(8):573-7.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021 Mar 29;372:n71. doi: 10.1136/bmj.n71. PMID: 33782057.
- Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, McKenzie JE. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. BMJ. 2021 Mar 29;372:n160. doi: 10.1136/bmj.n160. PMID: 33781993.
- Gagnier JJ, Riley D, Altman DG, Moher D, Sox H, Kienle G, for the Care Group. The CARE Guidelines: Consensus-Based Clinical Case Reporting Guideline Development. Dtsch Arztebl Int. 2013 Sep; 110(37): 603-608. Published online 2013 Sep 13. Doi: 10.3238/arztebl.2013.0603. PMCID: PMC3784031
- Riley, DS, Barber MS, Kienle GS, Aronson JK, von Schoen-Angerer T, Tugwell P, Kiene H, Helfand M, Altman DG, Sox H, Werthmann PG, Moher D, Rison RA, Shamseer L, Koch CA, Sun GH, Hanaway P, Sudak NL, Kaszkin-Bettag M, Carpenter JE, Gagnier JJ. CARE guidelines for case reports: explanation and elaboration document. J Clin Epidemiol. 2017. pii: S0895-4356(17)30037-9. PMID: 28529185
- Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin on adverse effects in abstracts of systematic reviews of orthodontic interventions: a cross-sectional study (part 2). Syst Rev. 2023 Jun 20;12(1):99. doi: 10.1186/s13643-023-02269-3.PMID: 37340504
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. PLoS Med. 2009; 6: e1000100. Epub 2009 Jul 21.
- 42. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med. 2009;6:e1000097.
- Chiu K, Grundy Q, Bero L. 'Spin' in published biomedical literature: A methodological systematic review. PLoS Biol. 2017 Sep 11;15(9):e2002173. doi: 10.1371/journal.pbio.2002173. PMID: 28892482; PMCID: PMC5593172.
- Zorzela L, Golder S, Liu Y, Pilkington K, Hartling L, Joffe A, Loke Y, Vohra S. Quality of reporting in systematic reviews of adverse events: systematic review. BMJ. 2014 Jan 8;348:f7668. doi: 10.1136/bmj.f7668. PMID: 24401468; PMCID: PMC3898583.
- 45. Qureshi R, Mayo-Wilson E, Rittiphairoj T, McAdams-DeMarco M, Guallar E, Li T. Summaries of harms in systematic reviews are unreliable Paper 2: Methods used to assess harms are neglected in systematic reviews of gabapentin, Journal of Clinical Epidemiology (2021), doi: <u>https://doi.org/10.1016/j.jclinepi.2021.10.024</u>
- Golder S, Loke YK, Wright K, Norman G. Reporting of Adverse Events in Published and Unpublished Studies of Health Care Interventions: A Systematic Review. PLoS Med. 2016 Sep 20;13(9):e1002127. doi: 10.1371/journal.pmed.1002127. eCollection 2016 Sep.
- Golder S, Loke YK, Wright K, Sterrantino C. Most systematic reviews of adverse effects did not include unpublished data. J Clin Epidemiol. 2016 Sep;77:125-133. doi: 10.1016/j.jclinepi.2016.05.003. Epub 2016 Jun 1. PMID: 27259470.
- Mayo-Wilson E, Fusco N, Li T, Hong H, Canner JK, Dickersin K; MUDS investigators. Harms are assessed inconsistently and reported inadequately part 1: systematic adverse events. J Clin Epidemiol. 2019 Sep;113:20-27. doi: 10.1016/j.jclinepi.2019.04.022. Epub 2019 May 2. PMID: 31055175.
- Mayo-Wilson E, Fusco N, Li T, Hong H, Canner JK, Dickersin K; MUDS investigators. Harms are assessed inconsistently and reported inadequately Part 2: nonsystematic adverse events. J Clin Epidemiol. 2019 Sep;113:11-19. doi: 10.1016/j.jclinepi.2019.04.020. Epub 2019 May 2. PMID: 31055176.
- Qureshi R, Mayo-Wilson E, Li T, Summaries of harms in systematic reviews are unreliable Paper 1: An introduction to research on harms, *Journal of Clinical Epidemiology* (2021), doi: https://doi.org/10.1016/j.jclinepi.2021.10.023
- 51. Tsichlaki A, O'Brien K, Benson PE, Marshman Z, Johal A, Colonio-Salazar FB, Harman NL, Fleming PS. Development of a core outcome set for use in routine orthodontic clinical trials. Am J Orthod

Dentofacial Orthop. 2020 Nov;158(5):650-660. doi: 10.1016/j.ajodo.2020.05.010. Epub 2020 Sep 16. PMID: 32950336.

- Peryer G, Golder S, Junqueira D, Vohra S, Loke YK. Chapter 19: Adverse effects. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.3 (updated February 2022). Cochrane, 2022. Available from <u>www.training.cochrane.org/handbook</u>. (Accessed 9 August 2023)
- Mayo-Wilson E, Qureshi R, Li T. Conducting separate reviews of benefits and harms could improve systematic reviews and meta-analyses. Syst Rev. 2023 Apr 15;12(1):67. doi: 10.1186/s13643-023-02234-0.PMID: 37061724
- Parsons R, Golder S, Watt I. More than one-third of systematic reviews did not fully report the adverse events outcome. J Clin Epidemiol. 2019 Apr;108:95-101. doi: 10.1016/j.jclinepi.2018.12.007. Epub 2018 Dec 13. PMID: 30553831.



CHAPTER 10

Summary and conclusions

Chapter 1 introduces the topic of this dissertation. This chapter also presents the objectives, and the definitions used in this dissertation.

Chapters 2, 3, 4, and 5, present the cross-sectional studies on defining, seeking and reporting on adverse effects in systematic reviews of orthodontic interventions. In part 1 of these studies, we assessed defining, seeking and reporting of adverse effects in these reviews. Part 2 focused on the reporting of adverse effects in abstracts of systematic reviews of orthodontic interventions and the presence and type of spin in these abstracts regarding adverse effects. The Cochrane Database of Systematic Reviews and five leading orthodontic journals were searched for eligible reviews published between August 1, 2009, and July 31, 2021. Systematic reviews of orthodontic interventions on human patients of any health status, sex, age, demographics, and socio-economic status and any adverse effect scored at any endpoint or timing were eligible, resulting in the inclusion of 98 systematic reviews.

All study selection and data extraction procedures were conducted by two researchers independently. The results showed that in 84 of 98 (86%) reviews, findings related to adverse effects of orthodontic interventions were sought, and in 83 of 98 (85%) reviews, reviewers reported on them. More than 90% of the reviews considered or discussed potential adverse effects. However, only 36% (35/98) of the reviews defined seeking of adverse effects as a research objective. The Journal of Orthodontics and Craniofacial research had approximately seven times the odds (OR 7.2, 95% CI 1.1 to 48.0) to report that adverse effects were sought in the research objectives compared to the eligible reviews in the Cochrane Database of Systematic Reviews. All identified adverse effects (N=195) were divided into 12 categories according to our new framework for defining adverse effects of orthodontic interventions. Five of these categories accounted for 83% (162/195) of all adverse effects sought and reported. These 5 categories are related to (1) tooth structures, (2) periodontal tissues, (3) undesired treatment results, (4) relapse and stability, and (5) negative qualitative experiences by the patient or carer(s) (Table 8 in chapter 3). Based on the findings of part 1 of our cross-sectional studies, we concluded that most of the included systematic reviews sought and reported adverse effects of orthodontic interventions. However, we have to be cautious when implementing these findings in clinical practice, because of several limitations, i.e., this sample does not represent the complete orthodontic literature, the risk of selective (non) reporting bias regarding adverse effects in the primary studies that were included in the reviews, and unequal assessments of different types of adverse effects, with certain effects being frequently assessed while others rarely or never. (Table 8 in chapter 3).

In part 2 of our cross-sectional studies, we investigated whether adverse effects in systematic reviews of orthodontic interventions were reported or considered in the abstract and whether spin on adverse effects was identified and what type of spin. We also compared the reported adverse effects in the abstract to those sought and reported in the main texts of these reviews. Associations between the presence of spin in the abstracts and variables like the journal, year, number of authors, conflict of interest reported, funding, and type of orthodontic intervention were also assessed. The results showed that 77% (75/98) of the included reviews reported or considered (i.e., discussed, weighed, etc.) potential adverse effects of orthodontic interventions in the abstract. In 41% (40/98) of these reviews, spin regarding adverse effects was present in the abstracts. Misleading reporting was the most

frequent type of spin (90% (36/40). Our univariable logistic regression models (95% CI) showed no associations between the presence of spin in the abstracts of systematic reviews of orthodontic interventions and any of the variables explored. The results of part 2 of our cross-sectional studies showed that findings related to adverse effects reported in the abstract should be interpreted with caution, because abstracts with spin regarding adverse effects, i.e., not being reported or misleading reporting, can make an intervention appear more favorable than it is. End-users should be aware of the presence of spin in abstracts of systematic reviews of orthodontic interventions. Editors and peer-reviewers have an essential role in tackling these problems before systematic reviews of orthodontic intervention.

Chapter 6 presents the critical appraisal of the systematic review by Arn et al. [1]. The authors assessed the potentially adverse effects of fixed retainers on periodontal health and compared different retainers according to their effects on periodontal health. The results pointed out that fixed orthodontic retainers appear compatible with periodontal health or at least unrelated to severe detrimental effects on the periodontium. However, it has to be considered that 5 out of the 29 included studies showed poorer periodontal health around fixed retainers and the majority of the included evidence was of low quality. In addition, the critical appraisal presented various limitations regarding the methods and reporting of the systematic review.

Chapter 7 presents, the critical appraisal of a randomized controlled trial by Krämer et al. [2]. The authors compared vacuum-formed retainers (VFR) to canine-to-canine bonded retainers in adolescents after five years of retention and assessed differences in dental stability, patient perceptions, compliance and retainer failures. The results showed that for both retention protocols, most post-treatment changes were small in both jaws, but the anterior alignment in the mandible was more stable in the bonded retainer group. This difference is possibly not related to the retention protocol but to the poor compliance identified in the group treated with the VFR and the fact that only adolescents were included. Additional limitations and the high risk of bias in the results of this randomized controlled trial should be considered before implementing its reported findings.

Chapter 8 presents a clinical case that showed unwanted tooth movements during the retention phase following orthodontic treatment. This adverse effect was probably related to the undesired forces of the fixed flexible spiral retainer wire in the upper jaw. After orthodontic retreatment, fixed braided rectangular retainer wires were placed in combination with a vacuum-formed retainer in the upper jaw. The knowledge and awareness of this adverse effect is important for researchers, patients, orthodontists, and dentists. This case report shows the importance of regular control visits after orthodontic treatment.

In chapter 9 the results of this dissertation are discussed. Section 1 presents the principal findings of this dissertation and how they compare with other studies. Section 2 presents the strengths and limitations of this dissertation, and Section 3 presents the meaning of the findings of this dissertation, reported unanswered questions, and future research.

Conclusions

Based on the answers to the 12 research questions that were assessed in this dissertation, the following conclusions were made (see 'highlights' page 7):

This dissertation presents a new framework for categorizing and defining adverse effects of orthodontic interventions. Relapse and stability issues and undesired treatment results were the predominant adverse effects sought and reported in systematic reviews of orthodontic interventions, but many adverse effects were underassessed and underreported. This dissertation showed that assessing and reporting of adverse effects of orthodontic interventions in systematic reviews is often not systematic, incomplete, and selective. This dissertation also showed that spin on adverse effects in abstracts of systematic reviews of orthodontic interventions was highly prevalent and that de predominant type of spin was misleading reporting. These findings implicate that what is reported on adverse effects in systematic reviews of orthodontic interventions should be interpreted with caution e.g., when making clinical decisions. Future research should focus on developing, assessing, and reporting of core adverse effects of orthodontic interventions in primary studies and in systematic reviews. Besides conducting traditional systematic reviews of interventions, it is also indicated to conduct systematic reviews that focus exclusively on adverse effects. Multiple stakeholders such as researchers, editors, peer reviewers, and policy makers have to take responsibility and clinicians and patients will ultimately benefit.

References

- Arn ML, Dritsas K, Pandis N, Kloukos D. The effects of fixed orthodontic retainers on periodontal health: A systematic review. Am J Orthod Dentofacial Orthop. 2020;157(2):156–164.e17. doi:10.1016/j.ajodo.2019.10.010
- Krämer A, Sjöström M, Apelthun C, Hallman M, Feldmann I. Post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers-a randomized controlled trial. Eur J Orthod 2022; DOI: 10.1093/ejo/cjac043.



CHAPTER 11

Samenvatting en conclusies

Hoofdstuk 1 introduceert het onderwerp van dit proefschrift. In dit hoofdstuk zijn ook de doelstellingen en de in dit proefschrift gebruikte definities opgenomen.

De hoofdstukken 2, 3, 4 en 5 presenteren de *cross-sectional studies* met betrekking tot het definiëren, zoeken en rapporteren van *adverse effects* in *systematic reviews* van orthodontische interventies. In deel 1 van deze studies beoordeelden we het definiëren, zoeken en rapporteren van *adverse effects* in deze *reviews*. Deel 2 richtte zich op het rapporteren van *adverse effects* in samenvattingen van *systematic reviews* van orthodontische interventies en de aanwezigheid en het type *spin* in de samenvatting ten aanzien van deze *adverse effects*. De '*Cochrane Database of Systematic Reviews*' en vijf toonaangevende orthodontische tijdschriften werden doorzocht op in aanmerking komende *reviews* gepubliceerd tussen 1 augustus 2009 en 31 juli 2021. *Systematic reviews* van orthodontische interventies bij patiënten van elke gezondheidsstatus, geslacht, leeftijd, demografie en sociaaleconomische status en elk nadelig effect waargenomen op elk eindpunt of tijdstip kwamen in aanmerking, hetgeen resulteerde in de inclusie van 98 *systematic reviews*.

Alle studie selectie en data-extractie procedures zijn door twee onderzoekers onafhankelijk van elkaar uitgevoerd. De resultaten lieten zien dat in 84 van de 98 (86%) reviews werd gezocht naar bevindingen met betrekking tot adverse effects van orthodontische interventies en dat in 83 van de 98 (85%) reviews de reviewers daarover rapporteerden. Meer dan 90% van de reviews beschouwde of bediscussieerde potentiële adverse effects in de reviews. Echter, slechts 36% (35/98) van de reviews definieerde het zoeken van adverse effects als een onderzoeksdoelstelling. In het tijdschrift 'The Journal of Orthodontics and Craniofacial research' was de kans ongeveer zeven keer zo groot (OR 7.2, 95% Cl 1.1 to 48.0) dat de gezochte adverse effects werden gerapporteerd in de onderzoeksdoelstellingen in vergelijking tot de in aanmerking komende reviews in de 'Cochrane Database of Systematic Reviews'. Alle geïdentificeerde adverse effects (N=195) werden ingedeeld in 12 categorieën volgens ons nieuwe framework voor het definiëren van adverse effects van orthodontische interventies. Drieëntachtig procent (162/195) van alle gezochte en gerapporteerde adverse effects vielen onder vijf van deze categorieën. Deze 5 categorieën hebben betrekking op (1) tandstructuren, (2) parodontale weefsels, (3) ongewenste behandelresultaten, (4) terugval en stabiliteit en (5) negatieve kwalitatieve ervaringen van de patiënt of verzorger(s) (tabel 8 in hoofdstuk 3). Op basis van de resultaten van deel 1 van onze cross-sectional studies, concludeerden wij dat de meeste van de geïncludeerde systematic reviews, adverse effects van orthodontische interventies zochten en rapporteerden. We moeten echter voorzichtig zijn met het interpreteren van deze data vanwege verschillende beperkingen: 1) deze studies vertegenwoordigen niet de volledige orthodontische literatuur, 2) het risico op selective (non) reporting bias in de primaire studies die in de reviews zijn opgenomen en 3) ongelijke beoordeling van verschillende adverse effects, waarbij bepaalde adverse effects vaak werden beoordeeld en andere zelden of nooit (tabel 8 in hoofdstuk 3).

In deel 2 van onze *cross-sectional studies* onderzochten wij of in *systematic reviews* van orthodontische interventies *adverse effects* werden gerapporteerd of beschouwd in de samenvatting en of er sprake was van *spin* met betrekking tot de onderzochte *adverse effects* en welk type *spin*. De in de samenvatting gerapporteerde *adverse effects* hebben we ook vergeleken met de gezochte en gerapporteerde *adverse effects* in de hoofdteksten van

deze reviews. Verbanden tussen de aanwezigheid van spin in de samenvattingen en variabelen zoals het tijdschrift, jaartal, aantal auteurs, gerapporteerde belangenverstrengeling, financiering en type orthodontische interventie zijn ook beoordeeld. Uit de resultaten bleek dat 77% (75/98) van de geïncludeerde reviews in de samenvatting mogelijke adverse effects van orthodontische interventies rapporteerden of beschouwden (d.w.z. bediscussieerde, afwoog, enz.). In 41% (40/98) van deze reviews was er sprake van spin met betrekking tot adverse effects in de samenvatting. Misleading reporting was het meest voorkomende type spin (90% (36/40)). Onze univariabele logistische regressiemodellen (95% CI) lieten geen verbanden zien tussen de aanwezigheid van spin in de samenvattingen van systematic reviews van orthodontische interventies en de onderzochte variabelen. De resultaten van deel 2 van onze cross-sectional studies toonden aan dat resultaten met betrekking tot adverse effects gerapporteerd in de samenvatting voorzichtig moeten worden geïnterpreteerd, omdat samenvattingen met spin ten aanzien van adverse effects, d.w.z. niet gerapporteerd of misleidend gerapporteerd, een interventie gunstiger kunnen doen lijken dan deze is. Eindgebruikers moeten zich bewust zijn van de aanwezigheid van spin in samenvattingen van systematic reviews van orthodontische interventies. Redacteuren en peer-reviewers spelen een essentiële rol bij het aanpakken van deze problemen alvorens systematic reviews van orthodontische interventies worden geaccepteerd voor publicatie.

Hoofdstuk 6 presenteert de kritische beoordeling van het *systematic review* van Arn et al. [1]. De auteurs vergeleken de mogelijke *adverse effects* van verschillende vaste retentiedraden op de parodontale gezondheid. De resultaten wezen erop dat vaste retentiedraden compatibel lijken met de parodontale gezondheid of in ieder geval geen verband houden met ernstige schadelijke effecten op het parodontium. Er moet echter rekening worden gehouden met het gegeven dat 5 van de 29 geïncludeerde studies een slechtere parodontale gezondheid lieten zien rondom vaste retentiedraden en dat het merendeel van het geïncludeerde *evidence* van lage kwaliteit was. Daarnaast liet de kritische beoordeling verschillende beperkingen zien met betrekking tot de methoden en de rapportage van het *systematic review*.

Hoofdstuk 7 presenteert de kritische beoordeling van een *RCT* van Krämer et al. [2]. De auteurs vergeleken vacuümgevormde retentie apparatuur met vaste retentiedraden van hoektand tot hoektand bij adolescenten na vijf jaar retentie en beoordeelden verschillen in gebitsstabiliteit, patiëntpercepties, therapietrouw en het falen van de retentie apparatuur. De resultaten lieten zien dat voor beide retentie protocollen de meeste veranderingen na de behandeling klein waren in beide kaken, maar dat het orthodontisch behandelde (opgelijnde) onderfront stabieler was in de groep met vaste retentiedraden. Dit verschil houdt mogelijk geen verband met het retentie protocol, maar met de slechte naleving van de therapie geïdentificeerd in de groep die behandeld werd met de vacuümgevormde retentie apparatuur en het feit dat alleen adolescenten werden geïncludeerd. Aanvullende beperkingen en het hoge risico op bias in de resultaten van deze *RCT* moeten in acht genomen worden alvorens de gerapporteerde resultaten worden geïmplementeerd.

Hoofdstuk 8 presenteert een klinische casus die ongewenste tandbewegingen liet zien tijdens de retentiefase na een orthodontische behandeling. Dit *adverse effect* hield waarschijnlijk verband met de ongewenste krachten van de vaste flexibele spiraalvormige

retentiedraad in de bovenkaak. Na orthodontische herbehandeling werden vaste gevlochten rechthoekige retentiedraden geplaatst in combinatie met vacuümgevormde retentie apparatuur in de bovenkaak. De kennis en het bewustzijn van dit *adverse effect* is belangrijk voor onderzoekers, patiënten, orthodontisten en tandartsen. Dit *case report* laat het belang zien van regelmatige controlebezoeken na een orthodontische behandeling.

In hoofdstuk 9 worden de resultaten van dit proefschrift besproken. Deel 1 presenteert de belangrijkste bevindingen van dit proefschrift en hoe deze zich verhouden tot andere studies. Deel 2 presenteert de sterke punten en beperkingen van dit proefschrift en deel 3 presenteert de betekenis van de bevindingen van dit proefschrift, gerapporteerde onbeantwoorde vragen en toekomstig onderzoek.

Conclusies

De volgende conclusies werden getrokken op basis van de antwoorden op de 12 onderzoeksvragen in dit proefschrift (zie 'highlights' pagina 7): Dit proefschrift presenteert een nieuw framework voor het categoriseren en definiëren van adverse effects van orthodontische interventies. Relapse, problemen met stabiliteit en ongewenste behandelresultaten waren de belangrijkste adverse effects die werden gezocht en gerapporteerd in systematic reviews van orthodontische interventies, maar veel adverse effects werden onvoldoende beoordeeld en onvoldoende gerapporteerd. Dit proefschrift toonde aan dat het beoordelen en rapporteren van adverse effects van orthodontische interventies in systematic reviews vaak onvolledig was, niet systematisch en selectief. Dit proefschrift toonde ook aan dat spin met betrekking tot adverse effects in samenvattingen van systematic reviews van orthodontische interventies veel voorkwam en dat het meest prevalente type spin misleading reporting was. Deze bevindingen impliceren dat wat gerapporteerd wordt over adverse effects in systematic reviews van orthodontische interventies voorzichtig geïnterpreteerd moet worden, bijvoorbeeld met het maken klinische beslissingen. Toekomstig onderzoek zou zich moeten richten op het ontwikkelen, beoordelen en rapporteren van 'core-adverse effects' van orthodontische interventies in primaire studies en in systematic reviews. Naast het uitvoeren van traditionele systematic reviews van interventies, is het ook geïndiceerd om systematic reviews die zich uitsluitend richten op adverse effects uit te voeren. Meerdere belanghebbenden zoals onderzoekers, redacteuren, peer reviewers en beleidsbepalers moeten hun verantwoordelijkheid nemen en uiteindelijk zullen clinici en patiënten hiervan profiteren.

Referenties

- Arn ML, Dritsas K, Pandis N, Kloukos D. The effects of fixed orthodontic retainers on periodontal health: A systematic review. Am J Orthod Dentofacial Orthop. 2020;157(2):156–164.e17. doi:10.1016/j.ajodo.2019.10.010
- Krämer A, Sjöström M, Apelthun C, Hallman M, Feldmann I. Post-treatment stability after 5 years of retention with vacuum-formed and bonded retainers-a randomized controlled trial. Eur J Orthod 2022; DOI: 10.1093/ejo/cjac043.



CHAPTER 12

List of publications

Peer-reviewed full-text publications

In this dissertation

Steegmans PAJ, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study. Syst Rev. 2019 Apr 5;8(1):89.

Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews. Res Integr Peer Rev. 2019 Dec 19;4:27.

Steegmans PAJ, Di Girolamo N, Bipat S, Reynders RAM. Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1). Syst Rev. 2023 Jul 3;12(1):112.

Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA. Spin on adverse effects in abstracts of systematic reviews of orthodontic interventions: a cross-sectional study (part 2). Syst Rev. 2023 Jun 20;12(1):99.

Steegmans PAJ, Meursinge Reynders RA. Fixed orthodontic retainers and periodontal health. Evid Based Dent. 2020 Dec;21(4):146-149.

Steegmans PAJ, Cavagnetto D, Meursinge Reynders RA. Which orthodontic retention protocol should I implement? A critical assessment of a randomized controlled trial. Evid Based Dent. 2022 Dec;23(4):162-165.

Steegmans PAJ, Jonkman REG, de Lange J. Fixed Flexible Spiral Wire Retainers and Unwanted Tooth movements: A Case Report. *Appl. Sci.* 2023, *13*(2), 922.

List of contributing authors

R.A. Meursinge Reynders (RMR)

Department of Oral- and maxillofacial surgery, Amsterdam University Medical Center (Amsterdam UMC), Amsterdam, the Netherlands | Studio di ortodonzia, Milan, Italy.

S. Bipat (SB) Department of radiology, Amsterdam University Medical Center (Amsterdam UMC), Amsterdam, the Netherlands

N. Di Girolamo (NDG) Department of Clinical Sciences, Cornell University, Ithaca, NY, USA |EBMVet, Cremona, Italy

D. Cavagnetto (DC) Department of Oral- and Maxillofacial surgery, Amsterdam University Medical Centre (Amsterdam UMC), Amsterdam, the Netherlands | Studio di ortodonzia, Milan, Italy

R.E.G. Jonkman (RJ) Department of orthodontics, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam, The Netherlands

J. de Lange (JL)

Department of Oral- and Maxillofacial surgery, Amsterdam University Medical Centre (Amsterdam UMC), Amsterdam, the Netherlands | Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam, Amsterdam, The Netherlands

Author's contributions

Chapter 2

Published as: Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study.

Authors: Steegmans PAJ, Bipat S, Meursinge Reynders RA

Published in: Systematic Reviews, 2019

Author contributions: Conceived and designed: RMR, PS Performed the study: RMR, PS Analyzed the data: RMR, SB, PS Critically revised the manuscript: RMR, SB, PS

Chapter 3

Published as: Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1)

Authors: Steegmans PAJ, Di Girolamo N, Bipat S, Meursinge Reynders RA

Published in: Systematic reviews, 2023

Author contributions: Conceived and designed: RMR, PS Performed the study: RMR, PS Analyzed the data: RMR, NDG, SB, PS Critically revised the manuscript: RMR, NDG, SB, PS

Chapter 4

Published as:

Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews.

Authors: Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA

Published in: Research Integrity and Peer Review, 2019

170

Author contributions: Conceived and designed: RMR, PS Performed the study: RMR, PS Analyzed the data: RMR, NDG, PS Critically revised the manuscript: RMR, NDG, PS

Chapter 5

Published as: Spin on adverse effects in abstracts of systematic reviews of orthodontic interventions: a cross-sectional study (part 2)

Authors: Steegmans PAJ, Di Girolamo N, Meursinge Reynders RA

Published in: Systematic reviews, 2023

Author contributions: Conceived and designed: RMR, PS Performed the study: RMR, PS Analyzed the data: RMR, NDG, PS Critically revised the manuscript: RMR, NDG, PS

Chapter 6

Published as: Fixed orthodontic retainers and periodontal health.

Authors: Steegmans PAJ, Meursinge Reynders RA.

Published in: Evidence Based Dentistry, 2020

Author contributions: Conceived and designed: RMR, PS Performed the study: RMR, PS Analyzed the data: RMR, PS Critically revised the manuscript: RMR, PS

Chapter 7

Published as: Which orthodontic retention protocol should I implement? A critical assessment of a randomized controlled trial

Authors: Steegmans PAJ, Cavagnetto D, Meursinge Reynders RA.

Published in: Evidence-Based Dentistry , 2022

Author contributions: Conceived and designed: DC, RMR, PS Performed the study: DC, RMR, PS Analyzed the data: DC, RMR, PS Critically revised the manuscript: DC, RMR, PS

Chapter 8

Published as: Fixed flexible spiral wire retainers and unwanted tooth movements: a Case Report

Authors: JL, RJ, PS

Published in: Applied Sciences

Author contributions: Conceived and designed: PS Performed the study: PS Analyzed the data: JL, RJ, PS Critically revised the manuscript: JL, RJ, PS

Funding

All expenses for conducting the protocols and research studies in this dissertation were paid evenly by each contributing author.

Conflict of interest

RMR and SB are both Associate Editors for Systematic Reviews. All authors declare that they have no further competing interests.

Data sharing statement

All data are presented in the additional files and raw data have been placed in the Open Science Framework (<u>https://osf.io/ka7mp/</u>)

Reprints and permission

All publishing companies of the published articles in this dissertation granted permission to reproduce these articles in this dissertation and its additional files. Pertinent URL's on this information are listed under here.

BioMed Central [online] Available from: <u>https://www.biomedcentral.com/about/policies/reprints-and-permissions</u> (accessed July 5th 2023)

Springer Nature [online] Available from: <u>https://www.nature.com/nature-portfolio/reprints-and-permissions</u> (accessed July 5th 2023

MDPI [online] Available from: <u>https://www.mdpi.com/authors/rights</u> (accessed July 5th 2023)

12

Acknowledgment

First of all, I would like to express my gratitude for the opportunity that was given to me to complete this PhD.

I would like to thank my esteemed supervisors dr. Reint Meursinge Reynders, Prof. dr. Jan de Lange and dr. Ronald Jonkman for their guidance, supervision, and invaluable support during the course of my PhD. Especially Reint, thank you for all the time and effort you put into this PhD process with me. I am very grateful for all the knowledge you have given me throughout the last 5 years. Without your passionate guidance, I could not have completed this PhD. I would also like to thank all participating researchers, dr. S. Bipat, prof. dr. N di Girolamo and Davide Cavagnetto for their contribution to my PhD.

Additionally, I would like to thank the members of the PhD committee, prof. dr. A.G. Becking, prof. dr. F. Abbas, prof. dr. L. Dubois, prof. dr. F.R. Rozema, dr. F.N. van der Weijden and dr. M.A. Jongsma for critically reading this dissertation and for being part of the PhD committee.

Furthermore, I would like to express gratitude to my family, my beloved father who I, unfortunately, have to miss since 2010, my dear mother who is always there for me, and my brother who supported me in different ways and today as my paranymph. Dear Walter, I also want to thank you for your support and for being a part of our family. In addition, I would like to express my gratitude to Paul Kerckhoffs, my uncle, who is like a second father to me and has supported me in many ways since I was young. Dear Floor, many thanks for being my paranymph and for your support in many ways over the last 15 years. My appreciation also goes out to my friends and colleagues; Lieke, Juliëtte, Emilie, Anne-Claire, Kiek, Marrit, Gabriëlla, Mojdeh, Mechteld, Jens, Cesar, Marieke, Amelia, Aletta, Evelyn and Amy for their encouragement and support during different stages of my PhD course.

Dear Edsart, my dear fiancé, thank you so much for your unconditional support and patience in completing this PhD. I am so happy and lucky to share my life with you and I am looking forward to our future together.

Curriculum vitae

Pauline Antoinette Josephine was born on September 21, 1991, in Leiderdorp, The Netherlands. She spent her early childhood in Leiden with her parents and brother Michiel. During her own orthodontic treatment, she realized that she wanted to become an orthodontist. She graduated from high school at the Rijnlands Lyceum in Oegstgeest in 2009

and moved to Groningen to study dentistry. During her study of dentistry, she did several internships in orthodontic practices and wrote her master's dissertation at the orthodontic department of the University Medical Center Groningen (UMCG). In July 2016 she completed her dentistry study and in October 2016 she started as a resident in orthodontics at the Academic Centre for Dentistry Amsterdam from 2016-2020. In 2018 she started her PhD at the department of Orthodontics of the Academic Center for Dentistry Amsterdam. In 2020 she started working as an orthodontist in several practices.





CHAPTER 13

PRISMA-P 2015 Checklist

This checklist has been adapted for use with systematic review protocol submissions to BioMed Central journals from Table 3 in Moher D et al: Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Systematic Reviews 2015 4:1

An Editorial from the Editors-in-Chief of Systematic Reviews details why this checklist was adapted - Moher D, Stewart L & Shekelle P. Implementing PRISMA-P: recommendations for prospective authors. Systematic Reviews 2016 5:15

Constantiants	-		Information reported	n reported	Line
Section/topic		Checklist Item	Yes	Ŷ	number(s)
ADMINISTRATIVE INFORMATION	NFORMA	Not			
Title: Seeking adver	rse effect	Title: Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study			
Identification	ta.	Identify the report as a protocol of a systematic review	×		line 2
Update	\$	If the protocol is for an update of a previous systematic review, identify as such		×	Not applicable
Registration	N	If registered, provide the name of the registry (e.g., PROSPERO) and registration number in the Abstract		×	Not applicable
Authors					
Contact	8	Provide name, institutional affiliation, and e-mail address of all protocol authors; provide physical mailing address of corresponding author	×		lines 24-28
Contributions	æ	Describe contributions of protocol authors and identify the guarantor of the review	×		Lines 360-365
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol identify as such and list changes; otherwise, state plan for documenting important protocol amendments		×	Not applicable
Support					
Sources	5a	Indicate sources of financial or other support for the review	×		Lines 356-358
Sponsor	8	Provide name for the review funder and/or sponsor		×	Not applicable
Role of sponsor/funder	3	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol		×	Not applicable
INTRODUCTION					
Rationale	9	Describe the rationale for the review in the context of what is already known	×		Lines 89-128

Additional files chapter 2

Additonal file 1. PRISMA-P Checklist

Section/topic		Checklist item	Informatio	nformation reported	
			Yes	Ŷ	number(s)
Objectives	~	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	×		Lines 132-142
METHODS					
Eligibility criteria	60	Spacify the study characteristics (e.g., PICO, study design, setting, time frame) and report characteristics (e.g., years considered, language, publication status) to be used as criteria for eligibility for the review	×		Lines 156-201
Information sources	m	Describe all intended information sources (a.g., electronic databases, contact with study authors, trial registers, or other grey literature sources) with planned dates of coverage	×		Lines 204-213
Search strategy	\$	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated	×		Lines 204-213
STUDY RECORDS					
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	×		Lines 217-226
Selection process	116	State the process that will be used for selecting studies (e.g., two independent reviewers) through each phase of the review (i.e., screening, eligibility, and inclusion in meta-analysis)	×		Lines 227-237
Data collection process	11c	Describe planned method of extracting data from reports (e.g., ploting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators	×		Lines 238-252
Data Items	5	List and define all variables for which data will be sought (e.g., PICO items, funding sources), any pre-planned data assumptions and simplifications	×		Lines 238-263
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	×		Lines 265-277
Risk of bias in Individual studies	7	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both, state how this information will be used in data synthesis		×	Not applicable
DATA					
	15a	Describe criteria under which study data will be quantitatively synthesized		×	Not applicable
Synthesis	156	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data, and methods of combining data from studies, including any planned exploration of consistency (e.g., <i>i</i> ² , Kendalf's tau)		×	Not applicable

		Chronolding Barne	Informatio	n reporte	d Line
Securioria			Yes	Ŷ	number(s)
	15c D	Describe any proposed additional analyses (e.g., sensitivity or subgroup analyses, meta- regression)		×	Not applicable
	154	If quantitative synthesis is not appropriate, describe the type of summary planned		×	Not applicable
Meta-bias(es)	2	Specify any planned assessment of meta-blas(es) (e.g., publication blas across studies, selective reporting within studies)		×	Not applicable
Confidence in cumulative evidence	11	17 Describe how the strength of the body of evidence will be assessed (e.g., GRADE)		×	Not applicable

Additional file 2. Pilot tests

Eligible systematic reviews of orthodontic interventions published in the eligible journals were used for our pilot tests and were stratified evenly over the eligible years and journals. We calculated the required sample size of our pilot study based on the probability of Yes scores for our primary research question 'Did the review seek any findings related to adverse effects of interventions in the included studies?

To calculate this sample size we used the following equation [24]:

$$n = \frac{\ln(1-Y)}{\ln(1-\pi)}$$

n = the sample size for the pilot study

 Υ = the threshold of confidence (95%)

 π = the probability of a 'Yes' score

We calculated the sample size separately for systematic reviews of orthodontic interventions in the Cochrane Database of Systematic Reviews and those published in the 5 leading orthodontic journals. We used the probabilities of the Yes scores identified during our scoping searches as our benchmark. These probabilities were respectively 1 and 0.25. Based on these probabilities we calculated the required sample sizes of respectively 1 and 10.4. To play it safe we planned to include at least 2 Cochrane reviews and 12 orthodontic reviews in our pilot study. We planned increasing the size of the pilot study when the probability of yes scores for the pertinent review types would be inferior in the pilot studies compared to those identified in the scoping searches. We used random number generator software [25] to select the pilot reviews. We searched the first eligible review published in the random issue number of the pertinent journal for the years 2009, 2010, and 2011. If no eligible review was identified for that issue, we searched the first eligible review published in subsequent issues of the stratified journal subgroup. If no eligible reviews were identified for that period, we searched in the issues published prior to the random issue number and back to the starting issue of the stratified journal subgroup. We applied this strategy for the 3 stratified subgroups, i.e., 2009-2011, 2012-2014, 2015-2017 (Table). The prevalence of eligible systematic reviews that sought any findings related to adverse effects in the included studies was 25% (3/12) for the orthodontic journals and 100% (2/2) for reviews identified in the Cochrane Database of Systematic reviews.

Journal	Year and	Random	tions selected for the pilot test Selected systematic review
Journal	number of	issue	Selected systematic review
	issues	number	
Cochrane Database	2009, 2010,	24	No reference identified
of Systematic	and 2011 (28	24	
reviews	issues)		
Cochrane Database	2012, 2013,	31	Agostino P, Ugolini A, Signori A, Silvestrini-Biavati A, Harrison
of Systematic	and 2014	51	JE, Riley P.
reviews	(36 issues)		Orthodontic treatment for posterior crossbites.
Terretto	(50 155005)		Cochrane Database Syst Rev. 2014 Aug 8;(8):CD000979.
Cochrane Database	2015, 2016,	12	Littlewood SJ, Millett DT, Doubleday B, Bearn DR, Worthington
of Systematic	and 2017 (36	12	HV. Retention procedures for stabilising tooth position after
reviews	issues)		treatment with orthodontic braces Cochrane Database Syst Rev.
Terretto	1554657		2016 Jan 29;(1):CD002283.
European Journal	2009, 2010,	10	Naoumova J, Kurol J, Kjellberg H. A systematic review of the
of Orthodontics	and 2011 (18	10	interceptive treatment of palatally displaced maxillary canines.
or or thoughties	issues)		Eur J Orthod. 2011 Apr;33(2):143-9.
European Journal	2012, 2013,	5	Zuccati G, Casci S, Doldo T, Clauser C. Expansion of maxillary
of Orthodontics	and 2014	5	arches with crossbite: a systematic review of RCTs in the last 12
of of thoughties	(18 issues)		years. Eur J Orthod. 2013 Feb;35(1):29-37.
European Journal	2015, 2016,	3	Koretsi V, Zymperdikas VF, Papageorgiou SN, Papadopoulos MA.
of Orthodontics	and 2017 (18	5	Treatment effects of removable functional appliances in patients
of of thoughties	issues)		with Class II malocclusion: a systematic review and meta-analysis.
	1354637		Eur J Orthod. 2015 Aug;37(4):418-34.
American Journal	2009, 2010,	18	Chen SS, Greenlee GM, Kim JE, Smith CL, Huang GJ. Systematic
of Orthodontics	and 2011 (36	10	review of self-ligating brackets. Am J Orthod Dentofacial Orthop.
and Dentofacial	issues)		2010 Jun;137(6):726.e1-726.e18; discussion 726-7.
Orthopedics	1554657		
American Journal	2012, 2013,	35	Yang X, Li C, Bai D, Su N, Chen T, Xu Y, Han X.
of Orthodontics	and 2014	55	Treatment effectiveness of Fränkel function regulator on
and Dentofacial	(36 issues)		the Class III malocclusion: a systematic review and meta-analysis.
Orthopedics	(00.00000)		Am J Orthod Dentofacial Orthop. 2014 Aug;146(2):143-54.
American Journal	2015, 2016,	9	Nucera R, Lo Giudice A, Rustico L, Matarese G, Papadopoulos
of Orthodontics	and 2017 (36	-	MA, Cordasco G.
and Dentofacial	issues)		Effectiveness of orthodontic treatment with functional
Orthopedics	,		appliances on maxillary growth in the short term: A systematic
			review and meta-analysis. Am J Orthod Dentofacial Orthop. 2016
			May;149(5):600-611.e3.
Angle Orthodontist	2009, 2010,	6	Leonardi R, Annunziata A, Licciardello V, Barbato E.
	and 2011 (18	-	Soft tissue changes following the extraction of premolars in nongr
	issues)		owing patients with bimaxillary protrusion. A systematic review.
	,		Angle Orthod. 2010 Jan;80(1):211-6.
Angle Orthodontist	2012, 2013,	4	Feng X, Li J, Li Y, Zhao Z, Zhao S, Wang J.
	and 2014		Effectiveness of TAD-
	(18 issues)		anchored maxillary protraction in latemixed dentition. Angle
	. ,		Orthod. 2012 Nov;82(6):1107-14.
Angle Orthodontist	2015, 2016,	13	Diar-Bakirly S, Feres MF, Saltaji H, Flores-Mir C, El-Bialy T.
	and 2017 (18		Effectiveness of the transpalatal arch in controlling orthodontic
	issues)		anchorage in maxillary premolar extraction cases: A systematic
	-		review and meta-analysis. Angle Orthod. 2017 Jan;87(1):147-158.
The Korean Journal	2009, 2010,	7	No reviews identified in all 18 issues
of Orthodontics	and 2011 (18		
	issues)		
The Korean Journal	2012, 2013,	14	No reviews identified in all 18 issues
of Orthodontics	and 2014		
	•	•	101

Table. Systematic reviews of orthodontic interventions selected for the pilot test
--

	(18 issues)		
The Korean Journal	2015, 2016,	7	Papageorgiou SN, Höchli D, Eliades T. Outcomes of
of Orthodontics	and 2017 (18		comprehensive fixed appliance orthodontic treatment: A
	issues)		systematic review with meta-analysis and methodological
			overview. Korean J Orthod. 2017 Nov; 47(6): 401–413.
Orthodontics and	2009, 2010,	1	Several reviews were identified, but none were eligible
Craniofacial	and 2011 (12		
Research	issues)		
Orthodontics and	2012, 2013,	3	Cordasco G, Matarese G, Rustico L, Fastuca S, Caprioglio A,
Craniofacial	and 2014		Lindauer SJ, Nucera R. Efficacy of orthopedic treatment with
Research	(12 issues)		protraction facemask on skeletal Class III malocclusion: a
			systematic review and meta-analysis. Orthod Craniofac Res. 2014
			Aug;17(3):133-43.
Orthodontics and	2015, 2016,	1	Al-Saleh MA, Alsufyani N, Flores-Mir C, Nebbe B, Major PW.
Craniofacial	and 2017 (12		Changes in temporomandibular joint morphology in class II patien
Research	issues)		ts treated with fixed mandibular repositioning and evaluated thro
			ugh 3D imaging: a systematic review. Orthod Craniofac Res. 2015
			Nov;18(4):185-201.

Additional file 3. Search terms and their derivatives

Search terms and their derivatives	Search terms for searching multiple words
	in a PDF
"adverse"	ADVERSE, Adverse, adverse
"effect", "effects"	EFFECT, Effect, effect
"reaction", "reactions"	REACTION, Reaction, reaction
"complication", "complications", "complicated",	COMPLICAT, Complicat, complicat
"complicating"	
"harm", "harms", "harmful"	HARM, Harm, harm
"risk", "risks", "risky"	RISK, Risk, risk
"safe", "safety"	SAFE, Safe, safe
"side"	SIDE, Side, side
"toxic", "toxicity"	TOXIC, Toxic, toxic
"benefit", "benefits"	BENEFIT, Benefit, benefit
"result", "results"	RESULT, Result, result
"finding", "findings"	FINDING, Finding, finding
"outcome", "outcomes"	OUTCOME, Outcome, outcome
"limitation", "limitations", limit	LIMIT, Limit, limit
"damage", "damages", "damaging"	DAMAGE, Damage, damage
"data"	DATA, Data, data
"information"	INFO, Info, info
"conflict", "conflicts", "conflicting"	CONFLICT, Conflict, conflict
"negative"	NEGATIVE, Negative, negative
"detrimental"	DETRIMENTAL, Detrimental, detrimental
"disadvantage", "disadvantages", "disadvantageous"	DISADVAN, Disadvan, disadvan
"down"	DOWN, Down, down
"injury", "injuries", "injured", "injurious"	INJUR, Injur, injur
"byproduct", "byproducts"	BYPRODUCT, Byproduct, byproduct
"collateral"	COLLATERAL, Collateral, collateral
"unfavorable", "unfavourable"	UNFAVO, Unfavo, unfavo
"destructive"	DESTRUCT, Destruct, destruct
"unsafe"	UNSAFE, Unsafe, unsafe
"undesired", "undesirable"	UNDESIR, Undesir, undesir
"recommend", "recommendation", "recommending"	RECOMMEND, Recommend, recommend
"emergency", "emergencies"	EMERGEN, Emergen, emergen

Table. Search terms and their derivatives

Additional file 4. Data collection forms

Table. Data collection forms*	1
Items for the main manuscript	Description
Journal	List the pertinent journal
Year	Year of publication
Binder page number	List the binder page number
Reference	List full reference (Authors, Title, Journal)
Is the article a systematic review?	Answer: Yes/No
	Consider definition of a systematic review
What type of systematic review?	List the type of systematic review.
	Consider different types of systematic reviews.
	When the publication is not an intervention
	systematic review describe what type it is or could
	be and classify. Types of systematic reviews will
	receive a final classification during the discussions
	between operators.
Were orthodontic interventions assessed?	Answer: Yes/No
	Consider the definition of orthodontic interventions.
What was the orthodontic intervention?	List the type of orthodontic intervention
	NA: When the article is not a systematic review or
	not a systematic review of interventions.
Is the systematic review eligible?	Answer: Yes/No
	Yes: The article is a systematic review of an
	orthodontic intervention.
	No: The article is not a systematic review of an
	orthodontic intervention.
	No: The article is a systematic review of an
	orthodontic intervention, but focusses exclusively on
	its adverse effects.
Page and potential comments**	Present the pertinent pages of reference for scoring
	the previous items and list the potential comments.
Was seeking of adverse effects of interventions	Answer: Yes/No
defined as a research objective of the review?	Yes: When seeking of adverse effects of
	interventions was defined as a research objective or
	as a research question or when adverse effects were
	predefined a priori as outcomes to assess.
	No: Seeking of adverse effects of interventions was
	not defined as a research objective or as a research
	question or when adverse effects were not
	predefined a priori as outcomes to assess.
What adverse effects of interventions were defined	Answer: List adverse effects/NA
as research objectives?	List all adverse effects of interventions that the
	reviewers defined as research objectives.
	NA: When the following question was answered with
	a 'No': 'Was seeking of adverse effects of
	interventions defined as a research objective of the
	review?'
Did the review seek any findings related to adverse	Answer: Yes/No
effects of interventions in the included studies?	Yes: Any findings related to adverse effects of
	interventions in the included studies were sought by
	the reviewers.
	Seeking any findings related to adverse effects of
	interventions in the included studies refers to
	reporting anywhere in the review (except in the

Table. Data collection forms*

	Abstract) that such adverse effects in the included studies were sought. Yes: Yes is also scored when reviewers only reported findings related to adverse effects of interventions in the included studies, but did not report that they actually sought them or planned to seek them. For example 'Yes' will be scored when outcomes on adverse effects of interventions in the included studies were reported in the review, but were not
	defined as objectives of the review. Yes: Yes is also scored when the reviewers reported that they planned to seek (for example in the research objectives) findings related to adverse
	effects of interventions in the included studies, but did not report on these findings. No: Findings related to adverse effects of interventions in the included studies were not
	sought by the reviewers.
Did the review report findings related to adverse effects of interventions sought in the included studies?	Answer: Yes/No/NA Yes: The review reported findings related to adverse effects of interventions sought in the included studies.
	'Yes' is also scored when the review reported that no findings on adverse effects of interventions in the included studies were identified.
	No: The review did not report any findings related to adverse effects of interventions sought in the included studies.
	NA: When the following question was answered with a No: 'Did the review seek any findings related to adverse effects of interventions in the included studies?'
What findings related to adverse effects of interventions sought in the included studies were reported in the review?	Answer: List of adverse effects/NA List all findings related to adverse effect(s) of interventions that were identified in the included studies and reported in the review. NA: When the following question was answered with a 'No': 'Did the review seek any findings related to adverse effects of interventions in the included studies?'
Rationale for assigning an effect as 'adverse' or 'not adverse' (In case of additional or ambivalent adverse effects)	Answer: Present the rationale for assigning an effect as 'adverse' or 'not adverse' (In case of additional or ambivalent adverse effects)
Page and potential comments**	Present the pertinent pages of reference for scoring the previous items and list the potential comments.
Were potential adverse effects of the intervention considered, discussed (weighed) anywhere in the review?	Answer: Yes/No Yes: Potential adverse effect(s) of interventions in the included studies were sought and reported by the reviewers. 'Yes' is also scored when potential adverse effect(s) of interventions were not sought, but only considered, discussed (weighed) anywhere in the review. 'No' is scored when potential adverse effects of the
	intervention were not considered, discussed (weighed) anywhere in the review.

Rationale for assigning an effect as 'adverse' or 'not adverse' (In case of additional or ambivalent adverse effects)	Answer: Present the rationale
Page and potential comments**	Present the pertinent pages of reference for scoring the previous items and list the potential comments.

*To address our research question we will not consider what was reported regarding this question in the abstract and in the protocol of the review.

**When referring to a particular page in the systematic review, we will use the page number of the systematic review and not the number in the binder document.

Additional files chapter 3

Additional file 1. STROBE checklist

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	ltem No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly	1
		used term in the title or the abstract	
		(b) Provide in the abstract an informative and	2 and
		balanced summary of what was done and what	3
		was found	
Introduction			
Background/rationale	2	Explain the scientific background and rationale	3 and
		for the investigation being reported	4
Objectives	3	State specific objectives, including any	5
		prespecified hypotheses	
Methods			
Study design	4	Present key elements of study design early in the	5
		paper	
Setting	5	Describe the setting, locations, and relevant	6 and
		dates, including periods of recruitment, exposure,	7
		follow-up, and data collection	
Participants	6	(a) Give the eligibility criteria, and the sources	6
		and methods of selection of participants	
Variables	7	Clearly define all outcomes, exposures,	8 and
		predictors, potential confounders, and effect	9
		modifiers. Give diagnostic criteria, if applicable	
Data sources/	8*	For each variable of interest, give sources of data	8 and
measurement		and details of methods of assessment	9
		(measurement). Describe comparability of	
		assessment methods if there is more than one	
		group	
Bias	9	Describe any efforts to address potential sources	NA
		of bias	
Study size	10	Explain how the study size was arrived at	8 and
			9
Quantitative variables	11	Explain how quantitative variables were handled	9
		in the analyses. If applicable, describe which	
		groupings were chosen and why	
Statistical methods	12	(a) Describe all statistical methods, including	9
		those used to control for confounding	

		(b) Describe any methods used to examine	9
		subgroups and interactions	
		(c) Explain how missing data were addressed	9
		(d) If applicable, describe analytical methods	NA
		taking account of sampling strategy	
		(<u>e</u>) Describe any sensitivity analyses	NA
Results			
Participants	13*	(a) Report numbers of individuals at each stage of	9 and
		study—eg numbers potentially eligible, examined	10
		for eligibility, confirmed eligible, included in the	
		study, completing follow-up, and analysed	
		(b) Give reasons for non-participation at each	NA
			NA
		stage	10
Deceriptive data	1.1*	(c) Consider use of a flow diagram	10
Descriptive data	14*	(a) Give characteristics of study participants (eg	10
		demographic, clinical, social) and information on	
		exposures and potential confounders	
		(b) Indicate number of participants with missing	NA
		data for each variable of interest	
Outcome data	15*	Report numbers of outcome events or summary	10
		measures	and
			11
Main results	16	(a) Give unadjusted estimates and, if applicable,	10
		confounder-adjusted estimates and their	and
		precision (eg, 95% confidence interval). Make	11
		clear which confounders were adjusted for and	
		why they were included	
		(b) Report category boundaries when continuous	NA
		variables were categorized	
		(c) If relevant, consider translating estimates of	NA
		relative risk into absolute risk for a meaningful	
		time period	
Other analyses	17	Report other analyses done—eg analyses of	11
o ther unuryses	17	subgroups and interactions, and sensitivity	
		analyses	
Discussion			1
Key results	18	Summarise key results with reference to study	11
-,		objectives	and
			12
Limitations	19	Discuss limitations of the study, taking into	12
	19	account sources of potential bias or imprecision.	and
			13
		Discuss both direction and magnitude of any	13
		potential bias	

Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	13 and 14
Generalisability	21	Discuss the generalisability (external validity) of the study results	14
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	15

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.

Additional file 2

Table of contents for additional file 2

Additional file item	Description
Additional file 2A	Difference between the protocol and the completed cross-sectional study
Additional file 2B	Selected journals and their 2018 impact factor (Clarivate Analytics 2021)
Additional file 2C	Data collection forms
Additional file 2D	Adverse effects hypothetically linked to orthodontic interventions according to Preoteasa et al. (2012)
Additional file 2E	Adverse effects hypothetically linked to orthodontic interventions
References for additional file 2	References for additional file 2

Additional file 2A. Differences between the protocol and the completed cross-sectional study

Differences between the protocol and the	Rationale
completed cross-sectional study	
Extension of the search period to identify eligible systematic reviews	We planned to complete our study on July 31 2019, but as a result of COVID 19-related delays we extended our search period to July 31 2021.
Excluding systematic reviews with Bayesian network meta-analysis	We excluded systematic reviews with Bayesian network meta- analyses, because such reviews include the results of multiple interventions, which could make it difficult to understand which adverse effect was assigned to which specific intervention.
Excluding systematic reviews that did not assess a specific orthodontic intervention, but refer to orthodontic interventions as a whole.	We excluded systematic reviews that did not assess a specific orthodontic intervention, but referred to orthodontic interventions as a whole.
Excluding systematic reviews that were conducted by one operator only	We excluded reviews that were conducted by one reviewer only, because such an approach is often conducted for narrative reviews, but is not congruent with the systematic approach of systematic reviews.
Implementing univariable logistic regression models in the statistical analyses	We implemented univariable logistic regression models to determine the association between each of the 4 outcomes and the journal in which the SR was published using the Cochrane Database of Systematic Reviews as reference. Chi- square tests of independence as planned in our protocol were not used, because of the small number of systematic reviews in each eligible journal and the low variability in the response scored (prevalence of 'no' ranging from 9.2% to 15.3%).
In the protocol the following definition was used for orthodontic interventions: Steegmans et al. (2019a): 'Orthodontic interventions refer to the use of any type of orthodontic appliance that are used to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodontic treatment, for example retainers.'	This definition was changed to: 'Orthodontic interventions refer to the use of any type of orthodontic appliance to move teeth or change the jaw size or position for orthodontic purposes. These interventions also include appliances to maintain or stabilize the results of orthodontic treatment, for example retainers.' Rationale for change: The modified definition is more accurate, but did not change the original meaning of the definition of orthodontic interventions
Inclusion of Dr. Nicola Di Girolama as an author	Dr. Di Girolamo was consulted for his assistance in statistical and methodological issues

Journal	Impact factor
Cochrane Database of Systematic reviews	7.755
American Journal of Orthodontics and Dentofacial Orthopedics	1.911
European Journal of Orthodontics	1.841
Korean Journal of Orthodontics	1.476
Orthodontics and Craniofacial Research	0.946

Additional file 2B. Selected journals and their 2018 impact factor (Clarivate Analytics 2021)

Additional file 2C. Data collection forms*

Items for the main manuscript	Description
Journal	List the pertinent journal
Year	Year of publication
Binder page number	List the binder page number
Reference	List full reference (Authors, Title, Journal)
Is the article a systematic review?	Answer: Yes/No
	Consider definition of a systematic review
Is the systematic review eligible?	Answer: Yes/No
	Consult the eligibility criteria for addressing this answer.
What was the orthodontic intervention?	List the type of orthodontic intervention.
Was seeking of adverse effects of interventions	Answer: Yes/No
defined as a research objective of the review?	Yes: When seeking of adverse effects of interventions was
	defined as a research objective or as a research question
	or when adverse effects were predefined a priori as
	outcomes to assess.
	No: Seeking of adverse effects of interventions was not
	defined as a research objective or as a research question
	or when adverse effects were not predefined a priori as
	outcomes to assess.
What adverse effects of interventions were defined as	Answer: List adverse effects/NA
research objectives?	List all adverse effects of interventions that the reviewers
	defined as research objectives.
	NA: When the following question was answered with a
	'No': 'Was seeking of adverse effects of interventions
	defined as a research objective of the review?'
Did the review seek any findings related to adverse	Answer: Yes/No
effects of interventions in the included studies?	Yes: Any findings related to adverse effects of
	interventions in the included studies were sought by the
	reviewers.
	Seeking any findings related to adverse effects of
	interventions in the included studies refers to reporting
	anywhere in the review (except in the Abstract) that such
	adverse effects in the included studies were sought. Yes: Yes is also scored when reviewers only reported
	findings related to adverse effects of interventions in the
	included studies, but did not report that they actually
	sought them or planned to seek them. For example 'Yes'
	will be scored when outcomes on adverse effects of
	interventions in the included studies were reported in the
	review, but were not defined as objectives of the review.
	Yes: Yes is also scored when the reviewers reported that
	they planned to seek (for example in the research

13

191

Did the review report findings related to adverse effects of interventions sought in the included studies?	No: Findings related to adverse effects of interventions in the included studies were not sought by the reviewers. Answer: Yes/No Yes: The review reported findings related to adverse effects of interventions sought in the included studies. 'Yes' is also scored when the review reported that no findings on adverse effects of interventions in the included studies were identified. No: The review did not report any findings related to adverse effects of interventions sought in the included studies.
What findings related to adverse effects of interventions sought in the included studies were reported in the review?	Answer: List of adverse effects List all findings related to adverse effect(s) of interventions that were identified in the included studies and reported in the review.
Rationale for assigning an effect as 'adverse' or 'not adverse' (In case of additional or ambivalent adverse effects)	Answer: Present the rationale for assigning an effect as 'adverse' or 'not adverse' (In case of additional or ambivalent adverse effects)
Were potential adverse effects of the intervention considered, discussed (weighed) anywhere in the review?	Answer: Yes/No Yes: Potential adverse effect(s) of interventions in the included studies were sought and reported by the reviewers. 'Yes' is also scored when potential adverse effect(s) of interventions were not sought, but only considered, discussed (weighed) anywhere (except in the Abstract) in the review. 'No' is scored when potential adverse effects of the intervention were not considered, discussed (weighed) anywhere in the review.
Rationale for assigning an effect as 'adverse' or 'not adverse' (In case of additional or ambivalent adverse effects)	Answer: Present the rationale

*To address our research question we will not consider what was reported regarding this question in the abstract and in the protocol of the review.

Additional file 2D. Adverse effects hypothetically linked to orthodontic interventions according to Preoteasa et al. (2012)*

Local adverse effects	
Subgroup	Description
Dental	 crown: decalcifications, decays, tooth wear, enamel cracks and fractures; discolorations, deterioration of prosthetic crown (as fracturing a ceramic one during debonding); root: root resorption, early closure of root apex, ankylosis; pulp: ischemia, pulpitis, necrosis;
Periodontal	 gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscences, fenestrations, interdental fold, dark triangles;
Temporomandibular joint	 condylar resorption, temporomandibular dysfunction;
Soft tissues of the oral and maxillofacial region	 trauma (e.g., long archwires, headgear related), mucosal ulcerations or hyperplasia, chemical burns (e.g., etching related), thermal injuries (e.g., overheated burs), stomatitis, clumsy handling of dental instruments;
Unsatisfactory treatment	inadequate morpho-functional, aesthetic or functional final result,
outcome	relapse, failure to complete treatment due to treatment dropout.
Systemic adverse effects	
Subgroup	Description
Psychological	 teasing, behavioral changes of patients and parents; discomfort associated with pain presence and aesthetic look discontents during orthodontic appliance usage;
Gastro-intestinal	 accidental swallowing of small parts of the orthodontic device (tubes, brackets);
Allergies	to nickel or latex;
Cardiac	infective endocarditis;
Chronic fatigue syndrome	
Cross infections	 from doctor to patient, patient to doctor, patient to patient.

*Permission to reproduce this table was obtained on August 16 2018 from InTech's Publishing Ethics and Legal Affairs Department.

Additional file 2E. Adverse effects hypothetically	v linked to orthodontic interventions*

Adverse effects related to	Description
Tooth structures	Tooth crown
	 decalcifications, decays, tooth wear, enamel cracks and fractures;
	discolorations, deterioration of prosthetic crown (as fracturing a
	ceramic one during debonding);
	• iatrogenic damage to the crown, e.g., fracture as a result of trauma
	Tooth root
	 root resorption, early closure of root apex, ankylosis;
	 iatrogenic damage to the root, e.g., fracture as a result of trauma
	Tooth pulp
	 ischemia, pulpitis, necrosis istragonia damaga to the pulping of fracture of a result of trauma
Periodontal tissues	 iatrogenic damage to the pulp, e.g., fracture as a result of trauma gingivitis, periodontitis, gingival recession or hypertrophy, alveolar
renouontar tissues	bone loss, dehiscences, fenestrations, interdental fold, dark
	triangles; tooth mobility, plague retention, bacterial count
Intraoral (non-tooth or	 intraoral tissue irritations and inflammation such as mucosal
periodontal) tissues	ulcerations or hyperplasia or irritations of the tongue (as a result of
· ,	trauma by appliances, e.g., breakage, failure, loosening etc. of
	appliances or long arch wires)
	Scar formation after suturing
	 chemical burns (e.g., etching related)
	 thermal injuries (e.g., overheated burs)
	nerve damage
	 tooth eruption, i.e., eruption disturbances (e.g., impactions) caused
	by orthodontic appliances
Extraoral tissues (non-	 cutting of lips or cheeks, eye injury (e.g., as a result of trauma by
temporomandibular tissues)	appliances, e.g., breakage, failure, loosening etc. of appliances or
	long arch wires or headgear-related trauma)
Townson and the low times and	discomfort on the lip
Temporomandibular tissues and disorders	 temporomandibular tissues and disorders
Appliance failure	 breakage, failure, loosening etc. of appliances
	 long archwires, headgear-related trauma
Undesired treatment results	inadequate morpho-functional, aesthetic or functional final result
	 inaccuracy of the treatment result
	 non predictability of the treatment result
	 Dental side effects e.g., unwanted tipping of teeth, anchorage loss
	etc.
	 Skeletal side effects, e.g., unwanted backward rotation of the
	mandible
Relapse and stability	Relapse and stability of the obtained treatment result
Undesired qualitative	Pain and discomfort
and the second	
experiences by the patient or	orthodontic tooth movement-related pain and discomfort
experiences by the patient or carer(s)	appliance (intervention)-related pain and discomfort: i.e., pain and
	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or
	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g.,
	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g., tension or pressure of the appliances (constriction of appliances),
	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g., tension or pressure of the appliances (constriction of appliances), speech difficulties, eating difficulties, swallowing difficulties, food
	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g., tension or pressure of the appliances (constriction of appliances), speech difficulties, eating difficulties, swallowing difficulties, food accumulation, bad tastes and smells
	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g., tension or pressure of the appliances (constriction of appliances), speech difficulties, eating difficulties, swallowing difficulties, food

	 Tolerability/acceptance/stress issues with the treatment procedures Absence from work or studies and difficulties in daily activities collaboration (compliance) issues or failure to complete treatment, e.g., dropout patient anxiety being teased social discomfort
	 embarrassment to wear the appliance behavioral changes of patients and parents, impaired family relationships aesthetic look discontents during orthodontic appliance usage concentration difficulties
	 concern ation difficulties reduced enjoyment of food and change in taste sleeping difficulties removal of appliance during sleep development of mannerisms
	 Satisfaction with the treatment procedures and final result not satisfied with the treatment procedures (Check in text what was measured, i.e., during or after) not satisfied with the final treatment result (Check in text what was measured, i.e., during or after)
Gastro-intestinal	 accidental swallowing parts of the orthodontic device (tubes, brackets);
Allergy	Allergies to nickel or latex;
Cardio	infective endocarditis;
Chronic fatigue	
Cross infections	 from doctor to patient, patient to doctor, patient to patient.
Non-defined	Adverse effects that were not defined by the authors of the review: referring to 'any adverse effect', 'any side effect' etc.
Additional adverse effects	Additional adverse effects that were identified during data extraction that could not be labeled under any of the categories of adverse effects given in this table

*Modified from Preoteasa et al. (Preoteasa 2012)

References for additional file 2

Clarivate Analytics 2021

Clarivate Analytics. [online] Available from: <u>https://clarivate.com/</u> (accessed December 4th 2021).

Preoteasa 2012

Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F. (editor). Orthodontics-Basic aspects and clinical considerations. March 9, 2012 under CC BY 3.0 license. www.intechopen.com. [online] Available from: https://cdn.intechopen.com/pdfs/31388/InTech-

Risks and complicationsassociated with orthodontic treatment.pdf (accessed December 4th 2021).

Steegmans 2019a

Steegmans PAJ, Bipat S, Meursinge Reynders RA. Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study. Syst Rev. 2019 Apr 5;8(1):89. doi: 10.1186/s13643-019-1000-1. PMID: 30953538; PMCID: PMC6449933.

Additional file 3. Included reviews

Journal*	Year	Reference
		Batista KB, Thiruvenkatachari B, Harrison JE, O'Brien KD. Orthodontic treatment for
Cochrane		prominent upper front teeth (Class II malocclusion) in children and adolescents.Cochrane
library	2018	Database Syst Rev. 2018 Mar 13;3:CD003452. doi: 10.1002/14651858.CD003452.pub4.
library	2010	Wang Y, Liu C, Jian F, McIntyre GT, Millett DT, Hickman J, Lai W. Initial arch wires used in
Cochrane		orthodontic treatment with fixed appliances.Cochrane Database Syst Rev. 2018 Jul
library	2018	31;7:CD007859. doi: 10.1002/14651858.CD007859.pub4.
library	2018	Borrie FR, Bearn DR, Innes NP, Iheozor-Ejiofor Z. Interventions for the cessation of non-
Cochrane		nutritive sucking habits in children. Cochrane Database Syst Rev. 2015 Mar
	2015	· · · ·
library	2015	31;(3):CD008694. doi: 10.1002/14651858.CD008694.pub2.
Casharan		Fleming PS, Fedorowicz Z, Johal A, El-Angbawi A, Pandis N. Surgical adjunctive procedures
Cochrane	2015	for accelerating orthodontic treatment. Cochrane Database Syst Rev. 2015 Jun
library	2015	30;(6):CD010572. doi: 10.1002/14651858.CD010572.pub2.
		El-Angbawi A, McIntyre GT, Fleming PS, Bearn DR. Non-surgical adjunctive interventions
		for accelerating tooth movement in patients undergoing fixed orthodontic treatment.
Cochrane		Cochrane Database Syst Rev. 2015 Nov 18;(11):CD010887. doi:
library	2015	10.1002/14651858.CD010887.pub2.
		Agostino P, Ugolini A, Signori A, Silvestrini-Biavati A, Harrison JE, Riley P. Orthodontic
Cochrane		treatment for posterior crossbites. Cochrane Database Syst Rev. 2014 Aug
library	2014	8;(8):CD000979. doi: 10.1002/14651858.CD000979.pub2.
		Jambi S, Walsh T, Sandler J, Benson PE, Skeggs RM, O'Brien KD. Reinforcement of
		anchorage during orthodontic brace treatment with implants or other surgical methods.
Cochrane		Cochrane Database Syst Rev. 2014 Aug 19;(8):CD005098. doi:
library	2014	10.1002/14651858.CD005098.pub3.
		Lentini-Oliveira DA, Carvalho FR, Rodrigues CG, Ye Q, Prado LB, Prado GF, Hu R.
Cochrane		Orthodontic and orthopaedic treatment for anterior open bite in children. Cochrane
library	2014	Database Syst Rev. 2014 Sep 24;(9):CD005515. doi: 10.1002/14651858.CD005515.pub3.
		Watkinson S, Harrison JE, Furness S, Worthington HV.Orthodontic treatment for
Cochrane		prominent lower front teeth (Class III malocclusion) in children. Cochrane Database Syst
library	2013	Rev. 2013 Sep 30;(9):CD003451. doi: 10.1002/14651858.CD003451.pub2.
		Jambi S, Thiruvenkatachari B, O'Brien KD, Walsh T. Orthodontic treatment for distalising
Cochrane		upper first molars in children and adolescents. Cochrane Database Syst Rev. 2013 Oct
library	2013	23;(10):CD008375. doi: 10.1002/14651858.CD008375.pub2.
		Afzal E, Fida M, Malik DS, Irfan S, Gul M. Comparison between conventional and
		piezocision-assisted orthodontics in relieving anterior crowding: a systematic review and
		meta-analysis. Eur J Orthod. 2021 Jun 8;43(3):360-366. doi: 10.1093/ejo/cjaa046. PMID:
EJO	2021	32812636.
		Kapetanović A, Theodorou CI, Bergé SJ, Schols JGJH, Xi T. Efficacy of Miniscrew-Assisted
		Rapid Palatal Expansion (MARPE) in late adolescents and adults: a systematic review and
		meta-analysis. Eur J Orthod. 2021 Jun 8;43(3):313-323. doi: 10.1093/ejo/cjab005. PMID:
EJO	2021	33882127; PMCID: PMC8186837.
		Rutili V, Mrakic G, Nieri M, Franceschi D, Pierleoni F, Giuntini V, Franchi L. Dento-skeletal
	1	effects produced by rapid versus slow maxillary expansion using fixed jackscrew
		expanders: a systematic review and meta-analysis. Eur J Orthod. 2021 Jun 8;43(3):301-
EJO	2021	312. doi: 10.1093/ejo/cjaa086. PMID: 33950178.
		Cornelis MA, Tepedino M, Riis NV, Niu X, Cattaneo PM. Treatment effect of bone-
		anchored maxillary protraction in growing patients compared to controls: a systematic
		review with meta-analysis. Eur J Orthod. 2021 Jan 29;43(1):51-68. doi:
		10 1000 / 1 / 1 010 00 / 000
EJO	2021	10.1093/ejo/cjaa016. PMID: 32815989.
EJO	2021	10.1093/ejo/cjaa016. PMID: 32815989. Shahabee M, Shafaee H, Abtahi M, Rangrazi A, Bardideh E. Effect of micro-

Included systematic reviews of orthodontic interventions

EJO		Consider Economics D. Contes M. Mondos CMDA. Nerrorada D. Maradibular service bits
EJO		González Espinosa D, Santos M, Mendes SMDA, Normando D. Mandibular propulsion
EJO		appliance for adults with Class II malocclusion: a systematic review and meta-analysis. Eur
	2020	J Orthod. 2020 Apr 1;42(2):163-173. doi: 10.1093/ejo/cjz089. PMID: 31786599.
		Mohammed H, Čirgić E, Rizk MZ, Vandevska-Radunovic V. Effectiveness of prefabricated
		myofunctional appliances in the treatment of Class II division 1 malocclusion: a systematic
		review. Eur J Orthod. 2020 Apr 1;42(2):125-134. doi: 10.1093/ejo/cjz025. PMID:
EJO	2020	31329848.
		Lyu C, Zhang L, Zou S. The effectiveness of supplemental vibrational force on enhancing
		orthodontic treatment. A systematic review. Eur J Orthod. 2019 Sep 21;41(5):502-512.
EJO	2019	doi: 10.1093/ejo/cjz018. PMID: 31065683.
		Algharbi M, Bazargani F, Dimberg L. Do different maxillary expansion appliances influence
		the outcomes of the treatment?Eur J Orthod. 2018 Jan 23;40(1):97-106. doi:
EJO	2018	10.1093/ejo/cjx035.
		Al Rahma WJ, Kaklamanos EG, Athanasiou AE. Performance of Hawley-type retainers: a
		systematic review of randomized clinical trials.Eur J Orthod. 2018 Apr 6;40(2):115-125.
EJO	2018	doi: 10.1093/ejo/cjx036.
		Feres MF, Abreu LG, Insabralde NM, de Almeida MR, Flores-Mir . Effectiveness of open
		bite correction when managing deleterious oral habits in growing children and
		adolescents: a systematic review and meta-analysis. Eur J Orthod. 2017 Feb;39(1):31-42.
EJO	2017	doi: 10.1093/ejo/cjw005. Epub 2016 Feb 3.
		Papageorgiou SN, Kutschera E, Memmert S, Gölz L, Jäger A, Bourauel C, Eliades T.
		Effectiveness of early orthopaedic treatment with headgear: a systematic review and
EJO	2017	meta-analysis. Eur J Orthod. 2017 Apr 1;39(2):176-187. doi: 10.1093/ejo/cjw041.
		Zymperdikas VF, Koretsi V, Papageorgiou SN, Papadopoulos MA. Treatment effects of
		fixed functional appliances in patients with Class II malocclusion: a systematic review and
EJO	2016	meta-analysis. Eur J Orthod. 2016 Apr;38(2):113-26. doi: 10.1093/ejo/cjv034.
		Feres MF, Abreu LG, Insabralde NM, Almeida MR, Flores-Mir C. Effectiveness of the open
		bite treatment in growing children and adolescents. A systematic review. Eur J Orthod.
EJO	2016	2016 Jun;38(3):237-50. doi: 10.1093/ejo/cjv048.
		Yang X, Zhu Y, Long H, Zhou Y, Jian F, Ye N, Gao M, Lai W. The effectiveness of the Herbst
		appliance for patients with Class II malocclusion: a meta-analysis. Eur J Orthod. 2016
EJO	2016	Jun;38(3):324-33. doi: 10.1093/ejo/cjv057.
		Mistakidis I, Katib H, Vasilakos G, Kloukos D, Gkantidis N. Clinical outcomes of lingual
		orthodontic treatment: a systematic review. Eur J Orthod. 2016 Oct;38(5):447-58. doi:
EJO	2016	10.1093/ejo/cjv061.
		Elkordy SA, Aboelnaga AA, Fayed MM, AboulFotouh MH, Abouelezz AM.Can the use of
		skeletal anchors in conjunction with fixed functional appliances promote skeletal
		changes? A systematic review and meta-analysis. Eur J Orthod. 2016 Oct;38(5):532-45.
EJO	2016	doi: 10.1093/ejo/cjv081.
		Pacha MM, Fleming PS, Johal A. A comparison of the efficacy of fixed versus removable
		functional appliances in children with Class II malocclusion: A systematic review. Eur J
EJO	2016	Orthod. 2016 Dec;38(6):621-630.
		Ehsani S, Nebbe B, Normando D, Lagravere MO, Flores-Mir C. Short-term treatment
		effects produced by the Twin-block appliance: a systematic review and meta-analysis. Eur
EJO	2015	J Orthod. 2015 Apr;37(2):170-6. doi: 10.1093/ejo/cju030.
		Zurfluh MA, Kloukos D, Patcas R, Eliades T. Effect of chin-cup treatment on the
		temporomandibular joint: a systematic review. Eur J Orthod. 2015 Jun;37(3):314-24. doi:
EJO	2015	10.1093/ejo/cju048.
		Koretsi V, Zymperdikas VF, Papageorgiou SN, Papadopoulos MA. Treatment effects of
		removable functional appliances in patients with Class II malocclusion: a systematic
EJO	2015	review and meta-analysis. Eur J Orthod. 2015 Aug;37(4):418-34. doi: 10.1093/ejo/cju071.
	_010	Liu S, Xu T, Zou W. Effects of rapid maxillary expansion on the midpalatal suture: a
EJO	2015	systematic review. Eur J Orthod. 2015 Dec;37(6):651-5. doi: 10.1093/ejo/cju100.

		Thou V Long H Vo N Vuo L Vong V Lipo L Loi W/ The effectiveness of non-survival
		Zhou Y, Long H, Ye N, Xue J, Yang X, Liao L, Lai W. The effectiveness of non-surgical maxillary expansion: a meta-analysis. Eur J Orthod. 2014 Apr;36(2):233-42. doi:
EJO	2014	10.1093/ejo/cjt044.
LJO	2014	Papageorgiou SN, Konstantinidis I, Papadopoulou K, Jäger A, Bourauel C. Clinical effects of
		pre-adjusted edgewise orthodontic brackets: a systematic review and meta-analysis. Eur J
EJO	2014	Orthod. 2014 Jun;36(3):350-63. doi: 10.1093/ejo/cjt064.
200		Zuccati G, Casci S, Doldo T, Clauser C. Expansion of maxillary arches with crossbite: a
		systematic review of RCTs in the last 12 years. Eur J Orthod. 2013 Feb;35(1):29-37. doi:
EJO	2013	10.1093/ejo/cjr140.
		Fleming PS, Johal A, Pandis N. The effectiveness of laceback ligatures during initial
I		orthodontic alignment: a systematic review and meta-analysis. Eur J Orthod. 2013
EJO	2013	Aug;35(4):539-46. doi: 10.1093/ejo/cjs033.
		Perillo L, Cannavale R, Ferro F, Franchi L, Masucci C, Chiodini P, Baccetti T. Meta-analysis
		of skeletal mandibular changes during Frankel appliance treatment. Eur J Orthod. 2011
EJO	2011	Feb;33(1):84-92. doi: 10.1093/ejo/cjq033.
		Naoumova J, Kurol J, Kjellberg H. A systematic review of the interceptive treatment of
		palatally displaced maxillary canines. Eur J Orthod. 2011 Apr;33(2):143-9. doi:
EJO	2011	10.1093/ejo/cjq045.
		Santana LG, de Campos França E, Flores-Mir C, Abreu LG, Marques LS, Martins-Junior PA.
		Effects of lip bumper therapy on the mandibular arch dimensions of children and
		adolescents: A systematic review. Am J Orthod Dentofacial Orthop. 2020 Apr;157(4):454-
AJODO	2020	465.e1. doi: 10.1016/j.ajodo.2019.10.014. PMID: 32241352.
		Sivarajan S, Ringgingon LP, Fayed MMS, Wey MC. The effect of micro-osteoperforations
		on the rate of orthodontic tooth movement: A systematic review and meta-analysis. Am J Orthod Dentofacial Orthop. 2020 Mar;157(3):290-304. doi: 10.1016/j.ajodo.2019.10.009.
AJODO	2020	PMID: 32115107.
AJUDU	2020	Theodorou CI, Kuijpers-Jagtman AM, Bronkhorst EM, Wagener FADTG. Optimal force
		magnitude for bodily orthodontic tooth movement with fixed appliances: A systematic
		review. Am J Orthod Dentofacial Orthop. 2019 Nov;156(5):582-592. doi:
AJODO	2019	10.1016/j.ajodo.2019.05.011. PMID: 31677666.
10000	2015	Kouvelis G, Dritsas K, Doulis I, Kloukos D, Gkantidis N. Effect of orthodontic treatment with
		4 premolar extractions compared with nonextraction treatment on the vertical dimension
		of the face: A systematic review.Am J Orthod Dentofacial Orthop. 2018 Aug;154(2):175-
AJODO	2018	187. doi: 10.1016/j.ajodo.2018.03.007.
		Aljabaa A, Almoammar K, Aldrees A, Huang G. Effects of vibrational devices on
		orthodontic tooth movement: A systematic review.Am J Orthod Dentofacial Orthop. 2018
AJODO	2018	Dec;154(6):768-779. doi: 10.1016/j.ajodo.2018.07.012.
		Woon SC, Thiruvenkatachari B. Early orthodontic treatment for Class III malocclusion: A
		systematic review and meta-analysis. Am J Orthod Dentofacial Orthop. 2017
AJODO	2017	Jan;151(1):28-52. doi: 10.1016/j.ajodo.2016.07.017.
		Antoszewska-Smith J, Sarul M, Łyczek J, Konopka T, Kawala B. Effectiveness of orthodontic
		miniscrew implants in anchorage reinforcement during en-masse retraction: A systematic
	2017	review and meta-analysis. Am J Orthod Dentofacial Orthop. 2017 Mar;151(3):440-455.
AJODO	2017	doi: 10.1016/j.ajodo.2016.08.029.
		Nucera R, Lo Giudice A, Rustico L, Matarese G, Papadopoulos MA, Cordasco G.
		Effectiveness of orthodontic treatment with functional appliances on maxillary growth in the short term: A systematic review and meta-analysis. Am J Orthod Dentofacial Orthop.
AJODO	2016	2016 May;149(5):600-611.e3. doi: 10.1016/j.ajodo.2015.09.030.
1000	2010	Ishaq RA, AlHammadi MS, Fayed MM, El-Ezz AA, Mostafa Y. Fixed functional appliances
		with multibracket appliances have no skeletal effect on the mandible: A systematic review
		and meta-analysis. Am J Orthod Dentofacial Orthop. 2016 May;149(5):612-24. doi:
AJODO	2016	10.1016/j.ajodo.2015.11.023.
		Silveira GS, de Almeida NV, Pereira DM, Mattos CT, Mucha JN. Prosthetic replacement vs
		space closure for maxillary lateral incisor agenesis: A systematic review. Am J Orthod
AJODO	2016	Dentofacial Orthop. 2016 Aug;150(2):228-37. doi: 10.1016/j.ajodo.2016.01.018.

		Thiruvenkatachari B, Harrison J, Worthington H, O'Brien K. Early orthodontic treatment
		for Class II malocclusion reduces the chance of incisal trauma: Results of a Cochrane
		systematic review. Am J Orthod Dentofacial Orthop. 2015 Jul;148(1):47-59. doi:
AJODO	2015	10.1016/j.ajodo.2015.01.030.
		Hoogeveen EJ, Jansma J, Ren Y. Surgically facilitated orthodontic treatment: a systematic
		review. Am J Orthod Dentofacial Orthop. 2014 Apr;145(4 Suppl):S51-64. doi:
AJODO	2014	10.1016/j.ajodo.2013.11.019.
I		Yang X, Li C, Bai D, Su N, Chen T, Xu Y, Han X. Treatment effectiveness of Fränkel function
	201.4	regulator on the Class III malocclusion: a systematic review and meta-analysis. Am J
AJODO	2014	Orthod Dentofacial Orthop. 2014 Aug;146(2):143-54. doi: 10.1016/j.ajodo.2014.04.017.
		Janson G, Sathler R, Fernandes TM, Branco NC, Freitas MR. Correction of Class II
	2012	malocclusion with Class II elastics: a systematic review.Am J Orthod Dentofacial Orthop.
AJODO	2013	2013 Mar;143(3):383-92. doi: 10.1016/j.ajodo.2012.10.015.
		Grec RH, Janson G, Branco NC, Moura-Grec PG, Patel MP, Castanha Henriques JF. Intraoral
AJODO	2013	distalizer effects with conventional and skeletal anchorage: a meta-analysis. Am J Orthod
AJUDU	2013	Dentofacial Orthop. 2013 May;143(5):602-15. doi: 10.1016/j.ajodo.2012.11.024.
		Millett DT, Cunningham SJ, O'Brien KD, Benson PE, de Oliveira CM. Treatment and stability of class II division 2 malocclusion in children and adolescents: a systematic review. Am J
		Orthod Dentofacial Orthop. 2012 Aug;142(2):159-169.e9. doi:
AJODO	2012	10.1016/j.ajodo.2012.03.022.
AUDO	2012	Marsico E, Gatto E, Burrascano M, Matarese G, Cordasco G. Effectiveness of orthodontic
		treatment with functional appliances on mandibular growth in the short term. Am J
AJODO	2011	Orthod Dentofacial Orthop. 2011 Jan;139(1):24-36. doi: 10.1016/j.ajodo.2010.04.028.
10000	2011	Fudalei P, Antoszewska J. Are orthodontic distalizers reinforced with the temporary
		skeletal anchorage devices effective? Am J Orthod Dentofacial Orthop. 2011
AJODO	2011	Jun;139(6):722-9. doi: 10.1016/j.ajodo.2011.01.019.
		Baratieri C, Alves M Jr, de Souza MM, de Souza Araújo MT, Maia LC. Does rapid maxillary
		expansion have long-term effects on airway dimensions and breathing? Am J Orthod
AJODO	2011	Dentofacial Orthop. 2011 Aug;140(2):146-56. doi: 10.1016/j.ajodo.2011.02.019.
		Chen SS, Greenlee GM, Kim JE, Smith CL, Huang GJ. Systematic review of self-ligating
		brackets. Am J Orthod Dentofacial Orthop. 2010 Jun;137(6):726.e1-726.e18; discussion
AJODO	2010	726-7. doi: 10.1016/j.ajodo.2009.11.009.
		Liu L, Zhan Q, Zhou J, Kuang Q, Yan X, Zhang X, Shan Y, Lai W, Long H. A comparison of the
		effects of Forsus appliances with and without temporary anchorage devices for skeletal
		Class II malocclusion. Angle Orthod. 2021 Mar 1;91(2):255-266. doi: 10.2319/051120-
AO	2021	421.1. PMID: 33378419; PMCID: PMC8028478.
		Torres D, Lopes J, Magno MB, Cople Maia L, Normando D, Leão PB. Effects of rapid
		maxillary expansion on temporomandibular joints. Angle Orthod. 2020 May 1;90(3):442-
AO	2020	456. doi: 10.2319/080619-517.1. PMID: 33378434; PMCID: PMC8032308.
		Sosly R, Mohammed H, Rizk MZ, Jamous E, Qaisi AG, Bearn DR. Effectiveness of
		miniscrew-supported maxillary incisor intrusion in deep-bite correction: A systematic
		review and meta-analysis. Angle Orthod. 2020 Mar;90(2):291-304. doi: 10.2319/061119-
AO	2020	400.1. Epub 2019 Dec 9. PMID: 31816252; PMCID: PMC8051239.
		Vieira EP, Watanabe BSD, Pontes LF, Mattos JNF, Maia LC, Normando D. The effect of
		bracket slot size on the effectiveness of orthodontic treatment: A systematic review. Angle
AO	2018	Orthod. 2018 Jan;88(1):100-106. doi: 10.2319/031217-185.1.
		Mohamed RN, Basha S, Al-Thomali Y. Maxillary molar distalization with miniscrew-
	2010	supported appliances in Class II malocclusion: A systematic review.Angle Orthod. 2018
AO	2018	Jul;88(4):494-502. doi: 10.2319/091717-624.1.
		Ellabban MT, Abdul-Aziz AI, Fayed MMS, AboulFotouh MH, Elkattan ES, Dahaba MM.
		Positional and dimensional temporomandibular joint changes after correction of posterior
10	2010	crossbite in growing patients: A systematic review.Angle Orthod. 2018 Sep;88(5):638-648.
AO	2018	doi: 10.2319/110217-749.1.

		Xu Y, Xie J. Comparison of the effects of mini-implant and traditional anchorage on
	2017	patients with maxillary dentoalveolar protrusion. Angle Orthod. 2017 Mar;87(2):320-327.
AO	2017	doi: 10.2319/051016-375.1.
		Janson G, Aliaga-Del Castillo A, Niederberger A. Changes in apical base sagittal
		relationship in Class II malocclusion treatment with and without premolar extractions: A
	2017	systematic review and meta-analysis. Angle Orthod. 2017 Mar;87(2):338-355. doi:
AO	2017	10.2319/030716-198.1.
		Yi J, Xiao J, Li Y, Li X, Zhao Z. Efficacy of piezocision on accelerating orthodontic tooth
	2017	movement: A systematic review. Angle Orthod. 2017 Jul;87(4):491-498. doi:
AO	2017	10.2319/01191-751.1.
		Fagundes NCF, Rabello NM, Maia LC, Normando D, Mello KCFR. Can rapid maxillary
	2017	expansion cause auditory improvement in children and adolescents with hearing loss? A
AO	2017	systematic review. Angle Orthod. 2017 Nov;87(6):886-896. doi: 10.2319/021517-111.1.
		Perinetti G, Primožič J, Furlani G, Franchi L, Contardo L. Treatment effects of fixed
		functional appliances alone or in combination with multibracket appliances: A systematic
		review and meta-analysis. Angle Orthod. 2015 May;85(3):480-92. doi: 10.2319/102813-
AO	2015	790.1.
		Lopes Filho H, Maia LH, Lau TC, de Souza MM, Maia LC. Early vs late orthodontic
		treatment of tooth crowding by first premolar extraction: A systematic review. Angle
AO	2015	Orthod. 2015 May;85(3):510-7. doi: 10.2319/050814-332.1.
		Rossini G, Parrini S, Castroflorio T, Deregibus A, Debernardi CL. Efficacy of clear aligners in
		controlling orthodontic tooth movement: a systematic review.Angle Orthod. 2015
AO	2015	Sep;85(5):881-9. doi: 10.2319/061614-436.1.
		Feres MF, Raza H, Alhadlaq A, El-Bialy T. Rapid maxillary expansion effects in Class II
	2015	malocclusion: a systematic review.Angle Orthod. 2015 Nov;85(6):1070-9. doi:
AO	2015	10.2319/102514-768.1.
		Long H, Pyakurel U, Wang Y, Liao L, Zhou Y, Lai W. Interventions for accelerating
	2012	orthodontic tooth movement: a systematic review. Angle Orthod. 2013 Jan;83(1):164-71.
AO	2013	doi: 10.2319/031512-224.1.
		Flores-Mir C, McGrath L, Heo G, Major PW. Efficiency of molar distalization associated
10	2012	with second and third molar eruption stage. Angle Orthod. 2013 Jul;83(4):735-42. doi:
AO	2013	10.2319/081612-658.1.
		Bazargani F, Feldmann I, Bondemark L. Three-dimensional analysis of effects of rapid
10	2012	maxillary expansion on facial sutures and bones. Angle Orthod. 2013 Nov;83(6):1074-82.
AO	2013	doi: 10.2319/020413-103.1.
		Feng X, Li J, Li Y, Zhao Z, Zhao S, Wang J. Effectiveness of TAD-anchored maxillary
10	2012	protraction in late mixed dentition. Angle Orthod. 2012 Nov;82(6):1107-14. doi:
AO	2012	10.2319/111411-705.1.
		Liu ZP, Li CJ, Hu HK, Chen JW, Li F, Zou SJ. Efficacy of short-term chincup therapy for mandibular growth retardation in Class III malocclusion. Angle Orthod. 2011
10	2011	
AO	2011	Jan;81(1):162-68. doi: 10.2319/050510-244.1. Li F, Hu HK, Chen JW, Liu ZP, Li GF, He SS, Zou SJ, Ye QS. Comparison of anchorage capacity
۸ 0	2011	between implant and headgear during anterior segment retraction. Angle Orthod. 2011
AO	2011	Sep;81(5):915-22. doi: 10.2319/101410-603.1.
		Leonardi R, Annunziata A, Licciardello V, Barbato E. Soft tissue changes following the
10	2010	extraction of premolars in nongrowing patients with bimaxillary protrusion. A systematic
AO	2010	review. Angle Orthod. 2010 Jan;80(1):211-6. doi: 10.2319/010709-16.1.
10	2040	Fleming PS, Johal A. Self-ligating brackets in orthodontics. A systematic review. Angle
AO	2010	Orthod. 2010 May;80(3):575-84. doi: 10.2319/081009-454.1.
		Gordon JM, Rosenblatt M, Witmans M, Carey JP, Heo G, Major PW, Flores-Mir C. Rapid
	2000	palatal expansion effects on nasal airway dimensions as measured by acoustic rhinometry.
AO	2009	A systematic review. Angle Orthod. 2009 Sep;79(5):1000-7. doi: 10.2319/082108-441.1.
KIO	2020	Giudice AL, Spinuzza P, Rustico L, Messina G, Nucera R. Short-term treatment effects
KJO	2020	produced by rapid maxillary expansion evaluated with computed tomography: A

		systematic review with meta-analysis. Korean J Orthod. 2020 Sep 25;50(5):314-323. doi:
		10.4041/kjod.2020.50.5.314. PMID: 32938824; PMCID: PMC7500570.
		Galan-Lopez L, Barcia-Gonzalez J, Plasencia E. A systematic review of the accuracy and
		efficiency of dental movements with Invisalign. Korean J Orthod. 2019 May;49(3):140-149.
KJO	2019	doi: 10.4041/kjod.2019.49.3.140. Epub 2019 May 21.
		Viwattanatipa N, Charnchairerk S. The effectiveness of corticotomy and piezocision on
		canine retraction: A systematic review.Korean J Orthod. 2018 May;48(3):200-211. doi:
KJO	2018	10.4041/kjod.2018.48.3.200.
		Bayome M, Park JH, Bay C, Kook YA. Distalization of maxillary molars using temporary
		skeletal anchorage devices: A systematic review and meta-analysis. Orthod Craniofac Res.
O&CR	2021	2021 Mar;24 Suppl 1:103-112. doi: 10.1111/ocr.12470. Epub 2021 Feb 8.
		Arvind P TR, Jain RK. Skeletally anchored forsus fatigue resistant device for correction of
		Class II malocclusions-A systematic review and meta-analysis. Orthod Craniofac Res. 2021
O&CR	2021	Feb;24(1):52-61. doi: 10.1111/ocr.12414. Epub 2020 Sep 7. PMID: 32772479.
		MacDonald, L., Zanjir, M., Laghapour Lighvan, N., da Costa, B. R., Suri, S., & Azarpazhooh,
		A. (2020). Efficacy and Safety of Different Interventions to Accelerate Maxillary Canine
		Retraction Following Premolar Extraction: A Systematic Review and Network Meta-
O&CR	2021	analysis. Orthodontics & Craniofacial Research. doi:10.1111/ocr.12409
		Santana LG, Avelar K, Flores-Mir C, Marques LS. Incremental or maximal mandibular
		advancement in the treatment of class II malocclusion through functional appliances: A
		systematic review with meta-analysis. Orthod Craniofac Res. 2020 Nov;23(4):371-384. doi:
O&CR	2020	10.1111/ocr.12388. Epub 2020 May 29. PMID: 32390332.
		Zhang B, Huang X, Huo S, Zhang C, Zhao S, Cen X, Zhao Z. Effect of clear aligners on oral
		health-related quality of life: A systematic review. Orthod Craniofac Res. 2020
O&CR	2020	Nov;23(4):363-370. doi: 10.1111/ocr.12382. Epub 2020 May 13. PMID: 32340082.
		Niu X, Di Carlo G, Cornelis MA, Cattaneo PM. Three-dimensional analyses of short- and
		long-term effects of rapid maxillary expansion on nasal cavity and upper airway: A
		systematic review and meta-analysis. Orthod Craniofac Res. 2020 Aug;23(3):250-276. doi:
O&CR	2020	10.1111/ocr.12378. Epub 2020 May 5. PMID: 32248642.
		Robertson L, Kaur H, Fagundes NCF, Romanyk D, Major P, Flores Mir C. Effectiveness of
		clear aligner therapy for orthodontic treatment: A systematic review. Orthod Craniofac
		Res. 2020 May;23(2):133-142. doi: 10.1111/ocr.12353. Epub 2019 Nov 13. PMID:
O&CR	2020	31651082.
		Dab S, Chen K, Flores-Mir C. Short- and long-term potential effects of accelerated
		osteogenic orthodontic treatment: A systematic review and meta-analysis. Orthod
O&CR	2019	Craniofac Res. 2019 May;22(2):61-68. doi: 10.1111/ocr.12272. Epub 2019 Mar 18.
		Lee WC, Tu YK2, Huang CS, Chen R, Fu MW, Fu E. Pharyngeal airway changes following
		maxillary expansion or protraction: A meta-analysis.Orthod Craniofac Res. 2018
O&CR	2018	Feb;21(1):4-11. doi: 10.1111/ocr.12208.
		Mohammed H, Rizk MZ, Wafaie K, Almuzian M. Effectiveness of nickel-titanium springs vs
		elastomeric chains in orthodontic space closure: A systematic review and meta-
O&CR	2018	analysis.Orthod Craniofac Res. 2018 Feb;21(1):12-19. doi: 10.1111/ocr.12210.
		Koletsi D, Makou M, Pandis N. Effect of orthodontic management and orofacial muscle
		training protocols on the correction of myofunctional and myoskeletal problems in
		developing dentition. A systematic review and meta-analysis.Orthod Craniofac Res. 2018
O&CR	2018	Nov;21(4):202-215. doi: 10.1111/ocr.12240.
		Zheng M, Liu R, Ni Z, Yu Z. Efficiency, effectiveness and treatment stability of clear
		aligners: A systematic review and meta-analysis. Orthod Craniofac Res. 2017
O&CR	2017	Aug;20(3):127-133. doi: 10.1111/ocr.12177.
		Cordasco G, Matarese G, Rustico L, Fastuca S, Caprioglio A, Lindauer SJ, Nucera R. Efficacy
		of orthopedic treatment with protraction facemask on skeletal Class III malocclusion: a
		systematic review and meta-analysis. Orthod Craniofac Res. 2014 Aug;17(3):133-43. doi:
O&CR	2014	10.1111/ocr.12040.
O&CR	2014	10.1111/ocr.12040. Papageorgiou SN, Konstantinidis I, Papadopoulou K, Jäger A, Bourauel C. A systematic

and archwire sequences. Orthod Craniofac Res. 2014 Nov;17(4):197-215. doi:
10.1111/ocr.12048.

*Description of the abbreviated journals: Cochrane library: Cochrane Database of Systematic Reviews AJODO: American Journal of Orthodontics and Dentofacial Orthopedics EJO: European Journal of Orthodontics AO: Angle Orthodontist KJO: Korean Journal of Orthodontics O&CR: Orthodontics and Craniofacial Research

Additional file 4. Excluded studies

Update

The following reference from 2017: Arora A, Khattri S, Ismail NM, Kumbargere Nagraj S, Prashanti E. School dental screening programmes for oral health.Cochrane Database Syst Rev. 2017 Dec 21;12:CD012595. doi: 10.1002/14651858.CD012595.pub2. was excluded prior to title and abstract screening, because it was an update of the following reference by the same authors in 2019: Arora A, Khattri S, Ismail NM, Kumbargere Nagraj S, Eachempati P. School dental screening programmes for oral health. Cochrane Database Syst Rev. 2019 Aug 8;8:CD012595. doi: 10.1002/14651858.CD012595.pub3.

Rationale for exclusion of studies

A total of 180 studies were excluded during the title and abstract screening and 45 were excluded during full text screening. The rationales for exclusion were given for each study. Only one rationale was given per study even when more than one rationales could have been applied.

Rationales for exclusion of studies:

- Assessed exclusively adverse effects
- Not an orthodontic intervention
- Not the effects of orthodontic interventions were assessed
- Empty review
- review was later updated
- Review of animal studies
- Review of laboratory studies
- Review included orthognathic surgical interventions
- Assessed exclusively patients with congenital anomalies
- Review did not assess the effect of a specific type of intervention(s), but assessed an undefined orthodontic intervention, e.g., orthodontic treatment as a whole
- Review was conducted by one operator only
- The review is about a specific outcome of an intervention, which is ambiguous and could also be an adverse effect
- A Bayesian network meta-analysis was used

Excluded studies during the title and abstract screening (n=180) with the rationale for exclusion

Journal	*	Year	Reference	Rationale for exclusion
Journal		Tear	Mulimani P, Abas AB, Karanth L, Colombatti R,	exclusion
			Kulkarni P. Treatment of dental and orthodontic	
			complications in thalassaemia. Cochrane Database	assessed
1.	Cochrane		Syst Rev. 2019 Aug 2;8:CD012969. doi:	exclusively
1.	library	2019	10.1002/14651858.CD012969.pub2.	adverse effects
	library	2015	Arora A, Khattri S, Ismail NM, Kumbargere Nagraj S,	
			Eachempati P. School dental screening programmes	not the effects of
			for oral health. Cochrane Database Syst Rev. 2019	orthodontic
2	Cochrane		Aug 8;8:CD012595. doi:	interventions
2.	library	2019	10.1002/14651858.CD012595.pub3.	were assessed
	library	2015	Millett DT, Cunningham SJ, O'Brien KD, Benson PE,	Were assessed
			de Oliveira CM. Orthodontic treatment for deep	
			bite and retroclined upper front teeth in	
			children.Cochrane Database Syst Rev. 2018 Feb	
3.	Cochrane		1;2:CD005972. doi:	
0.	library	2018	10.1002/14651858.CD005972.pub4.	empty review
		2010	Mandall NA, Hickman J, Macfarlane TV, Mattick RC,	empty renew
			Millett DT, Worthington HV. Adhesives for fixed	not the effects of
			orthodontic brackets.Cochrane Database Syst Rev.	orthodontic
4	Cochrane		2018 Apr 9;4:CD002282. doi:	interventions
	library	2018	10.1002/14651858.CD002282.pub2.	were assessed
	library	2010	Millett DT, Mandall NA, Mattick RC, Hickman J,	Were assessed
			Glenny AM. Adhesives for bonded molar tubes	not the effects of
			during fixed brace treatment. Cochrane Database	orthodontic
5.	Cochrane		Syst Rev. 2017 Feb 23;2:CD008236. doi:	interventions
5.	library	2017	10.1002/14651858.CD008236.pub3.	were assessed
	library	2017	Agnihotry A, Fedorowicz Z, Nasser M, Gill KS.	Were assessed
			Resorbable versus titanium plates for orthognathic	not the effects of
			surgery.Cochrane Database Syst Rev. 2017 Oct	orthodontic
6.	Cochrane		4;10:CD006204. doi:	interventions
	library	2017	10.1002/14651858.CD006204.pub3.	were assessed
			Monk AB, Harrison JE, Worthington HV, Teague A.	
			Pharmacological interventions for pain relief during	not the effects of
			orthodontic treatment. Cochrane Database Syst	orthodontic
7.	Cochrane		Rev. 2017 Nov 28;11:CD003976. doi:	interventions
	library	2017	10.1002/14651858.CD003976.pub2.	were assessed
			Ashley PF, Parekh S, Moles DR, Anand P,	
			MacDonald LC. Preoperative analgesics for	
			additional pain relief in children and adolescents	not the effects of
			having dental treatment. Cochrane Database Syst	orthodontic
8.	Cochrane		Rev. 2016 Aug 8;(8):CD008392. doi:	interventions
	library	2016	10.1002/14651858.CD008392.pub3.	were assessed
	1		Ghaeminia H, Perry J, Nienhuijs ME, Toedtling V,	
			Tummers M, Hoppenreijs TJ, Van der Sanden WJ,	
			Mettes TG. Surgical removal versus retention for	
			the management of asymptomatic disease-free	not the effects of
			impacted wisdom teeth. Cochrane Database Syst	orthodontic
9.	Cochrane		Rev. 2016 Aug 31;(8):CD003879. doi:	interventions
	library	2016	10.1002/14651858.CD003879.pub4.	were assessed
10.	Cochrane		Carvalho FR, Lentini-Oliveira DA, Prado LB, Prado	not the effects of
= 5.	library	2016	GF, Carvalho LB. Oral appliances and functional	orthodontic

		orthopaedic appliances for obstructive sleep	interventions
		apnoea in children. Cochrane Database Syst Rev.	were assessed
		2016 Oct 5;10:CD005520.	
		Millett DT, Glenny AM, Mattick RC, Hickman J,	not the effects of
		Mandall NA. Adhesives for fixed orthodontic bands.	orthodontic
11. Cochrane		Cochrane Database Syst Rev. 2016 Oct	interventions
library	2016	25;10:CD004485.	were assessed
		Fleming PS, Strydom H, Katsaros C, MacDonald L,	
		Curatolo M, Fudalej P, Pandis N. Non-	
		pharmacological interventions for alleviating pain	not the effects of
		during orthodontic treatment. Cochrane Database	orthodontic
12. Cochrane		Syst Rev. 2016 Dec 23;12:CD010263. doi:	interventions
library	2016	10.1002/14651858.CD010263.pub2.	were assessed
		Ahangari Z, Nasser M, Mahdian M, Fedorowicz Z,	
		Marchesan MA. Interventions for the management	
		of external root resorption. Cochrane Database Syst	
		Rev. 2015 Nov 24;(11):CD008003. doi:	assessed
13. Cochrane		10.1002/14651858.CD008003.pub3.(discuss met	exclusively
library	2015	Reint: moeten we deze wel includeren)	adverse effects
		Belmonte FM, Macedo CR, Day PF, Saconato H,	
		Fernandes Moça Trevisani V. Interventions for	
		treating traumatised permanent front teeth:	
		luxated (dislodged) teeth.Cochrane Database Syst	
14. Cochrane		Rev. 2013 Apr 30;(4):CD006203. doi:	
library	2013	10.1002/14651858.CD006203.pub2.	empty review
		Yu Y, Sun J, Lai W, Wu T, Koshy S, Shi Z.	
		Interventions for managing relapse of the lower	
		front teeth after orthodontic treatment. Cochrane	
15. Cochrane		Database Syst Rev. 2013 Sep 6;(9):CD008734. doi:	
library	2013	10.1002/14651858.CD008734.pub2.	empty review
		Hu H, Li C, Li F, Chen J, Sun J, Zou S, Sandham A, Xu	
		Q, Riley P, Ye Q. Enamel etching for bonding fixed	not the effects of
		orthodontic braces. Cochrane Database Syst Rev.	orthodontic
16. Cochrane		2013 Nov 25;(11):CD005516. doi:	interventions
library	2013	10.1002/14651858.CD005516.pub2.	were assessed
		Benson PE, Parkin N, Dyer F, Millett DT, Furness S,	
		Germain P. Fluorides for the prevention of early	
		tooth decay (demineralised white lesions) during	not the effects of
		fixed brace treatment. Cochrane Database Syst Rev.	orthodontic
17. Cochrane		2013 Dec 12;(12):CD003809. doi:	interventions
library	2013	10.1002/14651858.CD003809.pub3.	were assessed
		Alrashed M, Alqerban A. The relationship between	
		malocclusion and oral health-related quality of life	
		among adolescents: a systematic literature review	not the effects of
		and meta-analysis. Eur J Orthod. 2021 Apr	orthodontic
		3;43(2):173-183. doi: 10.1093/ejo/cjaa051. PMID:	interventions
18. EJO	2021	33009547.	were assessed
		Mulier D, Gaitán Romero L, Führer A, Martin C,	
		Shujaat S, Shaheen E, Politis C, Jacobs R. Long-term	
		dental stability after orthognathic surgery: a	review included
		systematic review. Eur J Orthod. 2021 Jan	orthognathic
		29;43(1):104-112. doi: 10.1093/ejo/cjaa022. PMID:	surgical
19. EJO	2021	32901268.	interventions
		Palikaraki G, Makrygiannakis MA, Zafeiriadis AA,	assessed
		Benetou V, Sanoudos M, Bitsanis I, Tsolakis AI. The	exclusively
	2021	effect of facemask in patients with unilateral cleft	patients with

		lip and palate: a systematic review and meta-	congenital
		analysis. Eur J Orthod. 2021 Jan 29;43(1):69-79. doi:	anomalies
		10.1093/ejo/cjaa027. PMID: 32274494.	
		Gandhi V, Mehta S, Gauthier M, Mu J, Kuo CL,	
		Nanda R, Yadav S. Comparison of external apical	
		root resorption with clear aligners and pre-adjusted	
		edgewise appliances in non-extraction cases: a	
		systematic review and meta-analysis. Eur J Orthod.	assessed
		2021 Jan 29;43(1):15-24. doi: 10.1093/ejo/cjaa013.	exclusively
21. EJO	2021	PMID: 32077935; PMCID: PMC7846172.	adverse effects
		Stucki S, Gkantidis N. Assessment of techniques	
		used for superimposition of maxillary and	
		mandibular 3D surface models to evaluate tooth	
		movement: a systematic review. Eur J Orthod. 2020	not an
		Nov 3;42(5):559-570. doi: 10.1093/ejo/cjz075.	orthodontic
22. EJO	2020	PMID: 31742598.	intervention
		Rodrigues AS, Antunes LS, Pinheiro LHM, Guimarães	
		LS, Calansans-Maia JA, Küchler EC, Antunes LAA. Is	
		dental agenesis associated with craniofacial	
		morphology pattern? A systematic review and	not an
		meta-analysis. Eur J Orthod. 2020 Nov 3;42(5):534-	orthodontic
23. EJO	2020	543. doi: 10.1093/ejo/cjz087. PMID: 31783403.	intervention
		Kaklamanos EG, Makrygiannakis MA, Athanasiou	
		AE. Does medication administration affect the rate	
		of orthodontic tooth movement and root	not the effects of
		resorption development in humans? A systematic	orthodontic
		review. Eur J Orthod. 2020 Sep 11;42(4):407-414.	interventions
24. EJO	2020	doi: 10.1093/ejo/cjz063. PMID: 31421637.	were assessed
		de Araujo CM, Schroder AGD, de Araujo BMM,	
		Cavalcante-Leão BL, Stechman-Neto J, Zeigelboim	
		BS, Santos RS, Guariza-Filho O. Impact of	review included
		orthodontic-surgical treatment on quality of life: a	orthognathic
		meta-analysis. Eur J Orthod. 2020 Jun 23;42(3):281-	surgical
25. EJO	2020	289. doi: 10.1093/ejo/cjz093. PMID: 31784741.	interventions
		Papadimitriou A, Kakali L, Pazera P, Doulis I, Kloukos	
		D. Social media and orthodontic treatment from	not the effects of
		the patient's perspective: a systematic review. Eur J	orthodontic
		Orthod. 2020 Jun 23;42(3):231-241. doi:	interventions
26. EJO	2020	10.1093/ejo/cjz029. PMID: 31107943.	were assessed
		Zymperdikas VF, Yavropoulou MP, Kaklamanos EG,	
		Papadopoulos MA. Effects of systematic	
		bisphosphonate use in patients under orthodontic	
		treatment: a systematic review. Eur J Orthod. 2020	not an
		Jan 27;42(1):60-71. doi: 10.1093/ejo/cjz021. PMID:	orthodontic
27. EJO	2020	31009953.	intervention
		Sardana D, Manchanda S, Ekambaram M, Yang Y,	
		McGrath CP, Yiu CKY. Effectiveness of self-applied	
		topical fluorides against enamel white spot lesions	
		from multi-bracketed fixed orthodontic treatment:	
		a systematic review. Eur J Orthod. 2019 Nov	
		15;41(6):661-668. doi: 10.1093/ejo/cjz015. Erratum	not an
		in: Eur J Orthod. 2019 Nov 15;41(6):669. PMID:	orthodontic
28. EJO	2019	31112229.	intervention
		Makrygiannakis MA, Kaklamanos EG, Athanasiou	
		AE. Does long-term use of pain relievers have an	review of animal
29. EJO	2019	impact on the rate of orthodontic tooth	studies
· · · · •			

	-		
		movement? A systematic review of animal studies.	
		Eur J Orthod. 2019 Sep 21;41(5):468-477. doi:	
		10.1093/ejo/cjy079. PMID: 30590549.	
		Kakali L, Alharbi M, Pandis N, Gkantidis N, Kloukos	
		D. Success of palatal implants or mini-screws placed	
		median or paramedian for the reinforcement of	not the effects of
		anchorage during orthodontic treatment: a	orthodontic
		systematic review. Eur J Orthod. 2019 Jan	interventions
30. EJO	2019	23;41(1):9-20. doi: 10.1093/ejo/cjy015.	were assessed
		Samandara A, Papageorgiou SN, Ioannidou-	
		Marathiotou I, Kavvadia-Tsatala S, Papadopoulos	
		MA.	
		Evaluation of orthodontically induced external root	
		resorption following orthodontic treatment using	
		cone beam computed tomography (CBCT): a	assessed
		systematic review and meta-analysis. Eur J Orthod.	exclusively
31. EJO	2019	2019 Jan 23;41(1):67-79. doi: 10.1093/ejo/cjy027.	adverse effects
		Bartolucci ML, Bortolotti F, Martina S2, Corazza G,	
		Michelotti A, Alessandri-Bonetti G. Dental and	
	1	skeletal long-term side effects of mandibular	
		advancement devices in obstructive sleep apnea	not the effects of
		patients: a systematic review with meta-regression	orthodontic
		analysis. Eur J Orthod. 2019 Jan 23;41(1):89-100.	interventions
32. EJO	2019	doi: 10.1093/ejo/cjy036.	were assessed
		Mohammed H, Rizk MZ, Wafaie K, Ulhaq A,	
		Almuzian M. Reminders improve oral hygiene and	
		adherence to appointments in orthodontic	not the effects of
		patients: a systematic review and meta-analysis.	orthodontic
		Eur J Orthod. 2019 Mar 29;41(2):204-213. doi:	interventions
33. EJO	2019	10.1093/ejo/cjy045.	were assessed
00. 200	2015	Makrygiannakis MA, Kaklamanos EG, Athanasiou	incre abbebbeu
		AE. Effects of systemic medication on root	
		resorption associated with orthodontic tooth	
		movement: a systematic review of animal studies.	
		Eur J Orthod. 2019 Aug 8;41(4):346-359. doi:	review of animal
34. EJO	2019	10.1093/ejo/cjy048.	studies
		De Grauwe A, Ayaz I, Shujaat S, Dimitrov S,	
		Gbadegbegnon L, Vande Vannet B, Jacobs R.	
		CBCT in orthodontics: a systematic review on	not the effects of
	1	justification of CBCT in a paediatric population prior	orthodontic
		to orthodontic treatment. Eur J Orthod. 2019 Aug	interventions
35. EJO	2019	8;41(4):381-389. doi: 10.1093/ejo/cjy066.	were assessed
		Chen J, Wan J, You L. Speech and orthodontic	
		appliances: a systematic literature review.Eur J	assessed
		Orthod. 2018 Jan 23;40(1):29-36. doi:	exclusively
36. EJO	2018	10.1093/ejo/cjx023.	adverse effects
-		Alyammahi AS, Kaklamanos EG, Athanasiou AE.	
		Effectiveness of extraction of primary canines for	
		interceptive management of palatally displaced	not the effects of
		permanent canines: a systematic review and meta-	orthodontic
		analysis.Eur J Orthod. 2018 Apr 6;40(2):149-156.	interventions
37. EJO	2018	doi: 10.1093/ejo/cjx042.	were assessed
-		Lo Giudice A, Barbato E, Cosentino L, Ferraro CM,	assessed
		Leonardi R Alveolar bone changes after rapid	exclusively
38. EJO	2018	maxillary expansion with tooth-born appliances: a	adverse effects

		systematic review.Eur J Orthod. 2018 May	
		25;40(3):296-303. doi: 10.1093/ejo/cjx057.	
		Haugland L, Kristensen KD, Lie SA, Vandevska-	
		Radunovic V. The effect of biologic factors and	
		adjunctive therapies on orthodontically induced	
		inflammatory root resorption: a systematic review	assessed
		and meta-analysis.Eur J Orthod. 2018 May	exclusively
39. EJO	2018	25;40(3):326-336. doi: 10.1093/ejo/cjy003.	adverse effects
	2010	Alharbi F, Almuzian M, Bearn D. Miniscrews failure	
		rate in orthodontics: systematic review and meta-	assessed
		analysis.Eur J Orthod. 2018 Sep 28;40(5):519-530.	exclusively
40. EJO	2018	doi: 10.1093/ejo/cjx093.	adverse effects
		Grisar K, Chaabouni D, Romero LPG,	
		Vandendriessche T, Politis C, Jacobs R. Autogenous	
		transalveolar transplantation of maxillary canines: a	
		systematic review and meta-analysis.Eur J Orthod.	not an
		2018 Nov 30;40(6):608-616. doi:	orthodontic
41. EJO	2018	10.1093/ejo/cjy026.	intervention
		Dumbryte I, Vebriene J, Linkeviciene L, Malinauskas	
		M. Enamel microcracks in the form of tooth	
		damage during orthodontic debonding: a	
		systematic review and meta-analysis of in vitro	review of
		studies.Eur J Orthod. 2018 Nov 30;40(6):636-648.	laboratory
42. EJO	2018	doi: 10.1093/ejo/cjx102.	studies
		Makrygiannakis MA, Kaklamanos EG, Athanasiou	
		AE. Does common prescription medication affect	
		the rate of orthodontic tooth movement? A	
		systematic review. Eur J Orthod. 2018 Nov	review of animal
43. EJO	2018	30;40(6):649-659. doi: 10.1093/ejo/cjy001.	studies
		Castroflorio T, Bargellini A, Rossini G, Cugliari G,	
		Deregibus A. Sleep bruxism in adolescents: a	
		systematic literature review of related risk factors.	not an
44. EJO	2017	Eur J Orthod. 2017 Feb;39(1):61-68. doi: 10.1093/ejo/cjw012.	orthodontic intervention
44. EJU	2017	Höchli D, Hersberger-Zurfluh M, Papageorgiou SN,	Intervention
		Eliades T. Interventions for orthodontically induced	
		white spot lesions: a systematic review and meta-	not an
		analysis. Eur J Orthod. 2017 Apr 1;39(2):122-133.	orthodontic
45. EJO	2017	doi: 10.1093/ejo/cjw065.	intervention
		Sonesson M, Bergstrand F, Gizani S, Twetman S.	
		Management of post-orthodontic white spot	
		lesions: an updated systematic review. Eur J	not an
		Orthod. 2017 Apr 1;39(2):116-121. doi:	orthodontic
46. EJO	2017	10.1093/ejo/cjw023.	intervention
		Dalessandri D, Parrini S, Rubiano R, Gallone D,	
		Migliorati M. Impacted and transmigrant	
		mandibular canines incidence, aetiology, and	not an
		treatment: a systematic review. Eur J Orthod. 2017	orthodontic
47. EJO	2017	Apr 1;39(2):161-169. doi: 10.1093/ejo/cjw027.	intervention
		Yi J, Ge M, Li M, Li C, Li Y, Li X, Zhao Z. Comparison	
		of the success rate between self-drilling and self-	not the effects of
		tapping miniscrews: a systematic review and meta-	orthodontic
		analysis. Eur J Orthod. 2017 Jun 1;39(3):287-293.	interventions
48. EJO	2017	doi: 10.1093/ejo/cjw036.	were assessed

	r		
		Mousoulea S, Kloukos D, Sampaziotis D, Vogiatzi T,	
		Eliades T. Condylar resorption in orthognathic	
		patients after mandibular bilateral sagittal split	assessed
40.510	2017	osteotomy: a systematic review. Eur J Orthod. 2017	exclusively
49. EJO	2017	Jun 1;39(3):294-309. doi: 10.1093/ejo/cjw045.	adverse effects
		Zimmerman JN, Lee J, Pliska BT. Reliability of upper	
		pharyngeal airway assessment using dental CBCT: a	not an
		systematic review. Eur J Orthod. 2017 Oct	orthodontic
50. EJO	2017	1;39(5):489-496. doi: 10.1093/ejo/cjw079.	intervention
		Becking BE, Verweij JP, Kalf-Scholte SM, Valkenburg	
		C, Bakker EWP, van Merkesteyn JPR. Impact of	
		adenotonsillectomy on the dentofacial	
		development of obstructed children: a systematic	not an
		review and meta-analysis. Eur J Orthod. 2017 Oct	orthodontic
51. EJO	2017	1;39(5):509-518. doi: 10.1093/ejo/cjx005.	intervention
		Sandhu SS, Piepho HP, Khehra HS. Comparing the	
		effectiveness profile of pharmacological	
		interventions used for orthodontic pain relief: an	
		arm-based multilevel network meta-analysis of	not an
		longitudinal data. Eur J Orthod. 2017 Nov	orthodontic
52. EJO	2017	30;39(6):601-614. doi: 10.1093/ejo/cjw088.	intervention
		Rakhshan V, Rakhshan H. Meta-analysis and	
		systematic review of the number of non-syndromic	
		congenitally missing permanent teeth per affected	not an
		individual and its influencing factors. Eur J Orthod.	orthodontic
53. EJO	2016	2016 Apr;38(2):170-7. doi: 10.1093/ejo/cjv008.	intervention
		Goracci C, Franchi L, Vichi A, Ferrari M3. Accuracy,	
		reliability, and efficiency of intraoral scanners for	
		full-arch impressions: a systematic review of the	not an
		clinical evidence. Eur J Orthod. 2016 Aug;38(4):422-	orthodontic
54. EJO	2016	8. doi: 10.1093/ejo/cjv077.	intervention
		Aragón ML, Pontes LF, Bichara LM, Flores-Mir C,	
		Normando D. Validity and reliability of intraoral	
		scanners compared to conventional gypsum models	
		measurements: a systematic review. Eur J Orthod.	not an
		2016 Aug;38(4):429-34. doi: 10.1093/ejo/cjw033.	orthodontic
55. EJO	2016	(geen PDF)	intervention
		lodice G, Danzi G, Cimino R, Paduano S, Michelotti	
		A. Association between posterior crossbite,	not an
		skeletal, and muscle asymmetry: a systematic	orthodontic
56. EJO	2016	review. Eur J Orthod. 2016 Dec;38(6):638-651.	intervention
		Dimberg L, Arnrup K, Bondemark L. The impact of	
		malocclusion on the quality of life among children	
		and adolescents: a systematic review of	not an
		quantitative studies. Eur J Orthod. 2015	orthodontic
57. EJO	2015	Jun;37(3):238-47. doi: 10.1093/ejo/cju046.	intervention
		Migliorati M, Isaia L, Cassaro A, Rivetti A, Silvestrini-	
		Biavati F, Gastaldo L, Piccardo I, Dalessandri D,	
		Silvestrini-Biavati A. Efficacy of professional hygiene	
		and prophylaxis on preventing plaque increase in	
		orthodontic patients with multibracket appliances:	not an
		a systematic review. Eur J Orthod. 2015	orthodontic
58. EJO	2015	Jun;37(3):297-307. doi: 10.1093/ejo/cju044.	intervention
50. 250	2015	Nazarali N, Altalibi M, Nazarali S, Major MP, Flores-	not an
		Mir C, Major PW. Mandibular advancement	orthodontic
59 EIO	2015	appliances for the treatment of paediatric	intervention
59. EJO	2013	appliances for the treatment of paeulatific	Intervention

		obstructive sleep apnea: a systematic review. Eur J	
		Orthod. 2015 Dec;37(6):618-26. doi:	
		10.1093/ejo/cju101.	
		Gomes Lde C, Horta KO, Gonçalves JR, Santos-Pinto	
		AD. Systematic review: craniocervical posture and	not an
		craniofacial morphology. Eur J Orthod. 2014	orthodontic
60. EJO	2014	Feb;36(1):55-66. doi: 10.1093/ejo/cjt004.	intervention
		Dalessandri D, Salgarello S, Dalessandri M,	
		Lazzaroni E, Piancino M, Paganelli C, Maiorana C,	
		Santoro F. Determinants for success rates of	not the effects of
		temporary anchorage devices in orthodontics: a	orthodontic
		meta-analysis (n > 50). Eur J Orthod. 2014	interventions
61. EJO	2014	Jun;36(3):303-13. doi: 10.1093/ejo/cjt049.	were assessed
	-	Kloukos D, Eliades T, Sculean A, Katsaros C.	
		Indication and timing of soft tissue augmentation at	
		maxillary and mandibular incisors in orthodontic	not an
		patients. A systematic review.Eur J Orthod. 2014	orthodontic
62 510	2014		intervention
62. EJO	2014	Aug;36(4):442-9. doi: 10.1093/ejo/cjt073.	
		Winsauer H, Vlachojannis C, Bumann A,	
		Vlachojannis J, Chrubasik S. Paramedian vertical	
		palatal bone height for mini-implant insertion: a	not an
60 FIF		systematic review. Eur J Orthod. 2014	orthodontic
63. EJO	2014	Oct;36(5):541-9. doi: 10.1093/ejo/cjs068.	intervention
		Forst D, Nijjar S, Khaled Y, Lagravere M, Flores-Mir	
		C. Radiographic assessment of external root	
		resorption associated with jackscrew-based	
		maxillary expansion therapies: a systematic review.	assessed
		Eur J Orthod. 2014 Oct;36(5):576-85. doi:	exclusively
64. EJO	2014	10.1093/ejo/cjt090.	adverse effects
		Livas C, Delli K. Subjective and objective perception	
		of orthodontic treatment need: a systematic	not an
		review.Eur J Orthod. 2013 Jun;35(3):347-53. doi:	orthodontic
65. EJO	2013	10.1093/ejo/cjr142.	intervention
		Alves M Jr, Baratieri C, Mattos CT, Araújo MT, Maia	
		LC. Root repair after contact with mini-implants:	
		systematic review of the literature.Eur J Orthod.	not an
		2013 Aug;35(4):491-9. doi: 10.1093/ejo/cjs025.	orthodontic
66. EJO	2013	Epub 2012 Apr 26.	intervention
		Perinetti G, Primozic J, Manfredini D, Di Lenarda R,	
		Contardo L. The diagnostic potential of static body-	
		sway recording in orthodontics: a systematic	not an
		review. Eur J Orthod. 2013 Oct;35(5):696-705. doi:	orthodontic
67 510	2012		intervention
67. EJO	2013	10.1093/ejo/cjs085.	intervention
		Iodice G, Danzi G, Cimino R, Paduano S, Michelotti	
		A. Association between posterior crossbite,	
		masticatory muscle pain, and disc displacement: a	not an
		systematic review. Eur J Orthod. 2013	orthodontic
68. EJO	2013	Dec;35(6):737-44. doi: 10.1093/ejo/cjt024.	intervention
		Altalibi M, Saltaji H, Edwards R, Major PW, Flores-	
		Mir C. Indices to assess malocclusions in patients	not an
		with cleft lip and palate. Eur J Orthod. 2013	orthodontic
69. EJO	2013	Dec;35(6):772-82. doi: 10.1093/ejo/cjt009.	intervention
		Walker SL, Tieu LD, Flores-Mir C. Radiographic	assessed
		comparison of the extent of orthodontically	exclusively
70. EJO	2013	comparison of the extent of orthodontically induced external apical root resorption in vital and	exclusively adverse effects

	1	reat filled teath, a sustainatic review. Fur I Orthad	
		root-filled teeth: a systematic review. Eur J Orthod. 2013 Dec;35(6):796-802. doi: 10.1093/ejo/cjs101.	
		Fudalej P, Dragan M, Wedrychowska-Szulc B.	
		Prediction of the outcome of orthodontic treatment	
		of Class III malocclusionsa systematic review. Eur J	not an
74 510	2011	Orthod. 2011 Apr;33(2):190-7. doi:	orthodontic
71. EJO	2011	10.1093/ejo/cjq052.	intervention
		Ahrens A, McGrath C, Hägg U. A systematic review	
		of the efficacy of oral appliance design in the	
		management of obstructive sleep apnoea. Eur J	not an
		Orthod. 2011 Jun;33(3):318-24. doi:	orthodontic
72. EJO	2011	10.1093/ejo/cjq079.	intervention
		Malekshoar M, Malekshoar M, Javanshir B.	
		Challenges, limitations, and solutions for	
		orthodontists during the coronavirus pandemic: A	
		review. Am J Orthod Dentofacial Orthop. 2021	not the effects of
		Jan;159(1):e59-e71. doi:	orthodontic
	1	10.1016/j.ajodo.2020.09.009. Epub 2020 Oct 19.	interventions
73. AJODO	2021	PMID: 33223376; PMCID: PMC7571895.	were assessed
73. AJUDU	2021	Tepedino M, Cattaneo PM, Niu X, Cornelis MA.	************************
		Interradicular sites and cortical bone thickness for	
		miniscrew insertion: A systematic review with	
		meta-analysis. Am J Orthod Dentofacial Orthop.	
		<i>,</i> , , , , , , , , , , , , , , , , , ,	
		2020 Dec;158(6):783-798.e20. doi:	not the effects of
		10.1016/j.ajodo.2020.05.011. Epub 2020 Oct 16.	orthodontic
		Erratum in: Am J Orthod Dentofacial Orthop. 2021	interventions
74. AJODO	2020	Jun;159(6):711. PMID: 33077369.	were assessed
		ElShehaby M, Mofti B, Montasser MA, Bearn D.	
		Powered vs manual tooth brushing in patients with	
		fixed orthodontic appliances: A systematic review	
		and meta-analysis. Am J Orthod Dentofacial	
		Orthop. 2020 Nov;158(5):639-649. doi:	not an
		10.1016/j.ajodo.2020.04.018. Epub 2020 Sep 17.	orthodontic
75. AJODO	2020	PMID: 32951930.	intervention
			review did not
			assess the effect
			of a specific type
			of
	1		intervention(s),
			but assessed an
	1	Macey R, Thiruvenkatachari B, O'Brien K, Batista	undefined
		KBSL. Do malocclusion and orthodontic treatment	orthodontic
	1	impact oral health? A systematic review and meta-	intervention, e.g.,
		analysis. Am J Orthod Dentofacial Orthop. 2020	orthodontic
	1	Jun;157(6):738-744.e10. doi:	treatment as a
76. AJODO	2020	10.1016/j.ajodo.2020.01.015. PMID: 32487303.	whole
	2020	Arn ML, Dritsas K, Pandis N, Kloukos D. The effects	
	1	of fixed orthodontic retainers on periodontal	
	1	health: A systematic review. Am J Orthod	assessed
	1	Dentofacial Orthop. 2020 Feb;157(2):156-164.e17.	
	2020		exclusively
77. AJODO	2020	doi: 10.1016/j.ajodo.2019.10.010. PMID: 32005466.	adverse effects
		Vásquez-Cárdenas J, Zapata-Noreña Ó, Carvajal-	
		Flórez Á, Barbosa-Liz DM, Giannakopoulos NN,	
	1	Faggion CM Jr. Systematic reviews in orthodontics:	not an
		I Impost at the DDICNAA for Abstracts checklist on	orthodontic
78. AJODO	2019	Impact of the PRISMA for Abstracts checklist on completeness of reporting. Am J Orthod Dentofacial	intervention

	1		
	1	Orthop. 2019 Oct;156(4):442-452.e12. doi:	
		10.1016/j.ajodo.2019.05.009. PMID: 31582116.	
		Barretto Dos Santos Lopes Batista K,	
		Thiruvenkatachari B, O'Brien K. Intention-to-treat analysis: Are we managing dropouts and missing	
		data properly in research on orthodontic	
			not on
		treatment? A systematic review. Am J Orthod	not an
70 41000	2010	Dentofacial Orthop. 2019 Jan;155(1):19-27.e3. doi:	orthodontic
79. AJODO	2019	10.1016/j.ajodo.2018.08.013. Currell SD, Liaw A, Blackmore Grant PD, Esterman	intervention
		A4, Nimmo A. Orthodontic mechanotherapies and	
		their influence on external root resorption: A systematic review. Am J Orthod Dentofacial Orthop.	accord
		, , , , , , , , , , , , , , , , , , , ,	assessed
80 41000	2010	2019 Mar;155(3):313-329. doi:	exclusively
80. AJODO	2019	10.1016/j.ajodo.2018.10.015.	adverse effects
		Yi J, Lu W, Xiao J, Li X, Li Y, Zhao Z. Effect of	
	1	conventional combined orthodontic-surgical	المعادية والمعاد
	1	treatment on oral health-related quality of life: A	review included
		systematic review and meta-analysis. Am J Orthod	orthognathic
01 41050	2010	Dentofacial Orthop. 2019 Jul;156(1):29-43.e5. doi:	surgical
81. AJODO	2019	10.1016/j.ajodo.2019.03.008.	interventions
	1	Al-Jewair T, Stellrecht E, Lewandowski L, Chakaki R. American Association of Orthodontists Foundation	
		Craniofacial Growth Legacy Collection in the	
		orthodontic literature-use and trends: A systematic	
		review.Am J Orthod Dentofacial Orthop. 2018	not an
02 41000	2010	Jan;153(1):15-25.e10. doi:	orthodontic
82. AJODO	2018	10.1016/j.ajodo.2017.07.015.	intervention
		Barber S, Bekker HL, Meads D, Pavitt S, Khambay B.	
		Identification and appraisal of outcome measures	
		used to evaluate hypodontia care: A systematic	
		review.Am J Orthod Dentofacial Orthop. 2018	not an
02 41050	2010	Feb;153(2):184-194.e18. doi:	orthodontic
83. AJODO	2018	10.1016/j.ajodo.2017.10.010.	intervention
		Eslami E, Barkhordar H, Abramovitch K, Kim J,	
		Masoud MI. Cone-beam computed tomography vs	
	1	conventional radiography in visualization of	
	1	maxillary impacted-canine localization: A	
	1	systematic review of comparative studies. Am J	not an
94 41000	2017	Orthod Dentofacial Orthop. 2017 Feb;151(2):248-	orthodontic
84. AJODO	2017	258. doi: 10.1016/j.ajodo.2016.07.018.	intervention
	1	Al-Moghrabi D, Salazar FC, Pandis N, Fleming PS.	and the offering of
		Compliance with removable orthodontic appliances	not the effects of
	1	and adjuncts: A systematic review and meta-	orthodontic
05 41050	2017	analysis. Am J Orthod Dentofacial Orthop. 2017	interventions
85. AJODO	2017	Jul;152(1):17-32. doi: 10.1016/j.ajodo.2017.03.019.	were assessed
	1	Al Makhmari SA, Kaklamanos EG, Athanasiou AE.	
		Short-term and long-term effectiveness of powered	
	1	toothbrushes in promoting periodontal health	
	1	during orthodontic treatment: A systematic review	
	1	and meta-analysis. Am J Orthod Dentofacial	not an
		Orthop. 2017 Dec;152(6):753-766.e7. doi:	orthodontic
86. AJODO	2017	10.1016/j.ajodo.2017.09.003.	intervention
	1	Rossini G, Parrini S, Castroflorio T, Deregibus A,	not an
		Debernardi CL. Diagnostic accuracy and	orthodontic
87. AJODO	2016	measurement sensitivity of digital models for	intervention

		and a straight and a straight and the straight s	
		orthodontic purposes: A systematic review. Am J	
		Orthod Dentofacial Orthop. 2016 Feb;149(2):161- 70. doi: 10.1016/j.ajodo.2015.06.029.	
		Peiró-Guijarro MA, Guijarro-Martínez R, Hernández-	
		Alfaro F. Surgery first in orthognathic surgery: A	
		systematic review of the literature. Am J Orthod	not an
		Dentofacial Orthop. 2016 Apr;149(4):448-62. doi:	orthodontic
88. AJODO	2016	10.1016/j.ajodo.2015.09.022.	intervention
88. AJODO	2010	Incerti-Parenti S, Checchi V, Ippolito DR, Gracco A,	intervention
		Alessandri-Bonetti G. Periodontal status after	
		surgical-orthodontic treatment of labially impacted	
		canines with different surgical techniques: A	
		systematic review. Am J Orthod Dentofacial Orthop.	not an
		2016 Apr;149(4):463-72. doi:	orthodontic
89. AJODO	2016	10.1016/j.ajodo.2015.10.019.	intervention
	2010	Ata-Ali F, Ata-Ali J, Ferrer-Molina M, Cobo T, De	
		Carlos F, Cobo J. Adverse effects of lingual and	
		buccal orthodontic techniques: A systematic review	
		and meta-analysis. Am J Orthod Dentofacial	assessed
		Orthop. 2016 Jun;149(6):820-9. doi:	exclusively
90. AJODO	2016	10.1016/j.ajodo.2015.11.031.	adverse effects
		Sandhu SS, Cheema MS, Khehra HS. Comparative	
		effectiveness of pharmacologic and	
		nonpharmacologic interventions for orthodontic	
		pain relief at peak pain intensity: A Bayesian	
		network meta-analysis. Am J Orthod Dentofacial	not an
		Orthop. 2016 Jul;150(1):13-32. doi:	orthodontic
91. AJODO	2016	10.1016/j.ajodo.2015.12.025.	intervention
		Cadenas-Perula M, Yañez-Vico RM, Solano-Reina E,	
		Iglesias-Linares A. Effectiveness of biologic methods	
		of inhibiting orthodontic tooth movement in animal	
		studies. Am J Orthod Dentofacial Orthop. 2016	review of animal
92. AJODO	2016	Jul;150(1):33-48. doi: 10.1016/j.ajodo.2016.01.015.	studies
		Parrini S, Rossini G, Castroflorio T, Fortini A,	
		Deregibus A, Debernardi C. Laypeople's perceptions	not the effects of
		of frontal smile esthetics: A systematic review. Am J	orthodontic
		Orthod Dentofacial Orthop. 2016 Nov;150(5):740-	interventions
93. AJODO	2016	750. doi: 10.1016/j.ajodo.2016.06.022.	were assessed
		Roscoe MG, Meira JB, Cattaneo PM. Association of	
		orthodontic force system and root resorption: A	accased
		systematic review. Am J Orthod Dentofacial Orthop.	assessed
04 41000	2015	2015 May;147(5):610-26. doi:	exclusively
94. AJODO	2015	10.1016/j.ajodo.2014.12.026.	adverse effects
		Afrand M, Ling CP, Khosrotehrani S, Flores-Mir C,	
		Lagravère-Vich MO. Anterior cranial-base time-	not an
		related changes: A systematic review. Am J Orthod Dentofacial Orthop. 2014 Jul;146(1):21-32.e6. doi:	not an orthodontic
95. AJODO	2014	10.1016/j.ajodo.2014.03.019.	intervention
55. AJODO	2014	Tsichlaki A, O'Brien K. Do orthodontic research	
		outcomes reflect patient values? A systematic	
		review of randomized controlled trials involving	not the effects of
		children. Am J Orthod Dentofacial Orthop. 2014	orthodontic
		Sep;146(3):279-85. doi:	interventions
96. AJODO	2014	10.1016/j.ajodo.2014.05.022.	were assessed
JU. AJUDU	2014	10.1010/ j.0j000.2017.05.022.	Were assessed

I			
		Katyal V, Pamula Y, Martin AJ, Daynes CN, Kennedy	
		JD, Sampson WJ. Craniofacial and upper airway	
		morphology in pediatric sleep-disordered	
		breathing: Systematic review and meta-analysis.Am	not an
		J Orthod Dentofacial Orthop. 2013 Jan;143(1):20-	orthodontic
97. AJODO	2013	30.e3. doi: 10.1016/j.ajodo.2012.08.021.	intervention
		Chen H, Liu X, Dai J, Jiang Z, Guo T, Ding Y. Effect of	
		remineralizing agents on white spot lesions after	
		orthodontic treatment: a systematic review. Am J	not an
		Orthod Dentofacial Orthop. 2013 Mar;143(3):376-	orthodontic
98. AJODO	2013	382.e3. doi: 10.1016/j.ajodo.2012.10.013.	intervention
		Fleming PS, Eliades T, Katsaros C, Pandis N. Curing	
		lights for orthodontic bonding: a systematic review	
		and meta-analysis. Am J Orthod Dentofacial	not an
		Orthop. 2013 Apr;143(4 Suppl):S92-103. doi:	orthodontic
99. AJODO	2013	10.1016/j.ajodo.2012.07.018.	intervention
33. AJODO	2013	Kloukos D, Pandis N, Eliades T. Bisphenol-A and	Intervention
		residual monomer leaching from orthodontic	
		adhesive resins and polycarbonate brackets: a	
		systematic review. Am J Orthod Dentofacial Orthop.	not an
		2013 Apr;143(4 Suppl):S104-12.e1-2. doi:	orthodontic
100.AJODO	2013	10.1016/j.ajodo.2012.11.015.	intervention
		Hua F, He H, Ngan P, Bouzid W. Prevalence of peg-	
		shaped maxillary permanent lateral incisors: A	
		meta-analysis. Am J Orthod Dentofacial Orthop.	not an
		2013 Jul;144(1):97-109. doi:	orthodontic
101.AJODO	2013	10.1016/j.ajodo.2013.02.025.	intervention
		Fleming PS, Johal A, Pandis N. Self-etch primers and	
		conventional acid-etch technique for orthodontic	
		bonding: a systematic review and meta-analysis.	not an
		Am J Orthod Dentofacial Orthop. 2012	orthodontic
102.AJODO	2012	Jul;142(1):83-94. doi: 10.1016/j.ajodo.2012.02.023.	intervention
		Hafez HS, Shaarawy SM, Al-Sakiti AA, Mostafa YA.	
		Dental crowding as a caries risk factor: a systematic	
		review. Am J Orthod Dentofacial Orthop. 2012	not an
		Oct;142(4):443-50. doi:	orthodontic
103.AJODO	2012	10.1016/j.ajodo.2012.04.018.	intervention
		Meursinge Reynders RA, Ronchi L, Ladu L, van	
		Etten-Jamaludin F, Bipat S. Insertion torque and	
		success of orthodontic mini-implants: a systematic	not the effects of
		review. Am J Orthod Dentofacial Orthop. 2012	orthodontic
		Nov;142(5):596-614.e5. doi:	interventions
104.AJODO	2012	10.1016/j.ajodo.2012.06.013.	were assessed
104.0000	2012	Papageorgiou SN, Zogakis IP, Papadopoulos MA.	
		Failure rates and associated risk factors of	
			not the effects of
		orthodontic miniscrew implants: a meta-analysis.	
		Am J Orthod Dentofacial Orthop. 2012	orthodontic
105 41000	2012	Nov;142(5):577-595.e7. doi:	interventions
105.AJODO	2012	10.1016/j.ajodo.2012.05.016.	were assessed
		Crismani AG, Bertl MH, Celar AG, Bantleon HP,	
		Burstone CJ. Miniscrews in orthodontic treatment:	not the effects of
		review and analysis of published clinical trials. Am J	orthodontic
		Orthod Dentofacial Orthop. 2010 Jan;137(1):108-	interventions
106.AJODO	2010	13. doi: 10.1016/j.ajodo.2008.01.027.	were assessed

		I	
		Weltman B, Vig KW, Fields HW, Shanker S, Kaizar	
		EE. Root resorption associated with orthodontic	
		tooth movement: a systematic review. Am J Orthod	assessed
		Dentofacial Orthop. 2010 Apr;137(4):462-76;	exclusively
107.AJODO	2010	discussion 12A. doi: 10.1016/j.ajodo.2009.06.021.	adverse effects
		Finnema KJ, Ozcan M, Post WJ, Ren Y, Dijkstra PU.	
		In-vitro orthodontic bond strength testing: a	
		systematic review and meta-analysis. Am J Orthod	review of
		Dentofacial Orthop. 2010 May;137(5):615-622.e3.	laboratory
108.AJODO	2010	doi: 10.1016/j.ajodo.2009.12.021.	studies
		Rogers S, Chadwick B, Treasure E. Fluoride-	
		containing orthodontic adhesives and	
		decalcification in patients with fixed appliances: a	
		systematic review. Am J Orthod Dentofacial Orthop.	not an
		2010 Oct;138(4):390.e1-8; discussion 390-1. doi:	orthodontic
109.AJODO	2010	10.1016/j.ajodo.2010.02.025.	intervention
		Ahrens A, McGrath C, Hägg U. Subjective efficacy of	
		oral appliance design features in the management	
		of obstructive sleep apnea: a systematic review. Am	not an
		J Orthod Dentofacial Orthop. 2010 Nov;138(5):559-	orthodontic
110.AJODO	2010	76. doi: 10.1016/j.ajodo.2010.01.030.	intervention
		Al-Riyami S, Moles DR, Cunningham SJ.	
		Orthognathic treatment and temporomandibular	
		disorders: a systematic review. Part 1. A new	
		quality-assessment technique and analysis of study	not the effects of
		characteristics and classifications. Am J Orthod	orthodontic
		Dentofacial Orthop. 2009 Nov;136(5):624.e1-15;	interventions
111.AJODO	2009	discussion 624-5. doi: 10.1016/j.ajodo.2009.02.021.	were assessed
		Al-Riyami S, Cunningham SJ, Moles DR.	
		Orthognathic treatment and temporomandibular	
		disorders: a systematic review. Part 2. Signs and	not the effects of
		symptoms and meta-analyses. Am J Orthod	orthodontic
		Dentofacial Orthop. 2009 Nov;136(5):626.e1-16,	interventions
112.AJODO	2009	discussion 626-7. doi: 10.1016/j.ajodo.2009.02.022.	were assessed
		Santana LG, Avelar K, Marques LS. Association	
		between arch perimeter management and the	
		occurrence of mandibular second molar eruption	not the effects of
		disturbances. Angle Orthod. 2021 Jul 1;91(4):544-	orthodontic
		554. doi: 10.2319/091720-799.1. PMID: 33587114;	interventions
113.AO	2021	PMCID: PMC8259765.	were assessed
		Li Z, Zhou J, Chen S. The effectiveness of locally	
		injected platelet-rich plasma on orthodontic tooth	
		movement acceleration. Angle Orthod. 2021 May	
		1;91(3):391-398. doi: 10.2319/061320-544.1. PMID:	review of animal
114.AO	2021	33566068; PMCID: PMC8084470.	studies
		Veginadu P, Tavva SR, Muddada V, Gorantla S.	
		Effect of pharmacological agents on relapse	
		following orthodontic tooth movement. Angle	
		Orthod. 2020 Jul 1;90(4):598-606. doi:	
		10.2319/092619-613.1. PMID: 33378496; PMCID:	review of animal
115.AO	2021	PMC8028453.	studies
		Yao K, Zhu G, Chen M, Zhang B, Wu Y, Li P. Effect of	review included
		surgery-first orthognathic approach on oral health-	orthognathic
		related quality of life. Angle Orthod. 2020 Sep	surgical
116.AO	2020	1;90(5):723-733. doi: 10.2319/112619-749.1.	interventions

		Erratum in: Angle Orthod. 2021 Mar 1;91(2):279-	
		281. PMID: 33378482; PMCID: PMC8032263.	
		Sam A, Currie K, Oh H, Flores-Mir C, Lagravére-Vich	
		M. Reliability of different three-dimensional	
		cephalometric landmarks in cone-beam computed	
		tomography: A systematic review. Angle Orthod.	not an
		2019 Mar;89(2):317-332. doi: 10.2319/042018-	orthodontic
117.AO	2019	302.1. Epub 2018 Nov 13. Bastos RTDRM, Blagitz MN, Aragón MLSC, Maia LC,	intervention
		Normando D.Periodontal side effects of rapid and	
		slow maxillary expansion: A systematic review.	assessed
		Angle Orthod. 2019 Jul;89(4):651-660. doi:	exclusively
118.AO	2019	10.2319/060218-419.1. Epub 2019 Feb 11.	adverse effects
1100.00	2015	Ponce-Garcia C, Lagravere-Vich M, Cevidanes LHS,	
		de Olivera Ruellas AC, Carey J, Flores-Mir C.	
		Reliability of three-dimensional anterior cranial	
		base superimposition methods for assessment of	
		overall hard tissue changes: A systematic review.	not an
		Angle Orthod. 2018 Mar;88(2):233-245. doi:	orthodontic
119.AO	2018	10.2319/071217-468.1.	intervention
		Lima IFP, de Andrade Vieira W, de Macedo	
		Bernardino Í, Costa PA, Lima APB, Pithon MM,	
		Paranhos LR. Influence of reminder therapy for	
		controlling bacterial plaque in patients undergoing	
		orthodontic treatment: A systematic review and	not an
		meta-analysis. Angle Orthod. 2018 Jul;88(4):483-	orthodontic
120.AO	2018	493. doi: 10.2319/111117-770.1.	intervention
		Sun W, Xia K, Tang L, Liu C, Zou L, Liu J. Accuracy of	
		panoramic radiography in diagnosing maxillary	
		sinus-root relationship: A systematic review and	not an
		meta-analysis. Angle Orthod. 2018 Nov;88(6):819-	orthodontic
121.AO	2018	829. doi: 10.2319/022018-135.1.	intervention
		Eslami E, Katz ES, Baghdady M, Abramovitch K,	
		Masoud MI. Are three-dimensional airway	
		evaluations obtained through computed and cone- beam computed tomography scans predictable	
		from lateral cephalograms? A systematic review of	not an
		evidence. Angle Orthod. 2017 Jan;87(1):159-167.	orthodontic
122.AO	2017	doi: 10.2319/032516-243.1.	intervention
122.00	2017	Yi J, Sun Y, Li Y, Li C, Li X, Zhao Z. Cone-beam	
		computed tomography versus periapical radiograph	
		for diagnosing external root resorption: A	
		systematic review and meta-analysis. Angle Orthod.	not an
		2017 Mar;87(2):328-337. doi: 10.2319/061916-	orthodontic
123.AO	2017	481.1.	intervention
		Currie K, Sawchuk D, Saltaji H, Oh H, Flores-Mir C,	
		Lagravere M. Posterior cranial base natural growth	
		and development: A systematic review. Angle	not an
		Orthod. 2017 Nov;87(6):897-910. doi:	orthodontic
124.AO	2017	10.2319/032717-218.1.	intervention
		Pachêco-Pereira C, Abreu LG, Dick BD, De Luca	
		Canto G, Paiva SM, Flores-Mir C. Patient satisfaction	
		after orthodontic treatment combined with	review included
		orthognathic surgery: A systematic review. Angle	orthognathic
105.10		Orthod. 2016 May;86(3):495-508. doi:	surgical
125.AO	2016	10.2319/040615-227.1.	interventions

		Gong A, Li J, Wang Z, Li Y, Hu F, Li Q, Miao D, Wang	
		L. Cranial base characteristics in anteroposterior	not an
		malocclusions: A meta-analysis.Angle Orthod. 2016	orthodontic
126.AO	2016	Jul;86(4):668-80. doi: 10.2319/032315-186.1.	intervention
120.80	2010	Maniewicz Wins S, Antonarakis GS, Kiliaridis S.	not the effects of
		Predictive factors of sagittal stability after	orthodontic
		treatment of Class II malocclusions. Angle Orthod.	interventions
127.AO	2016	2016 Nov;86(6):1033-1041.	were assessed
127.80	2010	Aminoshariae A, Aminoshariae A, Valiathan M,	were assessed
		Kulid JC. Association of genetic polymorphism and	assessed
		external apical root resorption. Angle Orthod. 2016	exclusively
128.AO	2016	Nov;86(6):1042-1049.	adverse effects
120.80	2010	Rossini G, Parrini S, Castroflorio T, Fortini A,	adverse effects
		Deregibus A, Debernardi CL. Children's perceptions	
		of smile esthetics and their influence on social	not an
		judgment. Angle Orthod. 2016 Nov;86(6):1050-	orthodontic
129.AO	2016	1055.	intervention
129.AU	2010		intervention
		Aljabaa A, McDonald F, Newton JT. A systematic review of randomized controlled trials of	
		interventions to improve adherence among orthodontic patients aged 12 to 18. Angle Orthod.	not an
		2015 Mar;85(2):305-13. doi: 10.2319/031214-	not an orthodontic
130.AO	2015	184.1.	intervention
150.AU	2015	Pachêco-Pereira C, De Luca Canto G, Major PW,	intervention
		Flores-Mir C. Variation of orthodontic treatment	not the effects of
		decision-making based on dental model type: A	orthodontic
		systematic review. Angle Orthod. 2015	interventions
121 40	2015	, .	
131.AO	2015	May;85(3):501-9. doi: 10.2319/051214-343.1.	were assessed
		Al-Jewair TS. Meta-analysis on the mandibular	
		dimensions effects of the MARA appliance in	
122.00	2045	patients with Class II malocclusions. Angle Orthod.	only one
132.AO	2015	2015 Jul;85(4):706-14. doi: 10.2319/052814-378.1.	reviewer
		Grewal Bach GK, Torrealba Y, Lagravère MO.	not the effects of
		Orthodontic bonding to porcelain: a systematic	orthodontic
122.40	2014	review. Angle Orthod. 2014 May;84(3):555-60. doi:	interventions
133.AO	2014	10.2319/083013-636.1.	were assessed
		Marquezan M, Mattos CT, Sant'Anna EF, de Souza	and the offerster f
		MM, Maia LC. Does cortical thickness influence the	not the effects of
		primary stability of miniscrews?: A systematic	orthodontic
124 40	2014	review and meta-analysis. Angle Orthod. 2014	interventions
134.AO	2014	Nov;84(6):1093-103. doi: 10.2319/093013-716.1.	were assessed
		Lione R, Franchi L, Cozza P. Does rapid maxillary	
		expansion induce adverse effects in growing	assessed
125 4.0	2012	subjects? Angle Orthod. 2013 Jan;83(1):172-82. doi:	exclusively
135.AO	2013	10.2319/041012-300.1.	adverse effects
		Al-Anezi SA, Harradine NW. Quantifying plaque	assessed
126.40	2012	during orthodontic treatment:. Angle Orthod. 2012	exclusively
136.AO	2012	Jul;82(4):748-53. doi: 10.2319/050111-312.1.	adverse effects
		Luu NS, Nikolcheva LG, Retrouvey JM, Flores-Mir C,	
		El-Bialy T, Carey JP, Major PW. Linear	
		measurements using virtual study models. Angle	not an
		Orthod. 2012 Nov;82(6):1098-106. doi:	orthodontic
137.AO	2012	10.2319/110311-681.1.	intervention
		Saltaji H, Major MP, Altalibi M, Youssef M, Flores-	assessed
		Mir C. Long-term skeletal stability after maxillary	exclusively
138.AO	2012	advancement with distraction osteogenesis in cleft	patients with

		lip and palate patients. Angle Orthod. 2012	congenital
		Nov;82(6):1115-22. doi: 10.2319/011212-27.1.	anomalies
		Santiago RC, de Miranda Costa LF, Vitral RW, Fraga	
		MR, Bolognese AM, Maia LC. Cervical vertebral	
		maturation as a biologic indicator of skeletal	not an
		maturity. Angle Orthod. 2012 Nov;82(6):1123-31.	orthodontic
139.AO	2012	doi: 10.2319/103111-673.1.	intervention
1000.00	2012	Archambault A, Lacoursiere R, Badawi H, Major PW,	
		Carey J, Flores-Mir C. Torque expression in stainless	
		steel orthodontic brackets. A systematic review.	review of
		Angle Orthod. 2010 Jan;80(1):201-10. doi:	laboratory
140.AO	2010	10.2319/080508-352.1.	studies
110.710	2010	Xiaoting L, Yin T, Yangxi C. Interventions for pain	not the effects of
		during fixed orthodontic appliance therapy. A	orthodontic
		systematic review. Angle Orthod. 2010	interventions
141.AO	2010	Sep;80(5):925-32. doi: 10.2319/010410-10.1.	were assessed
141.AU	2010		
		Magalhães IB, Pereira LJ, Marques LS, Gameiro GH.	not the effects of orthodontic
		The influence of malocclusion on masticatory	
142.40	2010	performance. A systematic review. Angle Orthod.	interventions
142.AO	2010	2010 Sep;80(5):981-7. doi: 10.2319/011910-33.1.	were assessed
		Giudice AL, Rustico L, Longo M, Oteri G,	
		Papadopoulos MA, Nucera R. Complications	
	1	reported with the use of orthodontic miniscrews: A	
		systematic review. Korean J Orthod. 2021 May	assessed
		25;51(3):199-216. doi: 10.4041/kjod.2021.51.3.199.	exclusively
143.KJO	2021	PMID: 33984227; PMCID: PMC8133901.	adverse effects
	1	Sivarajan S, Mani SA, John J, Fayed MMS, Kook YA,	
		Wey MC. The global distribution of permanent	not the effects of
		canine hypodontia: A systematic review. Korean J	orthodontic
		Orthod. 2021 Jan 25;51(1):55-74. doi:	interventions
		10.4041/kjod.2021.51.1.55. PMID: 33446621;	were assessed
144.KJO	2021	PMCID: PMC7837799.	
		Savoldi F, Papoutsi A, Dianiskova S, Dalessandri D,	
		Bonetti S, Tsoi JKH, Matinlinna JP, Paganelli C.	
		Resistance to sliding in orthodontics: misconception	
		or method error? A systematic review and a	review of
		proposal of a test protocol.Korean J Orthod. 2018	laboratory
145.KJO	2018	Jul;48(4):268-280. doi: 10.4041/kjod.2018.48.4.268.	studies
		Nowrin SA, Jaafar S, Ab Rahman N, Basri R, Alam	
		MK, Shahid F. Association between genetic	
		polymorphisms and external apical root resorption:	
		A systematic review and meta-analysis.Korean J	assessed
		Orthod. 2018 Nov;48(6):395-404. doi:	exclusively
146.KJO	2018	10.4041/kjod.2018.48.6.395.	adverse effects
		Hong SB, Kusnoto B, Kim EJ, BeGole EA, Hwang HS,	
		Lim HJ. Prognostic factors associated with the	
		success rates of posterior orthodontic miniscrew	not the effects of
		implants: A subgroup meta-analysis. Korean J	orthodontic
		Orthod. 2016 Mar;46(2):111-26. doi:	interventions
147 10	2016		
147.KJO	2016	10.4041/kjod.2016.46.2.111.	were assessed
		Sawchuk D, Currie K1 Vich ML, Palomo JM, Flores-	
		Mir C. Diagnostic methods for assessing maxillary	not the effects of
		skeletal and dental transverse deficiencies: A	orthodontic
		systematic review. Korean J Orthod. 2016	interventions
148.KJO	2016	Sep;46(5):331-42. doi: 10.4041/kjod.2016.46.5.331.	were assessed

			[
		Alessandri-Bonetti G, Ippolito DR, Bartolucci ML,	
		D'Antò V, Incerti-Parenti S. Cephalometric	
		predictors of treatment outcome with mandibular	and the offerster of
		advancement devices in adult patients with	not the effects of
		obstructive sleep apnea: a systematic review.	orthodontic
1 40 1/10	2015	Korean J Orthod. 2015 Nov;45(6):308-21. doi:	interventions
149.KJO	2015	10.4041/kjod.2015.45.6.308.	were assessed
		Berry S, Javed F, Rossouw PE, Barmak AB, Kalogirou	
		EM, Michelogiannakis D. Influence of thyroxine	
		supplementation on orthodontically induced tooth	
		movement and/or inflammatory root resorption: A	
		systematic review. Orthod Craniofac Res. 2021	
		May;24(2):206-213. doi: 10.1111/ocr.12428. Epub	review of animal
150.0&C	2021	2020 Oct 18. PMID: 32991769.	studies
		Roomaney IA, Chetty M. Sella turcica morphology in	
		patients with genetic syndromes: A systematic	
		review. Orthod Craniofac Res. 2021 May;24(2):194-	not the effects of
		205. doi: 10.1111/ocr.12426. Epub 2020 Sep 28.	orthodontcis
151.0&C	2021	PMID: 32920986.	were assessed
		Ahn HW, Kim SJ, Baek SH. Miniplate-anchored	
		maxillary protraction in adolescent patients with	assessed
		cleft lip and palate: A literature review of study	exclusively
		design, type and protocol, and treatment	patients with
		outcomes. Orthod Craniofac Res. 2021 Mar;24	congenital
		Suppl 1:21-30. doi: 10.1111/ocr.12446. Epub 2020	anomalies
152.0&C	2021	Dec 7. PMID: 33253469.	
		Kaklamanos EG, Makrygiannakis MA, Athanasiou	
		AE. Could medications and biologic factors affect	
		post-orthodontic tooth movement changes? A	
		systematic review of animal studies. Orthod	
		Craniofac Res. 2021 Feb;24(1):39-51. doi:	
		10.1111/ocr.12411. Epub 2020 Aug 5. PMID:	review of animal
153.0&C	2021	32654394.	studies
		Marques FBC, de Lima LS, Oliveira PLE, Magno MB,	
		Ferreira DMTP, de Castro ACR, Maciel JVB, Ruellas	
		ACO, Maia LC. Are temporomandibular disorders	
		associated with facial asymmetry? A systematic	
		review and meta-analysis. Orthod Craniofac Res.	not the effects of
		2021 Feb;24(1):1-16. doi: 10.1111/ocr.12404. Epub	orthodontcis
154.0&C	2021	2020 Jul 19. PMID: 32608091.	were assessed
		Pinheiro FHSL, Drummond RJ, Frota CM, Bartzela	
		TN, Dos Santos PB. Comparison of early and	assessed
		conventional autogenous secondary alveolar bone	exclusively
		graft in children with cleft lip and palate: A	patients with
		systematic review. Orthod Craniofac Res. 2020	congenital
		Nov;23(4):385-397. doi: 10.1111/ocr.12394. Epub	anomalies
155.O&C	2020	2020 Jun 28. PMID: 32446283.	
200.000	2020	Xiao WL, Jia KN, Yu G, Zhao N. Association between	
		forkhead box E1 polymorphisms and risk of non-	
		syndromic cleft lip with or without cleft palate: A	
		meta-analysis. Orthod Craniofac Res. 2020	not the effects of
		May;23(2):151-159. doi: 10.1111/ocr.12366. Epub	orthodontics
		1 181987521511731-132. UOI. 10.11111/0C1.12300. EDUD	orthouoritics
156 08 0	2020		wore accessed
156.O&C	2020	2020 Feb 5. PMID: 31944555.	were assessed
156.O&C	2020	2020 Feb 5. PMID: 31944555. Kaklamanos EG, Makrygiannakis MA, Athanasiou	
156.0&C	2020	2020 Feb 5. PMID: 31944555.	were assessed review of animal studies

	r		
		movement rate? A systematic review based on	
		animal studies. Orthod Craniofac Res. 2020	
		May;23(2):143-150. doi: 10.1111/ocr.12357. Epub	
		2019 Nov 9. PMID: 31705727.	
		Elsten EECM, Caron CJJM, Dunaway DJ, Padwa BL,	
		Forrest C, Koudstaal MJ. Dental anomalies in	
		craniofacial microsomia: A systematic review.	
		Orthod Craniofac Res. 2020 Feb;23(1):16-26. doi:	not the effects of
		10.1111/ocr.12351. Epub 2019 Oct 28. PMID:	orthodontics
158.0&C	2020	31608577; PMCID: PMC7003932.	were assessed
		Wu Z, Zhang X, Li Z, Liu Y, Jin H, Chen Q, Guo J. A	
		Bayesian network meta-analysis of orthopaedic	
		treatment in Class III malocclusion: Maxillary	
		protraction with skeletal anchorage or a rapid	
		maxillary expander. Orthod Craniofac Res. 2020	
		Feb;23(1):1-15. doi: 10.1111/ocr.12339. Epub 2019	Bayesian network
159.O&C	2020	Sep 15. PMID: 31452316.	analysis was used
100.000		Fang X, Qi R, Liu C. Root resorption in orthodontic	Linging that about
		treatment with clear aligners: A systematic review	
		and meta-analysis. Orthod Craniofac Res. 2019	assessed
		Nov;22(4):259-269. doi: 10.1111/ocr.12337. Epub	exclusively
160.O&C	2019		adverse effects
100.000	2019	2019 Aug 29. PMID: 31323701. Iliadi A, Koletsi D, Eliades T. Forces and moments	
		generated by aligner-type appliances for	
		orthodontic tooth movement: A systematic review	
		and meta-analysis. Orthod Craniofac Res. 2019	review of
		Nov;22(4):248-258. doi: 10.1111/ocr.12333. Epub	laboratory
161.0&C	2019	2019 Jul 9. PMID: 31237410.	studies
		Tarallo F, Chimenti C, Paiella G, Cordaro M,	
		Tepedino M. Biomarkers in the gingival crevicular	
		fluid used to detect root resorption in patients	
		undergoing orthodontic treatment: A systematic	not the effects of
		review. Orthod Craniofac Res. 2019 Nov;22(4):236-	orthodontic
		247. doi: 10.1111/ocr.12329. Epub 2019 Jul 2.	interventions
162.0&C	2019	PMID: 31207100.	were assessed
		Tasios T, Papageorgiou SN, Papadopoulos MA,	
		Tsapas A, Haidich AB. Prevention of orthodontic	
		enamel demineralization: A systematic review with	not the effects of
		meta-analyses. Orthod Craniofac Res. 2019	orthodontic
		Nov;22(4):225-235. doi: 10.1111/ocr.12322. Epub	interventions
163.0&C	2019	2019 May 27. PMID: 31081584.	were assessed
100.040	2015	Javed F, Akram Z, Barillas AP, Kellesarian SV, Ahmed	
		HB, Khan J, Almas K. Outcome of orthodontic	assessed
		· · · ·	
		palatal plate therapy for orofacial dysfunction in	exclusively
		children with Down syndrome: A systematic	patients with
101.005		review.Orthod Craniofac Res. 2018 Feb;21(1):20-26.	congenital
164.0&C	2018	doi: 10.1111/ocr.12211.	anomalies
		Papageorgiou SN, Xavier GM, Cobourne MT, Eliades	
		T. Effect of orthodontic treatment on the	
		subgingival microbiota: A systematic review and	assessed
		meta-analysis. Orthod Craniofac Res. 2018	exclusively
165.0&C	2018	Nov;21(4):175-185. doi: 10.1111/ocr.12237.	adverse effects
		Scariot R, Corso PFCL, Sebastiani AM, Vieira AR. The	not an
		many faces of genetic contributions to	orthodontic
166.O&C	2018	temporomandibular joint disorder: An updated	intervention
· · · · · · · ·			

		review. Orthod Craniofac Res. 2018 Nov;21(4):186-	
		201. doi: 10.1111/ocr.12239. Epub 2018 Sep 11.	
		Antonarakis GS, Palaska PK2 Suri S. Permanent	assessed
		tooth agenesis in individuals with non-syndromic	exclusively
		Robin sequence: a systematic review and meta-	patients with
		analysis. Orthod Craniofac Res. 2017 Nov;20(4):216-	congenital
167.0&C	2017	226. doi: 10.1111/ocr.12204.	anomalies
		Altmann AS, Collares FM, Leitune VC, Samuel SM.	
		The effect of antimicrobial agents on bond strength	
		of orthodontic adhesives: a meta-analysis of in vitro	review of
		studies. Orthod Craniofac Res. 2016 Feb;19(1):1-9.	laboratory
168.O&C	2016	doi: 10.1111/ocr.12100. Epub 2015 Aug 10.	studies
		Tee BC, Sun Z. Mandibular distraction osteogenesis	
		assisted by cell-based tissue engineering: a	review included
		systematic review. Orthod Craniofac Res. 2015	orthognathic
		Apr;18 Suppl 1:39-49. doi: 10.1111/ocr.12087. (in	surgical
169.0&C	2015	suppl issue)	interventions
		De Luca Canto G, Pachêco-Pereira C, Lagravere MO,	
		Flores-Mir C, Major PW. Intra-arch dimensional	
		measurement validity of laser-scanned digital	
		dental models compared with the original plaster	not an
		models: a systematic review. Orthod Craniofac Res.	orthodontic
170.0&C	2015	2015 May;18(2):65-76. doi: 10.1111/ocr.12068.	intervention
		Austin SL, Mattick CR, Waterhouse PJ. Distraction	
		osteogenesis versus orthognathic surgery for the	
		treatment of maxillary hypoplasia in cleft lip and	review included
		palate patients: a systematic review.	orthognathic
		Orthod Craniofac Res. 2015 May;18(2):96-108. doi:	surgical
171.0&C	2015	10.1111/ocr.12063.	interventions
		Koretsi V, Chatzigianni A, Sidiropoulou S. Enamel	
		roughness and incidence of caries after	
		interproximal enamel reduction: a systematic	not an
		review. Orthod Craniofac Res. 2014 Feb;17(1):1-13.	orthodontic
172.0&C	2014	doi: 10.1111/ocr.12030.	intervention
		Pittayapat P, Limchaichana-Bolstad N, Willems G,	
		Jacobs R. Three-dimensional cephalometric analysis	
		in orthodontics: a systematic review. Orthod	not an
		Craniofac Res. 2014 May;17(2):69-91. doi:	orthodontic
173.0&C	2014	10.1111/ocr.12034.	intervention
		Perinetti G, Primožič J, Castaldo A, Di Lenarda R,	
		Contardo L. Is gingival crevicular fluid volume	
		sensitive to orthodontic tooth movement? A	
		systematic review of split-mouth longitudinal	assessed
		studies. Orthod Craniofac Res. 2013 Feb;16(1):1-19.	exclusively
174.0&C	2013	doi: 10.1111/ocr.12005.	adverse effects
		Andrade DC, Loureiro CA, Araújo VE, Riera R,	
		Atallah AN. Treatment for agenesis of maxillary	
		lateral incisors: a systematic review. Orthod	
		Craniofac Res. 2013 Aug;16(3):129-36. doi:	
			1
175.0&C	2013	10.1111/ocr.12015.	empty review
175.0&C	2013		empty review
175.0&C	2013	Angelopoulou MV, Vlachou V, Halazonetis DJ.	empty review
175.0&C	2013	Angelopoulou MV, Vlachou V, Halazonetis DJ. Pharmacological management of pain during	
175.0&C	2013	Angelopoulou MV, Vlachou V, Halazonetis DJ.	empty review assessed exclusively

	Gritsch K, Laroche N, Morgon L, Al-Hity R, Vico L,	
	Colon P, Grosgogeat B. A systematic review of	
	methods for tissue analysis in animal studies on	
	orthodontic mini-implants. Orthod Craniofac Res.	
	2012 Aug;15(3):135-47. doi: 10.1111/j.1601-	review of animal
177.O&C 2012	6343.2012.01548.x.	studies
	Papadopoulos MA, Koumpridou EN, Vakalis ML,	
	Papageorgiou SN. Effectiveness of pre-surgical	assessed
	infant orthopedic treatment for cleft lip and palate	exclusively
	patients: a systematic review and meta-analysis.	patients with
	Orthod Craniofac Res. 2012 Nov;15(4):207-36. doi:	congenital
178.0&C 2012	10.1111/j.1601-6343.2012.01552.x.	anomalies
	Fleming PS, Marinho V, Johal A. Orthodontic	
	measurements on digital study models compared	
	with plaster models: a systematic review. Orthod	not an
	Craniofac Res. 2011 Feb;14(1):1-16. doi:	orthodontic
179.0&C 2011	10.1111/j.1601-6343.2010.01503.x.	intervention
	Joss-Vassalli I, Grebenstein C, Topouzelis N, Sculean	
	A, Katsaros C. Orthodontic therapy and gingival	
	recession: a systematic review. Orthod Craniofac	assessed
	Res. 2010 Aug;13(3):127-41. doi: 10.1111/j.1601-	exclusively
180.O&C 2010	6343.2010.01491.x.	adverse effects

*Description of the abbreviated journals:

Cochrane library: Cochrane Database of Systematic Reviews

AJODO: American Journal of Orthodontics and Dentofacial Orthopedics

EJO: European Journal of Orthodontics

AO: Angle Orthodontist

KJO: Korean Journal of Orthodontics

O&CR: Orthodontics and Craniofacial Research

				Rationale for
Journal	*	Year	Reference	exclusion
			Parkin N, Benson PE, Thind B, Shah A, Khalil I,	
			Ghafoor S. Open versus closed surgical exposure of	
			canine teeth that are displaced in the roof of the	not the effects of
	C. J.		mouth.Cochrane Database Syst Rev. 2017 Aug	orthodontic
1.	Cochrane	2017	21;8:CD006966. doi:	interventions
	library	2017	10.1002/14651858.CD006966.pub3.	were assessed
			Littlewood SJ, Millett DT, Doubleday B, Bearn DR,	
			Worthington HV. Retention procedures for stabilising tooth position after treatment with	
			orthodontic braces. Cochrane Database Syst Rev.	assessed
2.	Cochrane		2016 Jan 29;(1):CD002283. doi:	exclusively
۷.	library	2016	10.1002/14651858.CD002283.pub4.	adverse effects
	library	2010	10.1002/14051858.60002285.pdb4.	the review is
				about a specific
				outcome of an
		1		intervention,
			Rekhi U, Catunda RQ, Gibson MP. Surgically	which is
			accelerated orthodontic techniques and periodontal	ambiguous and
			response: a systematic review. Eur J Orthod. 2020	could also be an
			Jan 15:cjz103. doi: 10.1093/ejo/cjz103. Epub ahead	adverse effect
3.	EJO	2020	of print. PMID: 31942984.	
			Bortolotti F, Solidoro L, Bartolucci ML, Incerti	
			Parenti S, Paganelli C, Alessandri-Bonetti G. Skeletal	
			and dental effects of surgically assisted rapid palatal	
			expansion: a systematic review of randomized	review included
			controlled trials. Eur J Orthod. 2020 Sep	orthognathic
			11;42(4):434-440. doi: 10.1093/ejo/cjz057. PMID:	surgical
4.	EJO	2020	31365925.	interventions
			Papageorgiou SN, Koletsi D, Iliadi A, Peltomaki T,	
			Eliades T. Treatment outcome with orthodontic	
			aligners and fixed appliances: a systematic review	
			with meta-analyses. Eur J Orthod. 2020 Jun	review included
			23;42(3):331-343. doi: 10.1093/ejo/cjz094. PMID:	orthognathic
			31758191.	surgical
5.	EJO	2020		interventions
				the review is
				about a specific
		1	Bellini-Pereira SA, Pupulim DC, Aliaga-Del Castillo A,	outcome of an
			Henriques JFC, Janson G. Time of maxillary molar	intervention,
			distalization with non-compliance intraoral	which is
			distalizing appliances: a meta-analysis. Eur J Orthod.	ambiguous and
~	510	2010	2019 Nov 15;41(6):652-660. doi:	could also be an
6.	EJO	2019	10.1093/ejo/cjz030. PMID: 31107942.	adverse effect
			Swidi AJ, Griffin AE, Buschang PH. Mandibular	
			alignment changes after full-fixed orthodontic	accord
			treatment: a systematic review and meta-analysis.	assessed
7	EIO	2019	Eur J Orthod. 2019 Nov 15;41(6):609-621. Doi:	exclusively adverse effects
7.	EJO	2019	10.1093/ejo/cjz004. PMID: 30788505.	
			Phuong A, Fagundes NCF, Abtahi S, Roberts MR,	assessed
0	EIO	2019	Major PW, Flores-Mir C. Additional appointments	exclusively
8.	EJO	2013	and discomfort associated with compliance-free	adverse effects

Excluded studies during the full text screening (n=45) with the rationale for exclusion

			I
		fixed Class II corrector treatment: a systematic	
		review. Eur J Orthod. 2019 Aug 8;41(4):404-414.	
		Doi: 10.1093/ejo/cjy074.	
		Almuzian M, Rizk MZ, Ulhaq A, Alharbi F, Alomari S,	
		Mohammed H. Effectiveness of different debonding	
		techniques and adjunctive methods on pain and	
		discomfort perception during debonding fixed	not an
		orthodontic appliances: a systematic review. Eur J	orthodontic
		Orthod. 2019 Sep 21;41(5):486-494. doi:	intervention
9. EJO	2019	10.1093/ejo/cjz013. PMID: 30934051.	review
		Cassina C, Papageorgiou SN, Eliades T. Open versus	
		closed surgical exposure for permanent impacted	not the effects of
		canines: a systematic review and meta-analyses.Eur	orthodontic
		J Orthod. 2018 Jan 23;40(1):1-10. doi:	interventions
10. EJO	2018	10.1093/ejo/cjx047.	were assessed
		Sampaziotis D, Tsolakis IA, Bitsanis E, Tsolakis AI.	
		Open versus closed surgical exposure of palatally	
		impacted maxillary canines: comparison of the	not the effects of
		different treatment outcomes-a systematic	orthodontic
		review.Eur J Orthod. 2018 Jan 23;40(1):11-22. doi:	interventions
11. EJO	2018	10.1093/ejo/cjw077.	were assessed
		Papageorgiou SN, Papadelli AA, Eliades T. Effect of	
		orthodontic treatment on periodontal clinical	
		attachment: a systematic review and meta-	assessed
		analysis.Eur J Orthod. 2018 Apr 6;40(2):176-194.	exclusively
12. EJO	2018	doi: 10.1093/ejo/cjx052.	adverse effects
		Buzatta LN, Shimizu RH, Shimizu IA, Pachêco-Pereira	
		C, Flores-Mir C, Taba M Jr, Porporatti AL, De Luca	
		Canto G. Gingival condition associated with two	
		types of orthodontic fixed retainers: a meta-	assessed
		analysis. Eur J Orthod. 2017 Aug 1;39(4):446-452.	exclusively
13. EJO	2017	doi: 10.1093/ejo/cjw057.	adverse effects
		Buck LM, Dalci O, Darendeliler MA, Papageorgiou	
		SN, Papadopoulou AK. Volumetric upper airway	review included
		changes after rapid maxillary expansion: a	orthognathic
		systematic review and meta-analysis. Eur J Orthod.	surgical
14. EJO	2017	2017 Oct 1;39(5):463-473. doi: 10.1093/ejo/cjw048.	interventions
17. 600	2011	Bock NC, von Bremen J, Ruf S. Stability of Class II	
		fixed functional appliance therapya systematic	assessed
		review and meta-analysis. Eur J Orthod. 2016	exclusively
15. EJO	2016	Apr;38(2):129-39. doi: 10.1093/ejo/cjv009.	adverse effects
13. LIO	2010		review did not
			assess the effect
			of a specific type
			of
			intervention(s),
			but assessed an
			undefined
		Collonius O. Dotrón C. Diärnessen I. Newlund A.	orthodontic
		Sollenius O, Petrén S, Björnsson L, Norlund A,	intervention, e.g.,
		Bondemark L. Health economic evaluations in	orthodontic
16 510	2010	orthodontics: a systematic review. Eur J Orthod.	treatment as a
16. EJO	2016	2016 Jun;38(3):259-65. doi: 10.1093/ejo/cjv040.	whole
17 510	2010	Janson G, Mendes LM, Junqueira CH, Garib DG.	review included
17. EJO	2016	Soft-tissue changes in Class II malocclusion patients	orthognathic

		tracted with outractional a sustamatic ravious Fur L	ourgical
		treated with extractions: a systematic review. Eur J Orthod. 2016 Dec;38(6):631-637.	surgical interventions
		Rossini G, Parrini S, Castroflorio T, Deregibus A,	Interventions
		Debernardi CL. Periodontal health during clear	
		aligners treatment: a systematic review. Eur J	accord
			assessed
10 510	2015	Orthod. 2015 Oct;37(5):539-43. doi:	exclusively
18. EJO	2015	10.1093/ejo/cju083.	adverse effects
		Yepes E, Quintero P, Rueda ZV, Pedroza A. Optimal	
		force for maxillary protraction facemask therapy in	not the effects of
		the early treatment of class III malocclusion. Eur J	orthodontic
		Orthod. 2014 Oct;36(5):586-94. doi:	interventions
19. EJO	2014	10.1093/ejo/cjt091.	were assessed
		Pacha MM, Fleming PS, Johal A. Complications,	
		impacts, and success rates of different approaches	
		to treatment of Class II malocclusion in adolescents:	
		A systematic review and meta-analysis. Am J Orthod	
		Dentofacial Orthop. 2020 Oct;158(4):477-494.e7.	assessed
		doi: 10.1016/j.ajodo.2020.03.021. Epub 2020 Sep 2.	exclusively
20. AJODO	2020	PMID: 32888735.	adverse effects
		Vandersluis YR, Suri S. Infective endocarditis and	
		orthodontic implications in children: A review of the	not the effects of
		literature. Am J Orthod Dentofacial Orthop. 2020	orthodontic
		Jan;157(1):19-28. doi: 10.1016/j.ajodo.2019.03.027.	interventions
21. AJODO	2020	PMID: 31901273.	were assessed
			review did not
			assess the effect
			of a specific type
			of
			intervention(s),
			but assessed an
		Javidi H, Vettore M, Benson PE. Does orthodontic	undefined
		treatment before the age of 18 years improve oral	orthodontic
		health-related quality of life? A systematic review	intervention, e.g.,
		and meta-analysis. Am J Orthod Dentofacial Orthop.	orthodontic
		2017 Apr;151(4):644-655. doi:	treatment as a
22. AJODO	2017	10.1016/j.ajodo.2016.12.011.	whole
22. AJODO	2017	10.1010/j.aj000.2010.12.011.	
			review did not
			assess the effect
			of a specific type
			of
			intervention(s),
			but assessed an
			undefined
		Tsichlaki A, Chin SY, Pandis N, Fleming PS. How long	orthodontic
		does treatment with fixed orthodontic appliances	intervention, e.g.,
		last? A systematic review. Am J Orthod Dentofacial	orthodontic
		Orthop. 2016 Mar;149(3):308-18. doi:	treatment as a
23. AJODO	2016	10.1016/j.ajodo.2015.09.020.	whole
			review did not
			assess the effect
			r
		Pachêco-Pereira C, Pereira JR, Dick BD, Perez A,	of a specific type
		Pachêco-Pereira C, Pereira JR, Dick BD, Perez A, Flores-Mir C. Factors associated with patient and	of a specific type of
		Flores-Mir C. Factors associated with patient and	of
		Flores-Mir C. Factors associated with patient and parent satisfaction after orthodontic treatment: a	of intervention(s),

	1		intervention, e.g.,
			orthodontic
			treatment as a
			whole
		Mai W, He J, Meng H, Jiang Y, Huang C, Li M, Yuan	
		K, Kang N. Comparison of vacuum-formed and	
		Hawley retainers: a systematic review. Am J Orthod	assessed
		Dentofacial Orthop. 2014 Jun;145(6):720-7. doi:	exclusively
25. AJODO	2014	10.1016/j.ajodo.2014.01.019.	adverse effects
		Greenlee GM, Huang GJ, Chen SS, Chen J, Koepsell T, Hujoel P. Stability of treatment for anterior open-	
		bite malocclusion: a meta-analysis. Am J Orthod	review included orthognathic
		Dentofacial Orthop. 2011 Feb;139(2):154-69. doi:	surgical
26. AJODO	2011	10.1016/j.ajodo.2010.10.019.	interventions
20. / 0000	2011	Viglianisi A. Effects of lingual arch used as space	interventions
		maintainer on mandibular arch dimension: a	
		systematic review. Am J Orthod Dentofacial Orthop.	review was
		2010 Oct;138(4):382.e1-4; discussion 382-3. doi:	conducted by one
27. AJODO	2010	10.1016/j.ajodo.2010.02.026.	operator only
		Santana LG, Marques LS. Do adjunctive	
		interventions in patients undergoing rapid maxillary	
		expansion increase the treatment effectiveness?	review included
		Angle Orthod. 2021 Jan 1;91(1):119-128. doi: 10.2319/051320-431.1. PMID: 33289794; PMCID:	orthognathic
28. AO	2021	PMC8032281.	surgical interventions
20. AU	2021	Mecenas P, Espinosa DG, Cardoso PC, Normando D.	Interventions
		Stainless steel or titanium mini-implants? Angle	not the effects of
		Orthod. 2020 Jul 1;90(4):587-597. doi:	orthodontic
		10.2319/081619-536.1. PMID: 33378494; PMCID:	interventions
29. AO	2020	PMC8028470.	were assessed
		Moda LB, da Silva Barros ALC, Fagundes NCF,	
		Normando D, Maia LC, Mendes SMDA. Lower fixed	
		retainers: bonded on all teeth or only on canines? A	
		systematic review. Angle Orthod. 2020	
		Jan;90(1):125-143. doi: 10.2319/013019-63.1. Epub 2019 Sep 19. PMID: 31536378; PMCID:	assessed exclusively
30. AO	2020	PMC8087051.	adverse effects
50. AU	2020	Alakttash AM, Fawzi M, Bearn D. Adhesive	
		precoated bracket systems and operator coated	
		bracket systems: Is there any difference? A	
		systematic review and meta-analysis. Angle Orthod.	assessed
		2019 May;89(3):495-504. doi: 10.2319/051818-	exclusively
31. AO	2019	373.1. Epub 2018 Dec 17.	adverse effects
		Diar-Bakirly S, Feres MF, Saltaji H, Flores-Mir C, El-	
		Bialy T. Effectiveness of the transpalatal arch in	
		controlling orthodontic anchorage in maxillary	
		premolar extraction cases: A systematic review and	assessed exclusively
22.40		meta-analysis.Angle Orthod. 2017 Jan;87(1):147-	
	2017	158 doi: 10.2319/021216-120.1	anverse ettects
32. AO	2017	158. doi: 10.2319/021216-120.1. Almasoud NN. Extraction of primary canines for	adverse effects
32. AU	2017	Almasoud NN. Extraction of primary canines for	
32. AU	2017		not the effects of orthodontic
32. AU	2017	Almasoud NN. Extraction of primary canines for interceptive orthodontic treatment of palatally	not the effects of
32. AO 33. AO	2017	Almasoud NN. Extraction of primary canines for interceptive orthodontic treatment of palatally displaced permanent canines: A systematic review.	not the effects of orthodontic
		Almasoud NN. Extraction of primary canines for interceptive orthodontic treatment of palatally displaced permanent canines: A systematic review. Angle Orthod. 2017 Nov;87(6):878-885. doi:	not the effects of orthodontic interventions

		transfer and the Oral Uselth Import Deefile	august and
		treatment, and the Oral Health Impact Profile (OHIP-14): Systematic review and meta-analysis. Angle Orthod. 2015 May;85(3):493-500. doi: 10.2319/051414-348.1.	surgical interventions
35. AO	2013	Long H, Zhou Y, Pyakurel U, Liao L, Jian F, Xue J, Ye N, Yang X, Wang Y, Lai W. Comparison of adverse effects between lingual and labial orthodontic treatment. Angle Orthod. 2013 Nov;83(6):1066-73. doi: 10.2319/010113-2.1.	assessed exclusively adverse effects
		Janson G, Branco NC, Fernandes TM, Sathler R, Garib D, Lauris JR. Influence of orthodontic	review did not assess the effect of a specific type of intervention(s), but assessed an undefined orthodontic intervention, e.g.,
36. AO	2011	treatment, midline position, buccal corridor and smile arc on smile attractiveness. Angle Orthod. 2011 Jan;81(1):153-61. doi: 10.2319/040710-195.1.	orthodontic treatment as a whole
		Papageorgiou SN, Höchli D, Eliades T. Outcomes of comprehensive fixed appliance orthodontic treatment: A systematic review with meta-analysis and methodological overview. Korean J Orthod. 2017 Nov;47(6):401-413. doi:	review did not assess the effect of a specific type of intervention(s), but assessed an undefined orthodontic intervention, e.g., orthodontic treatment as a
37. KJO 38. O&CR	2017	10.4041/kjod.2017.47.6.401. Lee DW, Park JH, Bay RC, Choi SK, Chae JM. Cortical bone thickness and bone density effects on miniscrew success rates: A systematic review and meta-analysis. Orthod Craniofac Res. 2021 Mar;24 Suppl 1:92-102. doi: 10.1111/ocr.12453. Epub 2020 Dec 16.	whole not the effects of orthodontic interventions were assessed
		Lee DW, Park JH, Bay RC, Choi SK, Chae JM. Cortical bone thickness and bone density effects on miniscrew success rates: A systematic review and meta-analysis. Orthod Craniofac Res. 2021 Mar;24 Suppl 1:92-102. doi: 10.1111/ocr.12453. Epub 2020	assessed exclusively
39. O&CR 40. O&CR	2021	Dec 16. Lee DW, Park JH, Bay RC, Choi SK, Chae JM. Cortical bone thickness and bone density effects on miniscrew success rates: A systematic review and meta-analysis. Orthod Craniofac Res. 2021 Mar;24 Suppl 1:92-102. doi: 10.1111/ocr.12453. Epub 2020 Dec 16.	adverse effects not the effects of orthodontic interventions were assessed
41. O&CR	2021	Copello FM, Marañón-Vásquez GA, Brunetto DP, Caldas LD, Masterson D, Maia LC, Sant'Anna EF. Is the buccal alveolar bone less affected by mini- implant assisted rapid palatal expansion than by	assessed exclusively adverse effects

		conventional rapid palatal expansion?-A systematic	
		review and meta-analysis. Orthod Craniofac Res.	
		2020 Aug;23(3):237-249. doi: 10.1111/ocr.12374.	
		Epub 2020 Apr 16. PMID: 32187843.	
		Allen RK, Edelmann AR, Abdulmajeed A, Bencharit	
		S. Salivary protein biomarkers associated with	not the effects of
		orthodontic tooth movement: A systematic review.	orthodontic
		Orthod Craniofac Res. 2019 May;22 Suppl 1:14-20.	interventions
42. O&CR	2019	doi: 10.1111/ocr.12258.	were assessed
		Cannavale R, Chiodini P, Perillo L, Piancino MG.	
		Rapid palatal expansion (RPE): Meta-analysis of	assessed
		long-term effects.Orthod Craniofac Res. 2018	exclusively
43. O&CR	2018	Nov;21(4):225-235. doi: 10.1111/ocr.12244.	adverse effects
			the review is
			about a specific
		Al-Saleh MAQ, Alsufyani N, Flores-Mir C, Nebbe B,	outcome of an
		Major PW. Changes in temporomandibular joint	intervention.
		morphology in class II patients treated with fixed	which is
		mandibular repositioning and evaluated through 3D	ambiguous and
		imaging: a systematic review. Orthod Craniofac Res.	could also be an
44. O&CR	2015	2015 Nov;18(4):185-201. doi: 10.1111/ocr.12099.	adverse effect
	2010	von Bremen J, Ruf S. Orthodontic and dentofacial	
		orthopedic management of juvenile idiopathic	review included
		arthritis: a systematic review of the literature.	orthognathic
		Orthod Craniofac Res. 2011 Aug;14(3):107-15. doi:	surgical
45. O&CR	2011		interventions
45. U&CR	2011	10.1111/j.1601-6343.2011.01514.x.	interventions

*Description of the abbreviated journals:

Cochrane library: Cochrane Database of Systematic Reviews

AJODO: American Journal of Orthodontics and Dentofacial Orthopedics

EJO: European Journal of Orthodontics

AO: Angle Orthodontist

KJO: Korean Journal of Orthodontics

O&CR: Orthodontics and Craniofacial Research

Additional files chapter 4

PRISMA-P 2015 Checklist

Additional file 1. PRISMA-P Checklist

Not an update 2 and 301-306 Not an amendment number(s) Not a systematic review This checklist has been adapted for use with protocol submissions to Systematic Reviews from Table 3 in Moher D et al: Preferred reporting 297-299 297-299 297-299 110-117 61-109 Line N N Information reported Ŷ × × × Yes × 🗆 🗆 × 🔲 × × × × × × Items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Systematic Reviews 2015 4:1 If the protocol represents an amendment of a previously completed or published protocol, identify Provide name, institutional affiliation, and e-mail address of all protocol authors; provide physical If registered, provide the name of the registry (e.g., PROSPERO) and registration number in the Abstract as such and list changes; otherwise, state plan for documenting important protocol amendments Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO) Describe contributions of protocol authors and identify the guarantor of the review If the protocol is for an update of a previous systematic review, identify as such Describe the rationale for the review in the context of what is already known Indicate sources of financial or other support for the review Identify the report as a protocol of a systematic review Provide name for the review funder and/or sponsor mailing address of corresponding author Checklist item **ADMINISTRATIVE INFORMATION** . 2 ₽ 8 ŝ \$ \$ 38 . N 4 ~ Contributions INTRODUCTION Identification Section/topic sponsor/funder Amendments Sources Registration Sponsor Update Contact Role of Objectives Rationale Support Authors 물

Saction/tonic		Charklist Ham	Informatio	Information reported	Line
			Yes	Ŷ	number(s)
METHODS					
Eligibility criteria		Specify the study characteristics (e.g., PICO, study design, setting, time frame) and report characteristics (e.g., years considered, language, publication status) to be used as criteria for eligibility for the review	×		132-135
Information sources		Describe all intended information sources (e.g., electronic databases, contact with study authors, trial registers, or other grey literature sources) with planned dates of coverage	×		137-146
Search strategy	9	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated	×		138-141
STUDY RECORDS					
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	×		149-159
Selection process	115	State the process that will be used for selecting studies (e.g., two independent reviewers) through each phase of the review (i.e., screening, eligibility, and inclusion in meta-analysis)	×		160-170
Data collection process	11c	Describe planned method of extracting data from reports (e.g., piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators	×		171-184
Data items	12	List and define all variables for which data will be sought (e.g., PICO items, funding sources), any pre-planned data assumptions and simplifications	×		186-205
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	×		186-223
Risk of bias in individual studies	2	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis		×	Not applicable
DATA					
	15a	Describe criteria under which study data will be quantitatively synthesized		×	Not a systematic review
Synthesis	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data, and methods of combining data from studies, including any planned exploration of consistency (e.g., 1 ² , Kendall's tau)		×	Not a systematic review
	15c	Describe any proposed additional analyses (e.g., sensitivity or subgroup analyses, meta- regression)		×	Not a systematic review
	_		Informatio	Information reported	-Ine
Section to pic			Yes	Ŷ	number(s)
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned		×	Not a systematic review
Meta-bias(es)	\$	Specify any planned assessment of meta-blas(es) (e.g., publication blas across studies, selective reporting within studies)		×	Not a systematic review
Confidence in cumulative evidence	4	Describe how the strength of the body of evidence will be assessed (e.g., GRADE)		×	Not a systematic review

Additional file 2. Pilot tests

For our pilot studies we used the same sample of 14 reviews that was used for our pilot study of a previous protocol 'Seeking adverse effects in systematic reviews of orthodontic interventions: protocol for a cross-sectional study' [15]. The calculation of the sample size of this pilot study was based on the probability of the Yes scores for the question "Did the review seek any findings related to adverse effects of interventions in the included studies? This sample size was calculated using we the following equation [29]:

$$n = \frac{\ln(1-T)}{\ln(1-\pi)}$$

n = the sample size for the pilot study Υ = the threshold of confidence (95%) π = the probability of a 'Yes' score

Our pilot test on our sample of 14 reviews found that reviewers in 35.7% (5/14) of the abstracts reported or considered (discussed, weighed etc.) potential adverse effects of orthodontic interventions. In this sample we identified a prevalence of 14.3 % (2/14) of spin in the abstract on adverse effects of orthodontic interventions. Both cases of spin were 'Misleading reporting related spin'.

Additional file 3. Search terms and their derivatives

This additional file is identical to additional file 3 of our protocol on seeking adverse in systematic reviews of orthodontic interventions [15].

Search terms and their derivatives	Search terms for searching multiple words
	in a PDF
"adverse"	ADVERSE, Adverse, adverse
"effect", "effects"	EFFECT, Effect, effect
"reaction", "reactions"	REACTION, Reaction, reaction
"complication", "complications", "complicated",	COMPLICAT, Complicat, complicat
"complicating"	
"harm", "harms", "harmful"	HARM, Harm, harm
"risk", "risks", "risky"	RISK, Risk, risk
"safe", "safety"	SAFE, Safe, safe
"side"	SIDE, Side, side
"toxic", "toxicity"	TOXIC, Toxic, toxic
"benefit", "benefits"	BENEFIT, Benefit, benefit
"result", "results"	RESULT, Result, result
"finding", "findings"	FINDING, Finding, finding
"outcome", "outcomes"	OUTCOME, Outcome, outcome
"limitation", "limitations", limit	LIMIT, Limit, limit
"damage", "damages", "damaging"	DAMAGE, Damage, damage
"data"	DATA, Data, data
"information"	INFO, Info, info
"conflict", "conflicts", "conflicting"	CONFLICT, Conflict, conflict
"negative"	NEGATIVE, Negative, negative
"detrimental"	DETRIMENTAL, Detrimental, detrimental
"disadvantage", "disadvantages", "disadvantageous"	DISADVAN, Disadvan, disadvan
"down"	DOWN, Down, down
"injury", "injuries", "injured", "injurious"	INJUR, Injur, injur
"byproduct", "byproducts"	BYPRODUCT, Byproduct, byproduct
"collateral"	COLLATERAL, Collateral, collateral
"unfavorable", "unfavourable"	UNFAVO, Unfavo, unfavo
"destructive"	DESTRUCT, Destruct, destruct
"unsafe"	UNSAFE, Unsafe, unsafe
"undesired", "undesirable"	UNDESIR, Undesir, undesir
"recommend", "recommendation", "recommending"	RECOMMEND, Recommend, recommend
"emergency", "emergencies"	EMERGEN, Emergen, emergen

Table. Search terms and their derivatives

Additional file 4. Data collection forms

Items	Description
Journal	List the pertinent journal
Year	Year of publication
Binder page number	List the binder page number
Reference	List full reference (Authors, Title, Journal)
Is the article a systematic review?	Answer: Yes/No
	Consider definition of a systematic review
What type of systematic review?	List the type of systematic review.
	Consider different types of systematic reviews.
	When the publication is not an intervention
	systematic review describe what type it is or could
	be and classify. Types of systematic reviews will
	receive a final classification during the discussions
	between operators.
Were orthodontic interventions assessed?	Answer: Yes/No
	Consider the definition of orthodontic interventions.
What was the orthodontic intervention?	List the type of orthodontic intervention
	NA: When the article is not a systematic review or
	not a systematic review of interventions.
Is the systematic review eligible?	Answer: Yes/No
	Yes: The article is a systematic review of an
	orthodontic intervention.
	No: The article is not a systematic review of an
	orthodontic intervention.
	No: The article is a systematic review of an
	orthodontic intervention, but focusses exclusively on
	its adverse effects.
Page and potential comments*	Present the pertinent pages of reference for scoring
	the previous items and list the potential comments.

*When referring to a particular page in the systematic review, we will use the page number of the systematic review and not the number in the binder document.

Items	Description
Did the review seek any findings related to adverse	Answer: Yes/No
effects of interventions in the included studies?	Yes: Any findings related to adverse effects of
	interventions in the included studies were sought by
	the reviewers.
	Seeking any findings related to adverse effects of
	interventions in the included studies refers to
	reporting anywhere in the review (except in the
	Abstract) that such adverse effects in the included
	studies were sought.
	Yes: Yes is also scored when reviewers only reported
	findings related to adverse effects of interventions in
	the included studies, but did not report that they
	actually sought them or planned to seek them. For
	example 'Yes' will be scored when outcomes on
	adverse effects of interventions in the included
	studies were reported in the review, but were not
	defined as objectives of the review.
	Yes: Yes is also scored when the reviewers reported
	that they planned to seek (for example in the
	research objectives) findings related to adverse
	effects of interventions in the included studies, but
	did not report on these findings.
	No: Findings related to adverse effects of
	interventions in the included studies were not
	sought by the reviewers.
In abstracts of systematic reviews of orthodontic	Answer: Yes/No
interventions were potential adverse effects of these	Yes: In abstracts of systematic reviews of
interventions reported or considered (i.e., discussed,	orthodontic interventions potential adverse effects
weighed etc.)?	of these interventions were reported or considered
	(i.e., discussed, weighed etc.).
	No: In abstracts of systematic reviews of orthodontic
	interventions potential adverse effects of these
	interventions were not reported or considered (i.e.,
	discussed, weighed etc.).

Table 2. Data collection form on seeking any findings related to adverse effects of interventions in the included studies

Table 3a. Data collection form to identify spin in reviev	vs that did seek adverse effects of interventions
the set of	Cases

Items for misleading reporting (in the abstract) on adverse effects of interventions	Score
1) Not reporting in the abstract on the results of the adverse effects that were reported in the main text of the review.	Yes/no
2) Selective reporting in the abstract on the results of the adverse effects that were reported in the main text of the review.	Yes/no
Summary score on the presence of misleading reporting (in the abstract) on adverse effects of interventions	Yes/no Yes is scored when one or more of the 2 items is answered with a 'Yes'

	No is scored when both items are answered with a 'No'
Items for misleading interpretation (in the abstract) on adverse effects of interventions	
 Claiming in the abstract that the intervention is safe (has no or minimal adverse effects), despite concerning results on the adverse effects in the main text of the review, e.g., based on non-statistically significant results on adverse effects with wide confidence intervals [17]. 	Yes/no
 Downgrading in the abstract the importance of the adverse effects, despite concerning results on the adverse effects in the main text of the review. 	Yes/no
3) Recommendations are made in the abstract for clinical practice that are not congruent with the concerning results on the adverse effects in the main text of the review [17].	Yes/no
Summary score on the presence of misleading interpretation (in the abstract) on adverse effects of interventions	Yes/no Yes is scored when one or more of the 3 items is answered with a 'Yes' No is scored when all 3 items are answered with a 'No'
Items for misleading extrapolation (in the abstract) on adverse effects of interventions	
1) Results are extrapolated in the abstract to another population, intervention, outcome or setting than were assessed in the review despite evidence in the main text on concerning adverse effects on a different population, intervention, outcome or setting.	Yes/no
Summary score on the presence of misleading extrapolation (in the abstract) on adverse effects of interventions	Yes/no Yes is scored when the item is answered with a 'Yes' No is scored when the item is answered with a 'No'

Table 3b. Data collection form to identify spin in reviews that did not seek adverse effects of interventions

Items for misleading reporting (in the abstract) on	Score
adverse effects of interventions	
1) Reporting on results of adverse effects in the	Yes/no
abstract when adverse effects were not sought.	
2) Reporting in the abstract that adverse effects	Yes/no
were sought when they were not sought.	
Summary score on the presence of misleading	Yes/no
reporting (in the abstract) on adverse effects of	Yes is scored when one or more of the 2 items is
interventions	answered with a 'Yes'
	No is scored when both items are answered with a
	'No'
Items for misleading interpretation (in the abstract)	
on adverse effects of interventions	

1) Claiming in the abstract that the intervention is	Yes/no
safe (has no or minimal adverse effects) despite not	
having sought adverse effects.	
2) Downgrading in the abstract the importance of	Yes/no
the adverse effects, despite not having sought	
adverse effects.	
3) Recommendations are made in the abstract for	Yes/no
clinical practice despite not having sought adverse	
effects.	
Summary score on the presence of misleading	Yes/no
interpretation (in the abstract) on adverse effects of	Yes is scored when one or more of the 3 items is
interventions	answered with a 'Yes'
	No is scored when all 3 items are answered with a
	'No'
Items for misleading extrapolation (in the abstract)	
on adverse effects of interventions	
1) Results are extrapolated in the abstract to	Yes/no
another population, intervention, outcome or	
setting than were assessed in the review despite not	
having sought adverse effects.	
Summary score on the presence of misleading	Yes/no
extrapolation (in the abstract) on adverse effects of	Yes is scored when the item is answered with a 'Yes'
interventions	No is scored when the item is answered with a 'No'

Additional files chapter 5

Additional file 1. STROBE Checklist

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item		Page
	No	Recommendation	No
Title and abstract	1	(a) Indicate the study's design with a commonly	1
		used term in the title or the abstract	
		(b) Provide in the abstract an informative and	2
		balanced summary of what was done and what	
		was found	
Introduction			
Background/rationale	2	Explain the scientific background and rationale	3-5
		for the investigation being reported	
Objectives	3	State specific objectives, including any	5
		prespecified hypotheses	
Methods			
Study design	4	Present key elements of study design early in the	6
		paper	
Setting	5	Describe the setting, locations, and relevant	6-9
		dates, including periods of recruitment, exposure,	
		follow-up, and data collection	
Participants	6	(a) Give the eligibility criteria, and the sources	6-8
		and methods of selection of participants	
Variables	7	Clearly define all outcomes, exposures,	9-10
		predictors, potential confounders, and effect	
		modifiers. Give diagnostic criteria, if applicable	
Data sources/	8*	For each variable of interest, give sources of data	10-11
measurement		and details of methods of assessment	
		(measurement). Describe comparability of	
		assessment methods if there is more than one	
		group	
Bias	9	Describe any efforts to address potential sources	NA
		of bias	<u> </u>
Study size	10	Explain how the study size was arrived at	11-12
Quantitative variables	11	Explain how quantitative variables were handled	12
		in the analyses. If applicable, describe which	
		groupings were chosen and why	<u> </u>
Statistical methods	12	(a) Describe all statistical methods, including	11-12
		those used to control for confounding	<u> </u>
		(b) Describe any methods used to examine	11-12
		subgroups and interactions	<u> </u>
		(c) Explain how missing data were addressed	11-12

		(d) If applicable, describe analytical methods taking account of sampling strategy	11-12
		(<u>e</u>) Describe any sensitivity analyses	NA
Results			1
Participants	13*	(a) Report numbers of individuals at each stage of	13-15
	-	study—eg numbers potentially eligible, examined	
		for eligibility, confirmed eligible, included in the	
		study, completing follow-up, and analysed	
		(b) Give reasons for non-participation at each	13-15
		stage	
		(c) Consider use of a flow diagram	13
Descriptive data	14*	(a) Give characteristics of study participants (eg	13-15
		demographic, clinical, social) and information on	
		exposures and potential confounders	
		(b) Indicate number of participants with missing	13-15
		data for each variable of interest	
Outcome data	15*	Report numbers of outcome events or summary	13-15
		measures	
Main results	16	(a) Give unadjusted estimates and, if applicable,	13-15
		confounder-adjusted estimates and their	
		precision (eg, 95% confidence interval). Make	
		clear which confounders were adjusted for and	
		why they were included	
		(b) Report category boundaries when continuous	13-15
		variables were categorized	
		(c) If relevant, consider translating estimates of	NA
		relative risk into absolute risk for a meaningful	
		time period	
Other analyses	17	Report other analyses done—eg analyses of	NA
		subgroups and interactions, and sensitivity	
		analyses	
Discussion			
Key results	18	Summarise key results with reference to study	15
		objectives	
Limitations	19	Discuss limitations of the study, taking into	17
		account sources of potential bias or imprecision.	and
		Discuss both direction and magnitude of any	18
		potential bias	
Interpretation	20	Give a cautious overall interpretation of results	15-18
		considering objectives, limitations, multiplicity of	
		analyses, results from similar studies, and other	
		relevant evidence	
Generalisability	21	Discuss the generalisability (external validity) of	18
		the study results	
Other information			

Funding	22	Give the source of funding and the role of the	NA
		funders for the present study and, if applicable,	
		for the original study on which the present article	
		is based	

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.

Additional file 2

Additional file item	Description	
Additional file 2A	Differences between the protocol and the completed cross-sectional	
	study	
Additional file 2B	Search terms and their derivatives	
Additional file 2C	Data collection forms	
Additional file 2D	Adverse effects hypothetically linked to orthodontic interventions	
References	References for additional file 2	

Table of contents for additional file 2

Additional file 2A. Differences between the protocol and the completed cross-sectional study

Differences between the protocol and the	Rationale
completed cross-sectional study	
Fine-tuning of the definitions of the 3 types of spin	In our protocol we wanted to identify spin by
	comparing what was reported in the abstract with
	what was reported in the main text of the
	manuscript. We changed this to comparing what was
	reported in the abstract with what was found in the
	review. This fine-tuning was necessary, because on a
	few occasions what was reported in the main text of
	the review was not completely congruent with the
	findings of the review or pertinent information was
	not reported in the main text of the review, but only
	in supplementary files.
Explorative analyses to assess the presence of spin in	We assessed the association between the presence
the abstract and a series of predictors	of spin in the abstract and a series of predictors.
	These predictors were not defined a priori in our
	protocol, but several were explored in recent studies
	on spin in the field of orthodontics (Guo 2021,
	Makou 2021). We therefore conducted explorative
	analyses to determine associations between the
	presence of spin in the abstract and various
	predictors.

Additional file 2B. Search terms and their derivatives
Table. Search terms and their derivatives

Search terms and their derivatives	Search terms for searching multiple words	
	in a PDF	
"adverse"	ADVERSE, Adverse, adverse	
"effect", "effects"	EFFECT, Effect, effect	
"reaction", "reactions"	REACTION, Reaction, reaction	
"complication", "complications", "complicated",	COMPLICAT, Complicat, complicat	
"complicating"		
"harm", "harms", "harmful"	HARM, Harm, harm	
"risk", "risks", "risky"	RISK, Risk, risk	
"safe", "safety"	SAFE, Safe, safe	
"side"	SIDE, Side, side	
"toxic", "toxicity"	TOXIC, Toxic, toxic	
"benefit", "benefits"	BENEFIT, Benefit, benefit	
"result", "results"	RESULT, Result, result	
"finding", "findings"	FINDING, Finding, finding	
"outcome", "outcomes"	OUTCOME, Outcome, outcome	
"limitation", "limitations", limit	LIMIT, Limit, limit	
"damage", "damages", "damaging"	DAMAGE, Damage, damage	
"data"	DATA, Data, data	
"information"	INFO, Info, info	
"conflict", "conflicts", "conflicting"	CONFLICT, Conflict, conflict	
"negative"	NEGATIVE, Negative, negative	
"detrimental"	DETRIMENTAL, Detrimental, detrimental	
"disadvantage", "disadvantages", "disadvantageous"	DISADVAN, Disadvan, disadvan	
"down"	DOWN, Down, down	
"injury", "injuries", "injured", "injurious"	INJUR, Injur, injur	
"byproduct", "byproducts"	BYPRODUCT, Byproduct, byproduct	
"collateral"	COLLATERAL, Collateral, collateral	
"unfavorable", "unfavourable"	UNFAVO, Unfavo, unfavo	
"destructive"	DESTRUCT, Destruct, destruct	
"unsafe"	UNSAFE, Unsafe, unsafe	
"undesired", "undesirable"	UNDESIR, Undesir, undesir	
"recommend", "recommendation", "recommending"	RECOMMEND, Recommend, recommend	
"emergency", "emergencies"	EMERGEN, Emergen, emergen	

Additional file 2C. Data collection forms

Items	Description
Did the review seek any findings related to adverse	Answer: Yes/No
effects of interventions in the included studies?	Yes: Any findings related to adverse effects of
	interventions in the included studies were sought by the
	reviewers.
	Seeking any findings related to adverse effects of
	interventions in the included studies refers to reporting
	anywhere in the review (except in the Abstract) that
	such adverse effects in the included studies were sought.
	Yes: Yes is also scored when reviewers only reported
	findings related to adverse effects of interventions in the
	included studies, but did not report that they actually
	sought them or planned to seek them. For example 'Yes'
	will be scored when outcomes on adverse effects of
	interventions in the included studies were reported in
	the review, but were not defined as objectives of the
	review.
	Yes: Yes is also scored when the reviewers reported that
	they planned to seek (for example in the research
	objectives) findings related to adverse effects of
	interventions in the included studies, but did not report on these findings.
	No: Findings related to adverse effects of interventions
	in the included studies were not sought by the
	reviewers.
In abstracts of systematic reviews of orthodontic	Answer: Yes/No
interventions were potential adverse effects of these	Yes: In abstracts of systematic reviews of orthodontic
interventions were potential adverse eneces of these interventions reported or considered (i.e., discussed,	interventions potential adverse effects of these
weighed etc.)?	interventions were reported or considered (i.e.,
- 5	discussed, weighed etc.).
	No: In abstracts of systematic reviews of orthodontic
	interventions potential adverse effects of these
	interventions were not reported or considered (i.e.,
	discussed, weighed etc.).

Data collection form on reporting of adverse effects in the abstract of systematic reviews of orthodontic interventions

Data collection forms to identify spin of adverse effects of orthodontic interventions in abstracts of systematic reviews

To assign spin of adverse effects of orthodontic interventions in abstracts of systematic reviews we developed separate checklists for reviews that sought adverse effects of interventions and those that did not. These checklists were pilot-tested a priori during the development of our protocol. Because only 2 systematic reviews in this pilot study reported on adverse effects of interventions, we decided to conduct an additional pilot test to further fine-tune these checklists. Ten randomized controlled trials (RCTs) (2 per journal) of orthodontic interventions were selected consecutively from the websites of the 5 leading orthodontic journals from June 2021 backwards. RCTs were eligible for the pilot study if they assessed adverse effects of orthodontic interventions. The first 2 RCTs for each journal that were identified during this search process were selected. The fine-tuned checklists are reported in the tables under here and differ slightly from those reported in our protocol (Steegmans 2019). Changes were made to (1) reduce inter-operator differences in assigning various types of spin and (2) make the descriptions of spin more congruent with those given in the literature (Boutron 2018, Haneef 2017, Lazarus 2015) and with the definition of spin (in the abstract) on adverse effects of interventions presented in our protocol (Steegmans 2019), i.e., 'Incomplete or inadequate reporting, interpretation, or extrapolation (or a combination of these variables) of findings on adverse effects of interventions in the abstract that could be misleading for the reader' (See additional file 2A).

Items for misleading reporting (in the abstract) on	Score
adverse effects of interventions	
1) Not reporting in the abstract on the results of adverse effects found in the review.	Yes/no
2) Selective reporting in the abstract on the results of adverse effects found in the review.	Yes/no
Summary score on the presence of misleading reporting (in the abstract) on adverse effects of interventions	Yes/no Yes is scored when one or more of the 2 items is answered with a 'Yes' No is scored when both items are answered with a 'No'
Items for misleading interpretation (in the abstract) on adverse effects of interventions	
1) Claiming in the abstract that the intervention is safe (has no or minimal adverse effects), despite concerning results on adverse effects found in the review e.g., based on non-statistically significant results on adverse effects with wide confidence intervals (Yavchitz 2016)	Yes/no
 Downgrading in the abstract the importance of the adverse effects, despite concerning results on adverse effects found in the review. 	Yes/no
3) Recommendations are made in the abstract for clinical practice that are not supported by the findings in the review on adverse effects'(Yavchitz 2016)	Yes/no
Summary score on the presence of misleading interpretation (in the abstract) on adverse effects of interventions	Yes/no Yes is scored when one or more of the 3 items is answered with a 'Yes' No is scored when all 3 items are answered with a 'No'
Items for misleading extrapolation (in the abstract) on adverse effects of interventions	
1) Results are extrapolated in the abstract to another population, intervention, outcome or setting than were assessed in the review despite evidence on adverse effects on a different population, intervention, outcome or setting.	Yes/no
Summary score on the presence of misleading extrapolation (in the abstract) on adverse effects of interventions	Yes/no Yes is scored when the item is answered with a 'Yes' No is scored when the item is answered with a 'No'

Data collection form to identify spin in abstracts of reviews that did seek adverse effects of interventions

Data collection forms to identify spin in abstracts of reviews that did not seek adverse effects of interventions

interventions	- Coore
Items for misleading reporting (in the abstract) on	Score
adverse effects of interventions	
1) Reporting on results of adverse effects in the	Yes/no
abstract when adverse effects were not sought.	
Reporting in the abstract that adverse effects	Yes/no
were sought when they were not sought.	
Summary score on the presence of misleading	Yes/no
reporting (in the abstract) on adverse effects of	Yes is scored when one or more of the 2 items is
interventions	answered with a 'Yes'
	No is scored when both items are answered with a
	'No'
Items for misleading interpretation (in the abstract)	
on adverse effects of interventions	
1) Claiming in the abstract that the intervention is	Yes/no
safe (has no or minimal adverse effects) despite not	
having sought adverse effects.	
2) Downgrading in the abstract the importance of	Yes/no
the adverse effects, despite not having sought	
adverse effects.	
3) Recommendations are made in the abstract for	Yes/no
clinical practice despite not having sought adverse	
effects.	
Summary score on the presence of misleading	Yes/no
interpretation (in the abstract) on adverse effects of	Yes is scored when one or more of the 3 items is
interventions	answered with a 'Yes'
	No is scored when all 3 items are answered with a
	'No'
Items for misleading extrapolation (in the abstract)	
on adverse effects of interventions	
 Results are extrapolated in the abstract to 	Yes/no
another population, intervention, outcome or	
setting than were assessed in the review despite not	
having sought adverse effects.	
Summary score on the presence of misleading	Yes/no
extrapolation (in the abstract) on adverse effects of	Yes is scored when the item is answered with a 'Yes'
interventions	No is scored when the item is answered with a 'No'
	NO IS SCOLED WHEN THE REIT IS DISWELED WITH U NO

Adverse effects related to	Description		
Tooth structures	Tooth crown		
	 decalcifications, decays, tooth wear, enamel cracks and fractures; discolorations, deterioration of prosthetic crown (as fracturing a ceramic one during debonding); 		
	 iatrogenic damage to the crown, e.g., fracture as a result of trauma Tooth root 		
	 root resorption, early closure of root apex, ankylosis; 		
	 iatrogenic damage to the root, e.g., fracture as a result of trauma 		
	Tooth pulp		
	ischemia, pulpitis, necrosis		
	 iatrogenic damage to the pulp, e.g., fracture as a result of trauma 		
Periodontal tissues	 gingivitis, periodontitis, gingival recession or hypertrophy, alveolar bone loss, dehiscences, fenestrations, interdental fold, dark triangles; tooth mobility, plague retention, bacterial count 		
Intraoral (non-tooth or periodontal) tissues	 intraoral tissue irritations and inflammation such as mucosal ulcerations or hyperplasia or irritations of the tongue (as a result of trauma by appliances, e.g., breakage, failure, loosening etc. of appliances or long arch wires) 		
	 Scar formation after suturing 		
	 chemical burns (e.g., etching related) 		
	 thermal injuries (e.g., overheated burs) 		
	nerve damage		
	 tooth eruption, i.e., eruption disturbances (e.g., impactions) caused by orthodontic appliances 		
Extraoral tissues (non-	 cutting of lips or cheeks, eye injury (e.g., as a result of trauma by 		
temporomandibular tissues)	appliances, e.g., breakage, failure, loosening etc. of appliances or long		
	arch wires or headgear-related trauma)		
Temporomandibular tissues and	discomfort on the lip temporomandibular tissues and disorders		
disorders	 temporomandibular tissues and disorders 		
Appliance failure	 breakage, failure, loosening etc. of appliances 		
	 long archwires, headgear-related trauma 		
Undesired treatment results	 inadequate morpho-functional, aesthetic or functional final result 		
	 inaccuracy of the treatment result 		
	 non predictability of the treatment result Dental side affects a supported time of teach, and another set 		
	 Dental side effects e.g., unwanted tipping of teeth, anchorage loss etc. Skeletal side effects, e.g., unwanted backward rotation of the mandible 		
Relapse and stability	Relapse and stability of the obtained treatment result		
Undesired qualitative	Pain and discomfort		
experiences by the patient or	 orthodontic tooth movement-related pain and discomfort 		
carer(s)	 appliance (intervention)-related pain and discomfort: i.e., pain and discomfort as a result of the appliance (intervention) itself with or without pain and discomfort associated with tooth movement e.g., tension or pressure of the appliances (constriction of appliances), 		
	speech difficulties, eating difficulties, swallowing difficulties, food accumulation, bad tastes and smells		
	 additional intervention-related pain and discomfort, e.g., surgical and non-surgical adjunctive interventions to accelerate tooth movement 		
	 Tolerability/acceptance/stress issues with the treatment procedures Absence from work or studies and difficulties in daily activities 		
	 collaboration (compliance) issues or failure to complete treatment, e.g., dropout 		

Additional file 2D. Adverse effects hypothetically linked to orthodontic interventions (Steegmans 2023a)

	patient anxiety	
	being teased	
	social discomfort	
	 embarrassment to wear the appliance 	
	 behavioral changes of patients and parents, impaired family 	
	relationships	
	 aesthetic look discontents during orthodontic appliance usage 	
	concentration difficulties	
	 reduced enjoyment of food and change in taste 	
	 sleeping difficulties 	
	 removal of appliance during sleep 	
	development of mannerisms	
	Satisfaction with the treatment procedures and final result not satisfied with the treatment procedures (Check in text what was 	
	measured, i.e., during or after)	
	 not satisfied with the final treatment result (Check in text what was 	
	measured, i.e., during or after)	
Gastro-intestinal	 accidental swallowing of small parts of the orthodontic device (tubes, 	
	brackets);	
Allergy	Allergies to nickel or latex;	
Cardio	 infective endocarditis; 	
Chronic fatigue		
Cross infections	 from doctor to patient, patient to doctor, patient to patient. 	
Non-defined	Adverse effects that were not defined by the authors of the review: referring to	
	'any adverse effect', 'any side effect' etc.	
Additional adverse effects	Additional adverse effects that were identified during data extraction that could	
	not be labeled under any of the categories of adverse effects given in this table	

*Modified from Preoteasa et al. (Preoteasa 2012)

References for additional file 2

Boutron 2018

Boutron J, Ravaud P. Misrepresentation and distortion of research in biomedical literature. Proc Natl Acad Sci U <u>S A.</u> 2018 Mar 13;115(11):2613-2619. doi: 10.1073/pnas.1710755115.

Guo 2021

Guo F, Fang X, Li C, Qin D, Hua F, He H. The presence and characteristics of 'spin' among randomized controlled trial abstracts in orthodontics. Eur J Orthod. 2021 Oct 4;43(5):576-582. doi: 10.1093/ejo/cjab044. PMID: 34397084.

Haneef 2017

Haneef R, Yavchitz A, Ravaud P, Baron G, Oranksy I, Schwitzer G, Boutron I. <u>Interpretation of health</u> <u>news items reported with or without spin: protocol for a prospective meta-analysis of 16 randomised</u> <u>controlled trials</u>. BMJ Open. 2017 Nov 17;7(11):e017425. doi: 10.1136/bmjopen-2017-017425.

Lazarus 2015

Lazarus C, Haneef R, Ravaud P, Boutron I. Classification and prevalence of spin in abstracts of non-randomized studies evaluating an intervention. <u>BMC Med Res Methodol.</u> 2015 Oct 13;15:85. doi: 10.1186/s12874-015-0079-x.

Lefebvre 2021

Lefebvre C, Glanville J, Briscoe S, Littlewood A, Marshall C, Metzendorf M-I, Noel-Storr A, Rader T, Shokraneh F, Thomas J, Wieland LS. Chapter 4: Searching for and selecting studies. In: Higgins JPT, Thomas J, Chandler J,

Cumpston M, Li T, Page MJ, Welch VA (editors). *Cochrane Handbook for Systematic Reviews of Interventions* version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Makou 2021

Makou O, Eliades T, Koletsi D. Reporting, interpretation, and extrapolation issues (SPIN) in abstracts of orthodontic meta-analyses published from 2000 to 2020. Eur J Orthod. 2021 Mar 19:567-575. doi: 10.1093/ejo/cjab009. Epub ahead of print. PMID: 33740054.

Preoteasa 2012

Preoteasa CT, Ionescu E, Preoteasa E. Chapter 18: Risks and complications associated with orthodontic treatment. In: Bourzgui F. (editor). Orthodontics-Basic aspects and clinical considerations. March 9, 2012 under CC BY 3.0 license. www.intechopen.com. [online] Available from: <u>https://cdn.intechopen.com/pdfs/31388/InTech-</u>

Risks and complicationsassociated with orthodontic treatment.pdf (accessed December 4th 2021).

Steegmans 2019

Steegmans, P.A.J., Di Girolamo, N, Meursinge Reynders, R.A. Spin in the reporting, interpretation, and extrapolation of adverse effects of orthodontic interventions: protocol for a cross-sectional study of systematic reviews. *Res Integr Peer Rev* **4**, 27 (2019). <u>https://doi.org/10.1186/s41073-019-0084-4</u>

Steegmans 2023a

Steegmans PAJ, Di Girolamo N, Bipat S, Reynders RAM. Seeking adverse effects in systematic reviews of orthodontic interventions: a cross-sectional study (part 1). Syst Rev. 2023 Jul 3;12(1):112.

Yavchitz 2016


Yavchitz A, Ravaud P, Altman DG, Moher D, Hrobjartsson A, Lasserson T, Boutron I. A new classification of spin in systematic reviews and meta-analyses was developed and ranked according to the severity. J Clin Epidemiol. 2016 Jul;75:56-65. doi: 10.1016/j.jclinepi.2016.01.020. Epub 2016 Feb 2.

Additional file 3

Table of contents	s for additional file 3

Additional file item	Description
Additional file 3A	PRISMA flow diagram
Additional file 3B	Included reviews
Additional file 3C	Excluded studies with rationale

Additional file 3A. PRISMA flow diagram



*CDSR: Cochrane Database of Systematic Reviews, EJO: European Journal of Orthodontics, AJODO: American Journal of Orthodontics and Dentofacial Orthopedics, AO: Angle Orthodontist, KJO: Korean Journal of Orthodontics, O&CR: Orthodontics and Craniofacial Research ** The rationale for exclusion of each report are given in Additional file 3C

Additional file 3B. Included reviews

This additional file is identical to additional file 3 of chapter 3.

Additional file 3C. Excluded studies with rationale


This additional file is identical to additional file 4 of chapter 3.

Topic	Item	Checklist item description	Reported on Line
Title	-	The diagnosis or intervention of primary focus followed by the words "case report"	PI
Key Words	~	2 to 5 key words that identify diagnoses or interventions in this case report, including "case report" _	ы
Abstract	an se	Introduction: What is unique about this case and what does it add to the scientific literature?	Ы
(no references)	36	Main symptoms and/or important clinical findings	ы
	30	The main diagnoses, therapeutic interventions, and outcomes	PI
	3d	Conclusion—What is the main 'take-away' lesson(s) from this case?	PI
Introduction	4	One or two paragraphs summarizing why this case is unique (may include references)	P1-2
Patient Information	Sa	De-identified patient specific information	P2-3
	8	Primary concerns and symptoms of the patient	B3
	50	Medical, family, and psycho-social history including relevant genetic information	P3
	56	Relevant past interventions with outcomes	P3
Clinical Findings	9	Describe significant physical examination (PE) and important clinical findings	P4
Timeline	~	Historical and current information from this episode of care organized as a timeline	P4
Diagnostic	8a	Diagnostic testing (such as PE, laboratory testing, imaging, surveys).	P5-6
Vasessment	48	Diagnostic challenges (such as access to testing, financial, or cultural)	P5-6
	80	Diagnosis (including other diagnoses considered)	8
	98	Prognosis (such as staging in oncology) where applicable	B6
Intervention	9a	Types of therapeutic intervention (such as pharmacologic, surgical, preventive, self care)	P6-7
	8	Administration of therapeutic intervention (such as dosage, strength, duration)	P6-7
Follow-up and	90	Changes in therapeutic intervention (with rationale)	P6-7
Outcomes	10a	Cinician and patient-assessed outcomes (if available)	P8
	106	Important follow-up diagnostic and other test results	84
	10c	Intervention adherence and tolerability (How was this assessed?)	64
	104	Adverse and unanticipated events	64
Discussion	11a	A scientific discussion of the strengths AND limitations associated with this case report	P10
	116	Discussion of the relevant medical literature with references	P10
	11c	The scientific rationale for any condusions (including assessment of possible causes)	P11
	11d	The primary "take-away" lessons of this case report (without references) in a one paragraph conclusion	P11
Patient Perspective	12	The patient should share their perspective in one to two paragraphs on the treatment(s) they received	B
			and on the second

Additional files chapter 8

Additional file 1. CARE Checklist

Additional file 2



Supplementary file 2. Timeline

13

Additional file 3

Orthodontie



Gustav Mahlerlaan 3004, 1081 LA Amsterdam

Tel: 020-5980610 / 5980611

Patiënten enquête Esas 2019 2020

datum: 22 / 01 / 2020

UW geboorte datum:

NAAM van uw behandelaar:

geslacht M/V

ESASnummer:

https://esas.nu/

Geachte ouders en patiënt(e),

Wij doen ons best om op een prettige manier goede orthodontische zorg te verlenen, rekening houdend met de wensen en de verwachtingen van onze patiënten. Om na te gaan of dat lukt, worden zowel patiënten als ouders verzocht hun mening over onze afdeling te geven, via het invullen van deze enquête. Met de resultaten hiervan kunnen wij werken aan verbetering van onze zorgverlening. U hoeft uw naam niet te vermelden, want het onderzoek is gegarandeerd anoniem. Wij stellen het bijzonder op prijs als ook u hieraan wilt meewerken, het duurt slecht enkele minuten. Bij voorbaat hartelijk dank voor uw moeite en tijd om onze praktijkorganisatie mede door uw input te verbeteren.

A	Over de orthodontist	nie	t wa	ar (+	wel	l waa	ır			
	1. Is vriendelijk en goed aanspreekbaar	c	c	с	c	0	ĸ				
	2. Houdt me goed op de hoogte van het verloop van de behandeling	c	c	c	c	\$	×				
	3. Heeft mijn volledige vertrouwen	C	c	C	c	9	τ				
	4. Geeft mij alle informatie die ik nodig heb	c	c	c	c	9	R				
	5. Neemt de tijd voor me	c	c	c	c	6	ĸ				
	6. Luistert naar wat ik te zeggen heb	c	c	c	c	ð	ĸ				
	7. Is helder en duidelijk	c	r	c	c	0	ĸ				
	8. Gaat goed om met mensen	c	c	c	c	0	ĸ				
	9. Werkt netjes op tijd	c	c	c	c	0	×				
	10. Is goed in wat hij/zij doet	c	c	c	c	9	z				
в	Over de praktijkmedewerk(st) ers	niet waar ← → wel waar									
	11. Zijn vriendelijk				-	c	c	c	R		
	12. Zijn zorgzaam en plezierig				-	c	c	c	8		

Orthodontie

	13. Vormen een goed team	c	c	c	c	×
	14. Zijn goed in wat zij doen	c	c	c	c	æ
	15. De telefoon wordt prompt beantwoord	c	c	c	c	0
	16. Afspraken maken gaat gemakkelijk en snel	c	c	c	c	0
c	Over de praktijk					
	17. De praktijk ziet er goed uit	C	c	C	c	×
	18. De wachtkamer is aangenaam	c	C	x	c	c
	19. Er zijn altijd voldoende nieuwe tijdschriften aanwezig	C	c	R	c	с .
	20. De praktijk is goed bereikbaar	C	c	c	×	0
>	Over de behandeling	ni	et w	aar 📢		wel waar
	21. Ik heb nooit iets echt vervelends ervaren	C	C	C	C	× ·
	22. Het eindresultaat is zoals ik had verwacht	c	c	C	C	×
	23. De behandeling duurde net zo lang als was voorspeld	c	C	r	*	C C
	24. Mijn tanden zijn heel mooi geworden	c	c	c	c	×
Ę	Tenslotte					
	25. Ik zal deze praktijk aan mijn vrienden aanbevelen	C	C	C	X	

Supplementary file 3. ESAS patient satisfaction evaluation part

